

THE COMPLETE


Nursing School BUNDLE™

BROUGHT TO YOU BY





 www.etsy.com/shop/nurseinthemaking

 [@Kristine_nurseinthemaking](https://www.instagram.com/Kristine_nurseinthemaking)

 [@NurseInTheMaking](https://www.youtube.com/@NurseInTheMaking)

 [@nurseinthemakingkristine](https://www.tiktok.com/@nurseinthemakingkristine)

 [@anurseinthemaking](https://www.facebook.com/anurseinthemaking)





















 Kristine@anurseinthemaking.com




By purchasing this material, you agree to the following terms and conditions: you agree that this ebook and all other media produced by NurseInTheMaking LLC are simply guides and should not be used over and above your course material and teacher instruction in nursing school. When details contained within these guides and other media differ, you will defer to your nursing school's faculty/staff instruction. Hospitals and universities may differ on lab values; you will defer to your hospital or nursing school's faculty/staff instruction. These guides and other media created by NurseInTheMaking LLC are not intended to be used as medical advice or clinical practice; they are for educational use only. You also agree to not distribute or share these materials under any circumstances; they are for personal use only.

© 2021 NurseInTheMaking LLC. All content is property of NurseInTheMaking LLC and www.anurseinthemaking.com. Replication and distribution of this material is prohibited by law. All digital products (PDF files, ebooks, resources, and all online content) are subject to copyright protection. **Each product sold is licensed to an individual user and customers are not allowed to distribute, copy, share, or transfer the products to any other individual or entity, they are for personal use only. Fines of up to \$10,000 may apply and individuals will be reported to the BRN and their school of nursing.**

TABLE OF CONTENTS

	Head-To-Toe Assessment	5
	Dosage Calculation	9
	Lab Value Cheat Sheet with Memory Tricks	21
	Electrolyte Imbalances	25
	Fundamentals.....	31
	Mental Health.....	49
	Mother Baby	59
	Pediatrics.....	77
	Med-Surg	
	Renal / Urinary System	100
	Cardiac System	108
	Endocrine System	128
	Respiratory Disorders	136
	Hematology Disorders	140
	Gastrointestinal Disorders	144
	Neurological Disorders.....	148
	Critical Care (Burns & Shock)	153
	ABGs	159
	Musculoskeletal	163
	Pharmacology	167
	Templates & Planners	199
	Note from Kristine	209



HEAD-TO-TOE ASSESSMENT

BROUGHT TO YOU BY



HEAD-TO-TOE ASSESSMENT

- 1 INSPECT
- 2 PALPATE
- 3 PERCUSS
- 4 AUSCULTATE

Introduction

- * Knock
- * Introduce yourself
- * Wash hands
- * Provide privacy
- * Verify client ID and DOB
- * Explain what you are doing (using non-medical language)

Orientation

- * What is your name?
- * Do you know where you are?
- * Do you know what month it is?
- * Who is the current U.S. president?
- * What are you doing here?
- * **A&O X4** = Oriented to Person, Place, Time, and Situation


"Normal" Vital Signs

- * **PULSE:** 60-100 bpm
- * **BLOOD PRESSURE:** 120/80 mmHg
- * **O₂ SATURATION:** 95-100%
- * **TEMPERATURE:** 97.8-99.1°F
- * **RESPIRATIONS:** 12-20 breaths per min

Head & Face

HEAD

- * Inspect head/scalp/hair
- * Palpate head/scalp/hair



VII: FACIAL

- Raise eyebrows
- Smile
- Frown
- Show teeth
- Puff out cheeks
- Tightly close eyes

FACE

- * Inspect
- * Check for symmetry
 - * To assess **CRANIAL NERVE 7**, check....

EYES

- * Inspects external eye structures
- * Inspect color of conjunctiva and sclera
- * **PERRLA**
 - Pupils **E**qual, **R**ound, **R**eactive to **L**ight, & **A**ccommodation

PULSE SCALE


0	PULSE IS ABSENT
1+	DIMINISHED
2+	NORMAL
3+	FULL
4+	BOUNDED, STRONG

Assessing the strength of the pulse

Neck, (Chest (Lungs) & Heart


NECK

- * Inspect and palpate
- * Palpate carotid pulse
- * Check skin turgor (under clavicle)



POSTERIOR CHEST

- * Inspect
- * Auscultate lung sounds in posterior and lateral chest
 - Note any crackles or diminished breath sounds



ANTERIOR CHEST

- * Inspect:
 - Use of accessory muscles
 - AP to transverse diameter
 - Sternum configuration
- * Palpate: symmetric expansion
- * Auscultate lung sounds → anterior and lateral
 - Note any crackles or diminished breath sounds

ASSESS THE DEPTH OF THE RESPIRATIONS

EFFORT	note if it's LABORED or UNLABORED
RHYTHM	note if it's REGULAR or IRREGULAR

5 Areas for Listening to the Heart

ALL PEOPLE ENJOY TIME MAGAZINE



- Aortic**
- Pulmonic**
- Erb's Point**
- Tricuspid**
- Mitral**

HEART

- * Auscultate heart sounds (**A, P, E, T, M**) with diaphragm and bell
 - Note any murmurs, whooshing, bruits, or muffled heart sounds

HEAD-TO-TOE ASSESSMENT

Peripherals

UPPER EXTREMITIES

- ✦ Inspect and palpate
- ✦ Note any texture, lesions, temperature, moisture, tenderness, & swelling
- ✦ Palpate radial pulses bilaterally

0	PULSE IS ABSENT
1+	DIMINISHED
2+	NORMAL
3+	FULL
4+	BOUNGING, STRONG

SHOULDER

- ✦ Inspect, palpate, and assess

ELBOWS

- ✦ Inspect, palpate, and assess

HANDS AND FINGERS

- ✦ Inspect hands/fingers/nails
- ✦ Palpate hands and finger joints
- ✦ Check muscle strength of hands bilaterally
 - Does each hand grip evenly?

Spine

- ✦ Have the client stand up (if able)
- ✦ Inspect the skin on the back
- ✦ Inspect: spinal curvature (cervical/thoracic/lumbar)
- ✦ Palpate spine
- ✦ Note any lesions, lumps, or abnormalities

If we were to percuss + palpate before listening (auscultating), we would alter the bowel sounds. This would lead to inaccurate results.

Lower Extremities (hips, knees, ankles)

LOWER EXTREMITIES

- ✦ Inspect:
 - Overall skin coloration
 - Lesions
 - Hair distribution
 - Varicosities
 - Edema
- ✦ Palpate: Check for edema (pitting or non-pitting)
- ✦ Check capillary refill bilaterally

HIPS

- ✦ Inspect and palpate

KNEES

- ✦ Inspect and palpate

ANKLES

- ✦ Inspect and palpate
- ✦ Posterior pulse
- ✦ Dorsal pedis pulse bilaterally
 - Check strength bilaterally
 - Dorsiflexion flexion against resistance

0	PULSE IS ABSENT
1+	DIMINISHED
2+	NORMAL
3+	FULL
4+	BOUNGING, STRONG

CAPILLARY REFILL TIME (CRT)
Time taken for capillary bed to regain its color after pressure has been applied

NORMAL < 2-3 SECONDS

Abdomen

- ✦ Inspect:
 - Skin color
 - Contour
 - Scars
 - Aortic pulsations
- ✦ Auscultate bowel sounds: all 4 quadrants (start in RLQ and go clockwise)
- ✦ Light palpation: all 4 quadrants

Assess in different order:

- 1 INSPECT**
- 2 AUSCULTATE**
- 3 PERCUSS**
- 4 PALPATE**

ABSENT: Must listen for at least 5 minutes to chart absent bowel sounds

HYPOACTIONIVE: One bowel sound every 3-5 minutes

NORMOACTIONIVE: Gurgles 5-30 times per minute

HYPERACTIONIVE: Can sometimes be heard without a stethoscope. Constant bowel sounds (> 30 sounds per minute)

OVERALL

- ✦ Positions and drapes client appropriately during exam (gave client privacy)
- ✦ Gave client feedback/instructions
- ✦ Exhibits professional manner during exam, treated client with respect and dignity
- ✦ Organized: exam followed a logical sequence (order of exam "made sense")



DOSAGE CALCULATION

BROUGHT TO YOU BY



ABBREVIATIONS

TIMES OF MEDICATIONS

ac	before meals
pc	after meals
daily	every day
bid	two times a day
tid	three times a day
qid	four times a day
qh	every hour
ad lib	as desired
stat	immediately
q2h	every 2 hours
q4h	every 4 hours
q6h	every 6 hours
prn	as needed
hs	at bedtime

EXAMPLE

QUESTION: A patient is receiving 1 mg tid. How many mg will they receive in one day?

Remember: tid = 3X a day

ANSWER: If they are receiving 1 mg for 3X a day, that's $1 \text{ mg} \times 3 = 3 \text{ mg}$ per day

ROUTES OF ADMINISTRATION

PO	by mouth
IM	intramuscularly
PR	per rectum
SubQ	subcutaneously
SL	sublingual
ID	intra dermal
GT	gastrostomy tube
IV	intravenous
IVP	intravenous push
IVPB	intravenous piggyback
NG	nasogastric tube

DRUG PREPARATION

tab, tabs	tablet
cap, caps	capsule
gtt	drop
EC	enteric coated
CR	controlled release
susp	suspension
el, elix	elixir
sup, supp	suppository
SR	sustained release

METRIC

g (gm, Gm)	gram
mg	milligram
mcg	microgram
kg (Kg)	kilogram
L	liter
mL	milliliter
mEq	milliequivalent

APOTHECARY & HOUSEHOLD

gtt	drop
min, m, mx	minim
tsp	teaspoon
pt	pint
gal	gallon
dr	dram
oz	ounce
T, tbs, tbsp	tablespoon
qt	quart

CONVERSIONS

BASED ON VOLUME

$$1 \text{ mg} = 1,000 \text{ mcg}$$

$$1 \text{ g} = 1,000 \text{ mg}$$

$$1 \text{ oz} = 30 \text{ mL}$$

$$8 \text{ oz} = 1 \text{ cup}$$

$$1 \text{ tsp} = 5 \text{ mL}$$

$$1 \text{ dram} = 5 \text{ mL}$$

$$1 \text{ tbsp} = 15 \text{ mL}$$

$$1 \text{ tbsp} = 3 \text{ tsp}$$

$$1 \text{ L} = 1,000 \text{ mL}$$

$$1 \text{ mL} = 5 \text{ gtts (drops)}$$

THE METRIC SYSTEM

LARGE unit to **SMALL** unit → move decimal to the **RIGHT**

SMALL unit to **LARGE** unit → move decimal to the **LEFT**

MOVING TO A LARGER UNIT?

Move the decimal place to the **Left**
(Ex: mcg → mg)



Larger unit
think **L**eft

EXAMPLE

$$1500 \text{ mcg} = \underline{\hspace{1cm}} \text{ mg}$$

A **mg** is **larger** than a **mcg**
Therefore you move decimal
3 places to the **L**eft

$$1500. \text{ mcg} = \underline{1.500} \text{ mg (1.5 mg)}$$

BASED ON WEIGHT

$$1 \text{ kg} = 2.2 \text{ lbs}$$

$$1 \text{ lb} = 16 \text{ oz}$$

EXAMPLE

$$120 \text{ lbs} = \underline{\hspace{1cm}} \text{ kg}$$

$$120 \text{ lbs} / 2.2 = 54.545 \text{ kg}$$

kg → **lb**
MULTIPLY by 2.2

EXAMPLE

$$45.6 \text{ kg} = \underline{\hspace{1cm}} \text{ lbs}$$

$$45.6 \text{ kg} \times 2.2 = 100.32 \text{ lbs}$$

DOSAGE CALC RULES



Medication error kills,
prevention is crucial!

1 Show ALL your work.

2 Leading zeros must be placed before any decimal point.

The decimal point may be missed without the zero



.2 mg should be **0.2 mg**

WHY? **.2** could appear to be **2**

(0.2 mg of morphine is VERY different than 2 mg of morphine!)

3 No trailing zeros.



0.7 mL NOT **0.70 mL**

1 mg NOT **1.0 mg**

WHY? **1.0** could appear to be **10!**

4 Do not round until you have the final answer!

HOW TO ROUND YOUR FINAL ANSWER

If the number in the thousands place is **5 OR GREATER** → The # in the hundredth place is rounded up



1.995 mg is rounded to **2 mg**

1.985 mg is rounded to **1.99 mg**

If the number in the thousands place is **4 OR LESS** → The # is dropped



0.992 mg is rounded to **0.99 mg**

DECIMAL REFERENCE GUIDE

34.732

tens ↑
ones ↑
tenths ↑
hundredths ↑
thousandths ↑

5 Most nursing schools, if not all, do not give partial credit.

This means every step must be done correctly!

FORMULA METHOD

For Volume-Related Dosage Orders

$$\frac{D}{H} \times V = A$$

D = Desired

EXAMPLE: "The physician orders **120 mg**..."



Some medications like heparin and insulin are prescribed in **units/hour**

H = Dosage of medication available

EXAMPLE: "The medication is supplied as **100 mg/5 mL**"

V = Volume the medication is available in

EXAMPLE: "The medication is supplied as 100 mg/**5 mL**"

A = Amount of Medication required for administration

YOUR ANSWER



You should assume that all questions are asked "**per dose**" unless the question gives a timeframe (example: "how many tablets will you give in 24 hours?")

EXAMPLE 1

Ordered: **Drug C 150 mg**
Available: **Drug C 300 mg/tab**
How many tablets should be given?

$$\frac{D}{H} \times V = A$$

What's our desired? **Drug C 150mg PO**
What do we have? **Drug C 300mg/tab**
What's our quantity/volume? **tablets**

$$\frac{150 \text{ mg}}{300 \text{ mg}} \times 1 \text{ tab} = 0.5 \text{ tabs}$$
$$150 \div 300 = 0.5 \times 1 = 0.5 \text{ tabs}$$

FINAL ANSWER: **0.5 tabs**

EXAMPLE 2

Ordered: **Drug C 10,000 units SubQ**
Available: **Drug C 5,000 units/mL**
How many mL should be given?

$$\frac{D}{H} \times V = A$$

What's our desired? **Drug C 10,000 SubQ**
What do we have? **Drug C 5,000 units**
What's our quantity/volume? **1 mL**

$$\frac{10,000 \text{ units}}{5,000 \text{ units}} \times 1 \text{ mL} = 2 \text{ mL}$$
$$10,000 \div 5,000 = 2 \times 1 = 2 \text{ mL}$$

FINAL ANSWER: **2 mL**

IV FLOW RATES

$$\frac{\text{mL of solution}}{\text{total hours}} = \text{mL/hr}$$

What if the question is given in **MINUTES**?

Since there are 60 minutes in one hour, use this formula:

$$\frac{\text{mL of solution}}{\text{min}} \times 60_{\text{(minutes)}} = \text{mL/hr}$$

! If the question is asking for flow rate and you're given units of mL, you need to write the answers in **mL/hr!**

mL/hr is always rounded to the nearest **WHOLE** number!

EXAMPLE #1

ORDERED: 1000 mL D5W to infuse over 3 hours. What will the flow rate be?

$$\frac{1000 \text{ mL}}{3 \text{ hr}} = 333.333 \text{ mL/hr}$$

ANSWER: 333 mL/hr
(rounded to the nearest whole number)

EXAMPLE #2

ORDERED: Infuse 3 grams of Penicillin in 50 mL normal saline over 30 minutes.

$$\frac{50 \text{ mL}}{30 \text{ min}} \times 60 \text{ min} = 100 \text{ mL/hr}$$

ANSWER: 100 mL/hr

$$\frac{\text{mL of solution}}{\text{total minutes}} \times \text{drop factor} = \text{gtt/min}$$

What if the question is given in **HOURS**?

Since there are 60 minutes in one hour, use this formula:

Convert hours to minutes!

EXAMPLES:

1 hour = 60 minutes
2.5 hours = 150 minutes

! If a drop factor is included, the question is asking for flow rate in gtt/min.

You need to write the answers in **gtt/minute!**

REMEMBER OUR ABBREVIATIONS:
gtt means "drop"!

EXAMPLE #1

ORDERED: 1000 mL of Lactated Ringer's to infuse at 50 mL/hr. Drop factor for tubing is a 5 gtt/mL. (Convert: 1 hour = 60 min)

$$\frac{50 \text{ mL}}{60 \text{ min}} \times 5 \text{ gtt/mL} = 4 \text{ gtt/min}$$

$50 \div 60 = 0.833 \times 5 = 4.166$
Round to the nearest whole number $\rightarrow 4$

EXAMPLE #2

ORDERED: 100 mL of Metronidazole to infuse over 45 minutes. The tubing you are using has a drop factor of 10 gtt/mL.

$$\frac{100 \text{ mL}}{45 \text{ min}} \times 10 \text{ gtt/mL} = 22 \text{ gtt/min}$$

$100 \div 45 = 2.222 \times 10 = 22.222$
Round to the nearest whole number $\rightarrow 22$

FINAL ANSWER: 4 gtt/min

! **REMEMBER RULE #4**
Don't round till the end!

FINAL ANSWER: 22 gtt/min

! **REMEMBER RULE #4**
Don't round till the end!

PRACTICE QUESTIONS

Do all 10 questions without looking at the correct answers on the following pages. Don't forget to show all your work. After you are done, walk through each question...even the questions you got correct!

1 **ORDERED:** Rosuvastatin 3000 mcg PO ac
AVAILABLE: Rosuvastatin 2 mg tablet (scored)
How many tabs will you administer in 24 hours?

2 **ORDERED:** Tylenol supp 2 g PR q6h
AVAILABLE: Tylenol supp 700 mg
How many supp will you administer?
Round to nearest tenth.

3 **ORDERED:** Potassium chloride 0.525 mEq/lb PO dissolved in 6 oz of juice at 0930
AVAILABLE: Potassium chloride 12 mEq/mL
How many mL of potassium chloride will you add to the juice for a 66.75 kg patient? Round to nearest tenth.

4 **1000 mL D5W to infuse over 4 hours.**

5 **150 mL Cipro 250 mcg to infuse over 45 minutes.**

6 **250 mL normal saline over 5 hours.**
Tubing drop factor of 10 gtt/mL.

7 **Humulin R 200 units in 100 mL of normal saline to infuse at 4 units/hr.**

8 **Dopamine 600 mg in 200 mL of normal saline to infuse at 10mcg/kg/min. Pt weight = 190 lbs.**

9 **2.5 L normal saline to infuse over 48 hours.**

10 **ORDERED:** Morphine 100 mg IM q12h prn pain
AVAILABLE: Morphine 150 mg/2.6 mL
How many mL will you administer?
Round to nearest hundredth.

COMPREHENSIVE REVIEW

1

ORDERED: Rosuvastatin 3000 mcg PO ac
AVAILABLE: Rosuvastatin 2 mg tablet (scored)

How many tabs will you administer in 24 hours?

STEP 1: CONVERT DATA

mcg → mg

$$3000 \text{ mcg} = 3 \text{ mg}$$

REMEMBER **SMALL TO BIG:**
move the decimal point 3 to the left
unit is getting **L**arger think **L**eft

STEP 2: READY TO USE DATA

ORDERED: 3 mg
AVAILABLE: 2 mg
VOLUME: 1 tab
ADMINISTERED AC: before each meal
QUESTION IS ASKING: dosage in 24 hours

STEP 3: IRRELEVANT DATA

N/A

STEP 4: FORMULA USED

$$\frac{D}{H} \times V = A$$

SHOW YOUR WORK

$$\frac{3 \text{ mg}}{2 \text{ mg}} = 1.5$$

$$1.5 \times 1 \text{ tab} = 1.5$$

$$1.5 \times 3 = 4.5 \text{ tabs per day}$$

ROUND: No rounding necessary

! **DON'T FORGET TO CHECK TIMES OF MEDICATION!**

The medication is ordered to be given **AC**, which means **before each meal**. Since there are 3 meals in a day (24 hours), the answer must be multiplied by 3.

FINAL ANSWER: 4.5 tabs

2

ORDERED: Tylenol supp 2 g PR q6h
AVAILABLE: Tylenol supp 700 mg

How many supp will you administer? Round to nearest tenth.

STEP 1: CONVERT DATA

g → mg

$$2 \text{ g} = 2000 \text{ mg}$$

REMEMBER **BIG TO SMALL:**
move the decimal point 3 to the right

STEP 2: READY TO USE DATA

ORDERED: 2000 mg
AVAILABLE: 700 mg
VOLUME: 1 supp

STEP 3: IRRELEVANT DATA

N/A

STEP 4: FORMULA USED

$$\frac{D}{H} \times V = A$$

SHOW YOUR WORK

$$\frac{2000 \text{ mg}}{700 \text{ mg}} = 2.857$$

$$2.857 \times 1 \text{ supp} = 2.857 \text{ supp}$$

ROUND: Nearest tenth

$$2.857 \text{ supp} \rightarrow 2.9 \text{ supp}$$

! **REMEMBER RULE #4**
Don't round till the end!

FINAL ANSWER: 2.9 supp

COMPREHENSIVE REVIEW

3

ORDERED: Potassium chloride 0.525 mEq/lb PO dissolved in 6 oz of juice at 0930

AVAILABLE: Potassium chloride 12 mEq/mL

How many mL of potassium chloride will you add to the juice for a 66.75 kg patient? Round to nearest tenth.

STEP 1: CONVERT DATA

kg → lb

$$66.75 \text{ kg} \times 2.2 \text{ (lb/kg)} = 146.85 \text{ lb}$$

! In this case, ordered amount depends on patient weight

mEq/lb → mEq

$$(0.525 \text{ mEq/lb} \times 146.85 \text{ lb}) = 77.096 \text{ mEq}$$

STEP 2: READY TO USE DATA

ORDERED: 77.096 mEq

AVAILABLE: 12 mEq

VOLUME: 1 mL

STEP 3: IRRELEVANT DATA

Dissolved in 12 oz of juice at 0930

! Question asked for "per dose" because no timeframe was given

STEP 4: FORMULA USED

$$\frac{D}{H} \times V = A$$

SHOW YOUR WORK

$$\frac{77.096 \text{ mEq}}{12 \text{ mEq}} = 6.424$$

$$6.424 \times 1 \text{ mL} = 6.424 \text{ mL}$$

! **REMEMBER RULE #4**
Don't round till the end!

ROUND: Nearest tenth

$$6.424 \text{ mL} \rightarrow 6.4 \text{ mL}$$

FINAL ANSWER: 6.4 mL

4

1000 mL D5W to infuse over 4 hours.

STEP 1: CONVERT DATA

N/A

STEP 2: READY TO USE DATA

1000 mL

4 hr

STEP 3: IRRELEVANT DATA

N/A

STEP 4: FORMULA USED

$$\frac{\text{mL of solution}}{\text{total hours}} = \text{mL/hr}$$

SHOW YOUR WORK

$$\frac{1000 \text{ mL}}{4 \text{ hr}} = 250 \text{ mL/hr}$$

! mL/hr is always rounded to the nearest **WHOLE** number!

ROUND: No rounding necessary

FINAL ANSWER: 250 mL/hr

COMPREHENSIVE REVIEW

5

150 mL Cipro 250 mcg to infuse over 45 minutes.



If the question is asking for flow rate ("to infuse") and you're given mL of solution, you need to write the answer in mL/hours!

STEP 1: CONVERT DATA

N/A

STEP 2: READY TO USE DATA

ML OF SOLUTION: 150 mL
TOTAL HOURS: 45 min

STEP 3: IRRELEVANT DATA

Cipro 250 mcg

IMPORTANT: don't let this information lead you to use the wrong formula. In this example, we're asked for a flow rate which requires mL of solution and total time.

STEP 4: FORMULA USED

$$\frac{\text{mL of solution}}{\text{total minutes}} \times 60 = \text{mL/hr}$$

SHOW YOUR WORK

$$\frac{150 \text{ mL}}{45 \text{ min}} = 3.333 \times 60 = 200 \text{ mL/hr}$$

REMEMBER RULE #4
Don't round till the end!

REMEMBER!
mL/hr is always rounded to the nearest **WHOLE** number!

ROUND: No rounding necessary

FINAL ANSWER: 200mL/hr

6

250 mL normal saline over 5 hours. Tubing drop factor of 10 gtt/mL.

STEP 1: CONVERT DATA

hr → min

1 hour = 60 minutes

$$5 \text{ hr} \times \frac{60 \text{ min}}{1 \text{ hr}} = 300 \text{ min}$$

STEP 2: READY TO USE DATA

ML OF SOLUTION: 250 mL
TOTAL MINUTES: 300 min
DROP FACTOR: 10 gtt/mL

STEP 3: IRRELEVANT DATA

N/A

STEP 4: FORMULA USED

$$\frac{\text{mL of IV solution}}{\text{time in minutes}} \times \text{drop factor} = \text{gtt/min}$$

SHOW YOUR WORK

$$\frac{250 \text{ mL}}{300 \text{ min}} = 0.8333 \text{ mL/min}$$

REMEMBER RULE #4
Don't round till the end!

$$0.8333 \text{ mL/min} \times 10 \text{ gtt/mL} = 8.3333 \text{ gtt/min}$$

ROUND: gtt/mL is always rounded to the nearest whole number!

$$8.3333 \text{ gtt/min} \rightarrow 8 \text{ gtt/min}$$

REMEMBER!
The question may not specify to round the final answer to a whole number; you are expected to know this with gtt/min units.

FINAL ANSWER: 8 gtt/min

COMPREHENSIVE REVIEW

7

Humulin R **200 units** in **100 mL** of normal saline to infuse at **4 units/hr**.

STEP 1: CONVERT DATA

N/A

STEP 2: READY TO USE DATA

DESIRED: 4 units/hr
AVAILABLE: 200 units
VOLUME: 100 mL

STEP 3: IRRELEVANT DATA

N/A

STEP 4: FORMULA USED

$$\frac{D}{H} \times V = A$$

SHOW YOUR WORK

$$\frac{4 \text{ units/hr}}{200 \text{ units}} = 0.02 \text{ /hr}$$

$$0.02 \text{ /hr} \times 100 \text{ mL} = 2 \text{ mL/hr}$$

ROUND: No rounding necessary

!
 mL/hr is always rounded to the nearest **WHOLE** number!

FINAL ANSWER: 2 mL/hr

8

Dopamine **600 mg** in **200 mL** of normal saline to infuse at **10 mcg/kg/min**.
 Pt weight = 190 lbs.

! If the question is asking for flow rate ("to infuse") and you're given mL of solution, you need to write the answer in **mL/hours!**

STEP 1: CONVERT DATA

$$\text{mcg} \rightarrow \text{mg}$$

$$10 \text{ mcg} = 0.010 \text{ mg}$$

REMEMBER

SMALL TO BIG:
 move the decimal point 3 to the left
 unit is getting **L**arger think **L**eft

$$\text{lb} \rightarrow \text{kg}$$

$$190 \text{ lb} / 2.2 = 86.363 \text{ kg}$$

$$\frac{\text{mg}}{\text{kg}} \rightarrow \frac{\text{mg}}{\text{min}}$$

! In this case, ordered amount depends on patient weight

$$0.010 \text{ mg/kg/min} \times 86.363 \text{ kg} = 0.863 \text{ mg/min}$$

STEP 2: READY TO USE DATA

DESIRED: 0.863 mg/min
AVAILABLE: 600 mg
VOLUME: 200 mL

STEP 3: IRRELEVANT DATA

N/A

STEP 4: FORMULA USED

$$\frac{D}{H} \times V = A$$

SHOW YOUR WORK

$$\frac{0.863 \text{ mg/min}}{600 \text{ mg}} = 0.00143 \text{ /min}$$

$$0.00143 \text{ /min} \times 200 \text{ mL} = 0.2878 \text{ mL/min}$$

$$0.2878 \text{ mL/min} \times 60 \text{ min} = 17.2727 \text{ mL/hr}$$

ROUND: mL/hr is always rounded to nearest whole number!

$$17.2727 \text{ mL/hr} \rightarrow 17 \text{ mL/hr}$$

WAIT!
 This is in mL/min ... we need units of mL/hr!

FINAL ANSWER: 17 mL/hr

COMPREHENSIVE REVIEW

9

2.5 L normal saline to infuse over 48 hours.

! If the question is asking for flow rate ("to infuse") and you're given mL of solution, you need to write the answer in **mL/hours!**

STEP 1: CONVERT DATA

L → mL

REMEMBER **BIG TO SMALL:**
move the decimal point 3 to the right

$$2.5 \text{ L} = 2500 \text{ mL}$$

STEP 2: READY TO USE DATA

ML OF SOLUTION: 2500 mL
TOTAL HOURS: 48 hr

STEP 3: IRRELEVANT DATA

N/A

STEP 4: FORMULA USED

$$\frac{\text{mL of solution}}{\text{total hours}} = \text{mL/hr}$$

SHOW YOUR WORK

$$\frac{2500 \text{ mL}}{48 \text{ hours}} = 52.0833 \text{ mL/hr}$$

ROUND: mL/hr is always rounded to nearest whole number!

$$52.0833 \text{ mL/hr} \rightarrow 52 \text{ mL/hr}$$

FINAL ANSWER: 52 mL/hr

10

ORDERED: Morphine 100 mg IM q12h prn pain

AVAILABLE: Morphine 150 mg/2.6 mL

How many mL will you administer?

Round to nearest hundredth.

STEP 1: CONVERT DATA

N/A

STEP 2: READY TO USE DATA

ORDERED: 100 mg
AVAILABLE: 150 mg
VOLUME: 2.6 mL

STEP 3: IRRELEVANT DATA

IM q12h prn pain

! Question asked for "per dose" because no timeframe was given

STEP 4: FORMULA USED

$$\frac{D}{H} \times V = A$$

SHOW YOUR WORK

$$\frac{100 \text{ mg}}{150 \text{ mg}} = 0.6666$$

$$0.6666 \times 2.6 \text{ mL} = 1.7333 \text{ mL}$$

ROUND: nearest hundredth

$$1.7333 \text{ mL} \rightarrow 1.73 \text{ mL}$$

FINAL ANSWER: 1.73 mL



LAB VALUE CHEAT SHEET

WITH MEMORY TRICKS

BROUGHT TO YOU BY



LAB VALUE CHEAT SHEET

VITAL SIGNS

BLOOD PRESSURE	SYSTOLIC	120 mmHg
	DIASTOLIC	80 mmHg
HEART RATE		60 – 100 bpm
RESPIRATIONS		12 – 20 breaths/min
TEMPERATURE		97.8 – 99°F (36.5 – 37.2°C)
OXYGEN		95 – 100%
OXYGEN IN COPD PT.		as low as 88%



COPD pts are expected to have low O₂ levels

RENAL

CALCIUM	9 – 11 mg/dL
MAGNESIUM	1.5 – 2.5 mg/dL
PHOSPHORUS	2.5 – 4.5 mg/dL
SPECIFIC GRAVITY	1.010 – 1.030
GFR	90 – 120 mL/min/1.73 m ²
BUN	7 – 20 mg/dL
CREATININE	0.6 – 1.2 mg/dL



PANCREAS

AMYLASE	30 – 110 U/L
LIPASE	0 – 150 U/L



COMPLETE BLOOD COUNT (CBC)

WBCs	4,500 – 11,000 /μL
RBCs	4.5 – 5.5 X10 ⁶ /μL
PLTs	150,000 – 450,000 /μL
HEMOGLOBIN (HGB)	FEMALE: 12 – 16 g/dL MALE: 13 – 18 g/dL
HEMATOCRIT (HCT)	FEMALE: 36% – 48% MALE: 39% – 54%



BASIC METABOLIC PANEL (BMP)

SODIUM	135 – 145 mEq/L
POTASSIUM	3.5 – 5.0 mEq/L
CHLORIDE	95 – 105 mEq/L
CALCIUM	9 – 11 mg/dL
BUN	7 – 20 mg/dL
CREATININE	0.6 – 1.2 mg/dL
ALBUMIN	3.4 – 5.4 g/dL
TOTAL PROTEIN	6.2 – 8.2 g/dL

HBA1C

NON-DIABETIC	4 – 5.6%
PRE-DIABETIC	5.7 – 6.4%
DIABETIC	> 6.5%
Goal for diabetic:	< 6.5%

LIVER FUNCTION TEST (LFT)

ALT	7 – 56 U/L
AST	5 – 40 U/L
ALP	40 – 120 U/L
BILIRUBIN	0.1 – 1.2 mg/dL

BMI

UNDERWEIGHT	<18.5
HEALTHY WEIGHT	18.5 – 24.9
OVERWEIGHT	25.0 – 29.9
OBESITY	> 30.0

ABGs

PH	7.35 – 7.45
PaCO ₂	35 – 45 mmHg
PaO ₂	80 – 100 mmHg
HCO ₃	22 – 26 mEq/L



COAGs

PT	10 – 13 sec
PTT	25 – 35 sec
aPTT	NOT ON HEPARIN: 30–40 secs ON HEPARIN: 47–70 secs
INR	NOT ON WARFARIN: < 1 sec ON WARFARIN: 2 – 3 sec



LIPID PANEL

TOTAL CHOLESTEROL	<200 mg/dL
TRIGLYCERIDE	<150 mg/dL
LDL	<100 mg/dL
HDL	>60 mg/dL

LDL bad cholesterol - we want **LOW** levels

HDL Happy cholesterol - we want **HIGH** levels

OTHER

MAP (mean arterial pressure)	70 – 100 mmHg	
ICP (intracranial pressure)	5 – 15 mmHg	
GLASCOW COMA SCALE	BEST = 15	
MILD: 13-15	MODERATE: 9-12	SEVERE: 3-8

Lab values, instruments, and institutions differ based on the facility. Local policy should supersede. Author & publisher intend this reference to be free of errors but no guarantee can be made & assume no responsibility for any outcomes resulting from its use.



LAB VALUE MEMORY TRICKS



ELECTROLYTES

SODIUM: 135 - 145

*Commit to memory!



POTASSIUM: 3.5 - 5

BANANAS:

There are about 3-5 in every bunch & you want them half ripe (1/2)

So, think 3.5 - 5.0



PHOSPHORUS: 2.5 - 4.5

PHOR: 4

US: 2 (me + you = 2)



*don't forget the .5

CALCIUM: 9 - 11

CALL 911



MAGNESIUM: 1.5 - 2.5

MAGnifying glass you see 1.5 - 2.5 bigger than normal



CHLORIDE: 95 - 105

Think of a chlorinated pool that you want to go in when it's SUPER HOT: 95 - 105 °F



COMPLETE BLOOD COUNT (CBC)

- Hemoglobin (Hgb)
 - Female: 12 - 16 g/dL
 - Male: 13 - 18 g/dL
- Hematocrit (HCT)
 - Female: 36% - 48%
 - Male: 39% - 54%



To remember HCT, multiply Hgb by 3

$$12 \times 3 = 36 \text{ (Female)}$$

$$16 \times 3 = 48 \text{ (Female)}$$

$$13 \times 3 = 39 \text{ (Male)}$$

$$18 \times 3 = 54 \text{ (Male)}$$

BASAL METABOLIC PANEL (BMP)


BUN: 7 - 20 mg/dL


Think hamburger **BUN**s...
Hamburgers can cost anywhere from \$7 - \$20 dollars



CREATINE: 0.6 - 1.2 mg/dL

This is the same value as **LITHIUM**'s therapeutic range (0.6 - 1.2 mmol/L)
Lithium is excreted almost solely by the kidneys...
And creatine is a value that tests how well your kidneys filter





ELECTROLYTE IMBALANCES

BROUGHT TO YOU BY



SODIUM IMBALANCE

SODIUM is a major **ELECTROLYTE** found in ECF. Essential for acid-base, fluid balance, active & passive transport mechanism, irritability & **CONDUCTION** of nerve-muscle tissue

135 - 145 mEq/L

↑ HYPERNATREMIA > 145 mEq/L

↓ HYPONATREMIA < 135 mEq/L

SIGNS & SYMPTOMS

* **BIG & BLOATED**

- F**lushed skin
- R**estless, anxious, confused, irritable
- I**ncreased BP & fluid retention
- E**dema (pitting)
- D**ecreased urine output
- S**kin flushed & dry
- A**gitation
- L**ow-grade fever
- T**hirst (dry mucous membranes)

HYPOVOLEMIC HYPONATREMIA:

↓ of fluid & sodium

- S**tupor/coma
- A**norexia (nausea/vomiting)
- L**ethargy (weakness/fatigue)
- T**achycardia (thready pulse)

HYPERVOLEMIC HYPONATREMIA:

↑ body water that is greater than Na+

- L**imp muscles (muscle weakness)
- O**rthostatic hypotension
- S**eizures/headache
- S**tomach cramping (hyperactive bowels)

RISK FACTORS

- * Increased sodium intake
 - ↳ Excess oral sodium ingestion
 - ↳ Excess administration of IV fluids w/ sodium
 - ↳ Hypertonic IV fluids
- * **LOSS OF FLUIDS!**
 - ↳ Fever
 - ↳ Watery diarrhea
 - ↳ Diabetes insipidus
 - ↳ Excessive diaphoresis
 - ↳ Infection
- * Decreased sodium excretion
 - ↳ Kidney problems

HEMOCONCENTRATION = INCREASED SODIUM!

- * Increased sodium excretion
 - ↳ Diaphoresis (ex: high fever)
 - ↳ Diarrhea & vomiting
 - ↳ Drains (NGT suction)
 - ↳ Diuretics (Thiazide & loop diuretics)

4 D's

- * SIADH
- * Adrenal insufficiency (adrenal crisis)
- * Inadequate sodium intake
 - ↳ Fasting, NPO, Low-salt diet
- * Kidney disease
- * Heart failure

MANAGEMENT

- * If due to fluid loss:
 - ↳ Administer IV infusions
- * If the cause is inadequate renal excretion of sodium:
 - ↳ Give diuretics that promote sodium loss
- * Restrict sodium & fluid intake as prescribed

A D D S A L T

- A** **ADMINISTER** IV sodium chloride infusions (Only if due to hypovolemia)
- D** **DIURETICS** (If due to hypervolemia) Hyponatremia → high fluids & low salt = hemodilution
- D** **DAILY WEIGHTS** Where sodium goes, water FLOWS
- S** **SAFETY** (orthostatic hypotension AKA risk for falls)
- A** **AIRWAY PROTECTION** (NPO) Don't give food to a lethargic, confused client (INCREASED RISK FOR ASPIRATION)
- L** **LIMIT WATER INTAKE** Hypervolemic hyponatremia (high fluid & low salt)
- T** **TEACH** about foods high in sodium (Canned food, packaged/processed meats, etc.)

POTASSIUM IMBALANCE

POTASSIUM imbalance plays a vital role in cell METABOLISM, and TRANSITION of nerve impulses, the functioning of cardiac, lung, muscle tissues, & acid-base balance.

3.5 - 5 mEq/L

↑ HYPERKALEMIA > 5 mEq/L

↓ HYPOKALEMIA < 3.5 mEq/L

SIGNS & SYMPTOMS

* TIGHT & CONTRACTED

- M uscle cramps & weakness
- U rine abnormalities
- R espiratory distress
- D ecreased cardiac contractility (↓HR, ↓BP)
- E CG changes
 - Tall peaked T waves
 - Flat P waves
 - Widened QRS complexes
 - Prolonged PR intervals
- R eflexes (↑ DTR)

- * Thready, weak, irregular pulse
- * Orthostatic hypotension
- * Shallow respirations
- * Anxiety, lethargy, confusion, coma
- * Paresthesias
- * Hyporeflexia
- * Hypoactive bowel sounds (constipation)
- * Nausea, vomiting, abdominal distention
- * ECG changes
 - ST depression
 - Shallow or inverted T wave
 - Prominent U wave

RISK FACTORS

- * Medication
 - ↳ Potassium-sparing diuretics (Spironolactone)
 - ↳ Ace inhibitors
 - ↳ NSAIDs
- * Excessive potassium intake
(Example: rapid infusion of potassium-containing IV solutions)
- * Kidney disease or those on Dialysis
 - ↳ Decreased potassium excretion
- * Adrenal insufficiency (Addison's disease)
- * Tissue damage
- * Acidosis
- * Hyperuricemia
- * Hypercatabolism

- * Actual total body potassium loss
- * Inadequate potassium intake
 - ↳ Fasting, NPO
- * Movement of potassium from the extracellular fluid to the intracellular fluid
 - ↳ Alkalosis
 - ↳ Hyperinsulinism
- * Dilution of serum potassium
 - ↳ Water intoxication
 - ↳ IV therapy with potassium-deficient solutions

POTASSIUM IMBALANCE CAN CAUSE CARDIAC DYSRHYTHMIAS THAT CAN BE LIFE-THREATENING!

MANAGEMENT

- * Monitor EKG
- * Discontinue IV & PO potassium
- * Initiate a potassium-restricted diet
- * Potassium-excreting diuretics
- * Prepare the client for dialysis
- * Prepare for administration:
 - ↳ IV calcium gluconate & IV sodium bicarb
- * Avoid the use of salt substitutes or other potassium-containing substances

- * Oral potassium supplements
- * Liquid potassium chloride
- * Potassium-retaining diuretic
- * Potassium is **NEVER** administered by IV push, IM, or subcut routes.
 - ↳ IV potassium is always diluted & administered using an infusion device!



POTASSIUM & SODIUM = OPPOSITES

EXAMPLE: ↑ NA = ↓ K+

CALCIUM IMBALANCE

CALCIUM is found in the body's cells, bones, and teeth. Needed for proper functioning of the **CARDIOVASCULAR, NEUROMUSCULAR, ENDOCRINE** systems, blood clotting & teeth formation

9 - 11 mg/dL

↑
HYPERCALCEMIA > 11 mg/dL

↓
HYPOCALCEMIA < 9 mg/dL

SIGNS & SYMPTOMS

- B**one pain
- A**rrhythmias
- C**ardiac arrest (bounding pulses)
- K**idney stones
- M**uscle weakness ↓ (DTR)
- E**xcessive urination

- C**onvulsions
- A**rrhythmias (diminished pulses)
- T**etany
- S**pasms & stridor
- GO NUMB**ness in the fingers, face, & limbs



POSITIVE TROUSSEAU'S:
Carpal spasm caused by inflating a blood pressure cuff



CHVOSTEK'S SIGNS:
Contraction of facial muscles w/ light tap over the facial nerve.

MEMORY TRICK
Think "C"
for Cheesy smile

RISK FACTORS

- * Increased calcium absorption
- * Decreased calcium excretion
- * Kidney disease
- * Thiazide diuretics
- * Increased bone resorption of calcium
 - ↳ Hyperparathyroidism / Hyperthyroidism
 - ↳ Malignancy (bone destruction from metastatic tumors)
- * Hemoconcentration
- * Inhibition of calcium absorption from the GI tract
- * Increased calcium excretion
 - ↳ Kidney disease, diuretic phase
 - ↳ Diarrhea & steatorrhea
 - ↳ Wound drainage
- * Conditions that decrease the ionized fraction of calcium

MANAGEMENT

- * D/C IV or PO calcium
- * D/C Thiazide diuretics
- * Administer phosphorus, calcitonin, bisphosphonates, & prostaglandin synthesis inhibitors (NSAIDs)
- * Avoid foods high in calcium
- * Adm. calcium PO or IV
 - ↳ For IV, warm before & adm. slowly
- * Adm. aluminum hydroxide & Vit D
- * Initiate seizure precautions
- * 10% calcium (acute calcium deficit)
- * Consume foods high in calcium

A CLIENT WITH A CALCIUM IMBALANCE IS AT RISK FOR A PATHOLOGICAL FRACTURE. MOVE THE CLIENT CAREFULLY AND SLOWLY

CALCIUM & PHOSPHATE = INVERSE

EXAMPLE: ↑ CA+ = ↓ PO4

MAGNESIUM IMBALANCE

Most of the **MAGNESIUM** found in the body is found in the bones.
Regulates BP, blood sugar, muscle contraction & nerve function.

1.5 - 2.5 mg/dL

↑
HYPERMAGNESEMIA > 2.5 mg/dL

↓
HYPOMAGNESEMIA < 1.5 mg/dL



MAGNESIUM IS A SEDATIVE!

SIGNS & SYMPTOMS

* LOW EVERYTHING AKA **SEDATED**

- * Low energy (drowsiness / coma)
- * Low HR (bradycardia)
- * Low BP (hypotension)
- * Low RR (bradypnea)
- * ↓ Respirations (shallow)
- * ↓ Bowel sounds
- * ↓ DTR's (deep tendon reflex)

* HIGH EVERYTHING AKA **NOT SEDATED**

- * High HR (tachycardia)
- * High BP (hypertension)
- * Increased deep tendon reflex (hyperreflexia)
- * Shallow respirations
- * Twitches, paresthesias
- * Tetany & seizures
- * Irritability & confusion

REMEMBER!
Also seen in hypocalcemia.
Ca & Mg rise and fall together!

POSITIVE TROUSSEAU'S:

Carpal spasm caused by inflating a blood pressure cuff

CHVOSTEK'S SIGNS:

Contraction of facial muscles w/ light tap over the facial nerve

RISK FACTORS

- * Increased magnesium intake
 - ↳ Magnesium-containing antacids (TUMS) & laxatives
 - ↳ Excessive adm. of magnesium IV
- * Renal insufficiency
 - ↳ ↓ renal excretion of Mg = ↑ Mg in the blood
- * DKA (Diabetic Ketoacidosis)

- * Insufficient magnesium intake
 - ↳ Malnutrition/vomiting/diarrhea
 - ↳ Malabsorption syndrome
 - ↳ Celiac & Crohn's disease
- * Increased magnesium excretion
 - ↳ Diuretics or chronic alcoholism
- * Intracellular movement of magnesium
 - ↳ Hyperglycemia & Insulin adm.
 - ↳ Sepsis

MANAGEMENT

- * Diuretics
- * IV adm. calcium chloride or calcium gluconate
- * Restrict dietary intake of Mg containing foods
- * Avoid the use of laxatives & antacids containing magnesium
- * Hemodialysis
- * Magnesium sulfate IV or PO
- * Seizure precautions
- * Instruct the client to increase magnesium-containing foods

MAGNESIUM & CALCIUM = SAME

EXAMPLE: ↑ MG = ↑ CA+




FUNDAMENTALS

BROUGHT TO YOU BY








BLOOD TYPES



CENTRIFUGE

A device that uses force to separate components of fluids. It separates fluids of different densities. This is how labs separate blood.



ANTIGEN:	A
ANTIBODIES:	B
RECIPIENT:	A, O
DONOR:	A, AB

ANTIGEN:	B
ANTIBODIES:	A
RECIPIENT:	B, O
DONOR:	B, AB

ANTIGEN:	A & B
ANTIBODIES:	NONE
RECIPIENT:	ALL
DONOR:	AB

ANTIGEN:	NONE
ANTIBODIES:	A & B
RECIPIENT:	O
DONOR:	ALL

PLASMA ANTIBODIES

- Protects body from "invaders" (think ANTI)
- Opposite of the type of antigen that is found on the RBC

ANTIGENS


- Proteins that elicit immune response
- Identifies the cell

DONOR BLOOD TYPES

RECIPIENT BLOOD TYPES

	O-	O+	A-	A+	B-	B+	AB-	AB+
O-	♥							
O+	♥	♥						
A-	♥		♥					
A+	♥	♥	♥	♥				
B-	♥				♥			
B+	♥	♥			♥	♥		
AB-	♥		♥		♥		♥	
AB+	♥	♥	♥	♥	♥	♥	♥	♥

Rh FACTOR



ABBREVIATIONS

AAA Abdominal Aortic Aneurysm
Abd Abdomen
Ac Before Meals
ACLS Advanced Cardiac Life Support
AD Admitting Diagnosis
A&D Admission and Discharge
Ad lib As Desired
ALL Acute Lymphocytic Leukemia
ADL Activities of Daily Living
Adm. Admission
Amb Ambulation
AKA Above-the-Knee Amputation
AV Atrioventricular
AP or A.P. Appendectomy
Bid Twice a Day
BLS Basic Life Support
BM Bowel Movement
BP Blood Pressure
BKA Below-the-Knee Amputation
BUN Blood Urea Nitrogen
BPH Benign Prostatic Hyperplasia

BX Biopsy
CABG Coronary Artery Bypass Graft
C/O Complaining Of
CAD Coronary Artery Disease
CBC Complete Blood Count
CCU Cardiac Care Unit / Coronary Care Unit
C&S Culture & Sensitivity
CF Cystic Fibrosis
CHF Congestive Heart Failure
CKD Chronic Kidney Disease
CPR Cardiopulmonary Resuscitation
COPD Chronic Obstructive Pulmonary Disease
CVA Cerebrovascular Accident (stroke)
CVC Central Venous Catheter
D/C Discontinue or Discharge
D&C Dilatation and Curettage
DI Diabetes Insipidus
DIC Disseminated Intravascular Coagulation
DKA Diabetic Ketoacidosis
DM Diabetes Mellitus
DVT Deep Vein Thrombosis

DX Diagnosis
ECG or EKG Electrocardiogram
ED Emergency Department
EENT Eye, Ears, Nose and Throat
ETT Endotracheal Tube
FBS Fasting Blood Sugar
Fx Fracture
Gtt or G.T.T. Glucose Tolerance Test
HOB Head of Bed
HS Bedtime
Hx History
ICU Intensive Care Unit
LMP Last Menstrual Period
LOC Level of Consciousness
LES Lower Esophageal Sphincter
LP Lumbar Puncture
I&O Intake and Output
MAP Mean Arterial Pressure
MRI Magnetic Resonance Imaging
MVA Motor Vehicle Accident
NGT Nasogastric Tube

NPO Nothing by Mouth
NKA No Known Allergies
O₂ Oxygen
OB Obstetrics
OOB Out of Bed
OR Operating Room
OA Osteoarthritis
Ortho Orthopedics
OT Occupational Therapist
Pc After Meals
Prn or p.r.n. As Needed
Pre op Before Surgery
PFT Pulmonary Function Test
PLT Platelets
PTCA Percutaneous Transluminal Coronary Angioplasty
PRBC Packed Red Blood Cells
PVC Premature Ventricular Contraction
Rom/R.O.M. Range of Motion
RBC Red Blood Cell
RT Respiratory Therapist
RA Rheumatoid Arthritis

SOB Shortness of Breath
SBAR Situation, Background, Assessment, Recommendation
SSE or S.S.E. Soap Suds Enema
Stat At Once, Immediately
SLE Systemic Lupus Erythematosus
STD Sexually Transmitted Disease
SIADH Syndrome of Inappropriate Antidiuretic Hormone Secretion
Tid Three Times a Day
T&S Type and Screen
TPN Total Parenteral Nutrition
TIA Transient Ischemic Attack
TB Tuberculosis
TURP Transurethral Resection of the Prostate
UA Urinalysis
UTI Urinary Tract Infection
US Ultrasound
VS Vital Signs
WBC White Blood Count
WNL Within Normal Limits

DO NOT USE

POTENTIAL PROBLEM

INSTEAD, WRITE:

U	Mistaken for "0" (zero) or "cc"	unit
IU	Mistaken for IV (intravenous) or the number 10 (ten)	"international unit"
Q.D., QD, q.d., qd, Q.O.D., QOD, q.o.d., qod	Mistaken for each other	"daily" or "every other day"
Trailing zero (X.0 mg) Lack of leading zero (.X mg)	Decimal point is missed	"X mg" "0.X mg"
MS, MSO ₄ , MgSO ₄	Can mean morphine sulfate or magnesium sulfate	"morphine sulfate" "magnesium sulfate"
@	Mistaken for the number "2" (two)	"at"
cc	Mistaken for U (units) when poorly written	"mL" or "milliliters"

THE NURSING PROCESS

"A DELICIOUS PIE"

SUBJECTIVE DATA

What the client tells the nurse

OBJECTIVE DATA

Data the nurse obtains through their assessment & observation

SET SMART GOALS

SPECIFIC
MEASURABLE
ACHIEVABLE
RELEVANT
TIME FRAME

ASSESS

Gather information
 Verify the information collected is clear & accurate

EVALUATE

Determine the outcome of goals
 Evaluate client's compliance
 Document client's response to pain
 Modify & assess for needed changes

DIAGNOSE

Interpret the information collected

Identify & prioritize the problem through a nursing diagnosis (be sure it's NANDA approved)

IMPLEMENT

Reaching those goals through performing the nursing actions
 "Implementing" the goals set above in the planning stage

PLAN

Set goals to solve the problem.
 Prioritize the outcomes of care

PRIORITY QUESTIONS

ABCs

- A** AIRWAY
- B** BREATHING
- C** CIRCULATION



YOU KNOW YOU ARE BEING ASKED A **PRIORITY QUESTION** WHEN THE QUESTION ASKS:

- What is the **most important**?
- What is the **initial response**?
- Which action should the nurse take **first**?

When you see these questions, you should immediately think of

MASLOW'S HIERARCHY OF NEEDS

as well as **ABCs!**

#1 PATENT AIRWAY

- Patent means "open"; the airway is clear!
- *ASK YOURSELF: Can they successfully breathe oxygen in and breathe CO₂ out?*

#2 BREATHING

- Gas exchange taking place inside the lungs
- *ASK YOURSELF: Can gas exchange successfully happen in their lungs?*

#3 CIRCULATION

- Can they circulate blood through their body and are their organs being perfused?
- *ASK YOURSELF: Is there a reason that the blood isn't pumping/circulating in the body?*
(Example: The heart is working to pump the blood to the vital organs)

MASLOW'S HIERARCHY OF BASIC NEEDS

This shows the **5** levels of human needs

PHYSIOLOGICAL NEEDS being the most important (Oxygen, fluids, nutrition, shelter). **ABCs** fall into Maslow's **PHYSIOLOGICAL** need!

NCLEX TIP
Pain is considered "psychological" meaning it does not take priority.
*Pain rarely kills people



NURSING ETHICS & LAW

ETHICAL PRINCIPLES

AUTONOMY

Respect for an individual's right to make their own decisions

NONMALEFICENCE

Obligation to do & cause no harm to others

BENEFICENCE

Duty to do good to others

JUSTICE

Distribution of benefits & services fairly

VERACITY

Obligation to tell the truth

FIDELITY

Following through with a promise

HIPAA

THE HEALTH INSURANCE PORTABILITY & ACCOUNTABILITY ACT

- Clients records are private & they have the right to ensure the medical information is not shared without permission
- All health care professionals must inform the client how their health information is used
- The client has the right to obtain a copy of their personal health information

PATIENT RIGHTS

THE RIGHT TO...

- Privacy
- Considerate & respectful care
- Be informed
- Know the names & roles of the persons who are involved in care
- Consent or refuse treatment
- Have an advance directive
- Obtain their own medical records & results

CONSENT

TYPES OF CONSENT:

- Admission agreement
 - Immunization consent
 - Blood transfusion consent
 - Surgical consent
 - Research consent
 - Special consents
- Treatment can not be done without a client's consent
 - In the case of an emergency when a client cannot give consent, then consent is implied through emergency laws
 - Minors (under 18), consent must be obtained from a parent or legal guardian



Before signing the consent, the client must be informed of the following: risks & benefits of surgery, treatments, procedures, & plan of care in layman's terms so the client understands clearly what is being done.


INFECTION CONTROL

PPE → PERSONAL PROTECTIVE EQUIPMENT


DONNING PUTTING ON PPE

- Put on PPE before entering the client's room
- Do not touch your face while wearing PPE
- Minimize contact with items in the client's room

1 HAND HYGIENE



2 GOWN



3 MASK / RESPIRATOR



4 GOGGLES / FACE SHIELD



5 GLOVES



DOFFING REMOVING PPE

- Remove PPE at the client's doorway or outside the room
- If hands become soiled while removing PPE, stop & perform hand hygiene
- After hand hygiene, continue with PPE removal


1 REMOVE GLOVES




2 REMOVE PROTECTIVE EYEWEAR




3 REMOVE GOWN



4 REMOVE & DISCARD RESPIRATOR



5 PERFORM HAND HYGIENE



HOSPITAL-ASSOCIATED INFECTIONS (HAIs)

CAUTI..... Catheter-associated urinary tract infection


SSI Surgical site infection

CLABSI..... Central line-associated blood infection

VAP Ventilator-associated pneumonia

Meticulous hand hygiene practices and use of chlorhexidine washes helps in preventing HAIs

INFECTION CONTROL



STAGES OF INFECTION

INCUBATION

Interval between the pathogen entering the body & the presentation of the first symptom

PRODROMAL STAGE

Onset of general symptoms to more distant symptoms; the pathogen is multiplying

ILLNESS STAGE

Symptoms specific to the infection appear

CONVALESCENCE

Acute symptoms disappear and total recovery could take days to months

TRANSMISSION BASED PRECAUTIONS

AIRBORNE

- Single room under negative pressure
- Door remains closed
- Health care workers wear a respiratory mask (N95 or higher level)

M measles

T tuberculosis

V varicella (Chickenpox) & Disseminated herpes-zoster (Shingles)



**Airborne precaution is no longer needed when all lesions have crusted over.*

DROPLET

- Private room or a client whose body cultures contain the same organism
- Wear a surgical mask
- Place a mask on the client whenever they leave the room

- Adenovirus
- Diphtheria (pharyngeal)
- Epiglottitis
- Influenza (flu)
- Meningitis
- Mumps
- Parvovirus B19
- Pertussis
- Pneumonia
- Rubella
- Scarlet fever
- Sepsis
- Streptococcal pharyngitis

CONTACT

- Private room or cohort client
- Use gloves & a gown whenever entering the client's room

- Colonization or infection with a multidrug-resistant organism

- Enteric infections (Clostridium difficile)
- Respiratory infections (RSV, Influenza)


- Wound & skin infections (cutaneous diphtheria, herpes simplex, impetigo, pediculosis, scabies, staphylococci, & varicella-zoster)
- Eye infections (conjunctivitis)

When in contact with C. Diff, patient's hands must be washed with soap & water when performing hand hygiene

IV THERAPY: TYPES OF IV SOLUTIONS

HYPERTONIC

MEMORY TRICK: "ENTER THE VESSEL FROM THE CELLS"



MORE SALT in the solution, **LESS WATER** in the solution. The vessel becomes **MORE** concentrated than the cell. Water then **LEAVES** the cell. Therefore, the cells will **SHRINK**.

EXAMPLES:

- 5% saline
- 3% saline
- 5% dextrose in 0.9% saline (D5NS)
- 5% dextrose in 0.45% saline (D5 1/2 NS)
- 5% dextrose in LR (D5LR)
- 10% dextrose in water (D10W)

USED FOR:

- Cerebral edema
- Hyponatremia (low levels of sodium)
- Metabolic alkalosis
- Maintenance fluid
- Hypovolemia

DO NOT GIVE WITH:

- ↑ ICP
- Burns
- Trauma



HYPERtonic think HIGH numbers


*The only exception to this memory trick is 5% DEXTROSE IN WATER (D5W)

→ 5% DEXTROSE IN WATER (D5W) starts as ISOTONIC and then changes to HYPOTONIC when the dextrose is metabolized.

ISO means EQUAL

ISOTONIC

MEMORY TRICK: "STAYS WHERE I PUT IT"



Same osmolality as body fluids (Equal water & particle ratio)

EXAMPLES:

- 0.9% sodium chloride (NS) (normal saline)
- 5% dextrose in water (D5W)*
- Lactated Ringers (LR)

USED FOR:

- Blood loss (hemorrhage, burns, surgery)
- Dehydration (vomiting & diarrhea)
- Fluid maintenance

EXPANDS intravascular fluid volume & replaces fluid loss


Use with BLOOD PRODUCTS

NORMAL SALINE

is the only solution compatible to use with blood or blood products

HYPOTONIC

MEMORY TRICK: "GO OUT OF THE VESSEL" + INTO THE CELL



LESS SALT in the solution, **MORE WATER** in the solution. The vessel becomes **LESS** concentrated than the cell. Water then **ENTERS** the cell. Therefore, the cells will **SWELL**.

EXAMPLES:

- 0.45% saline (1/2 NS)
- 0.33% saline (1/3 NS)
- 0.225% saline (1/4 NS)
- 5% dextrose in water (D5W)*

USED FOR:

- Diabetic ketoacidosis (DKA)
- Helps kidneys excrete excess fluids
- Hypernatremia (high levels of sodium)

MONITOR FOR:

- Fluid Volume Overload

In DKA, there is so much glucose in the cells, they need water!

IV THERAPY: BASICS

Fluid in our body is found in **2** places:

INTRACELLULAR (ICF) & EXTRACELLULAR (ECF)

fluid **INSIDE** the cell

fluid **OUTSIDE** the cell


(Millions of these cells in our body)

INTERSTITIAL FLUID (ISF)

fluid that surrounds the cell
AKA fluid in the tissues

INTRAVASCULAR (IV)

plasma/fluid in
the blood vessels



THE CELLS & HOMEOSTASIS

The cells love to have everything equal (homeostasis).
But when fluids/solutes shift, **DIFFUSION/OSMOSIS**
occurs to get back to homeostasis again.

DIFFUSION

the movement of a **SOLUTE** from a **HIGHER** concentration to a **LOWER** concentration
(until there is equal concentration)

TIP

Sodium is a solute!

OSMOSIS

the movement of **WATER** through a semipermeable membrane from a **LOWER** solute concentration to a **HIGHER** solute concentration
(until there is equal concentration)

said another way...
from a **HIGHER** water concentration to a **LOWER** water concentration
(until there is equal concentration)

SODIUM & WATER



WHERE SODIUM GOES WATER FLOWS!

Sodium is the cool kid,
so water wants to be his friend.



EXAMPLE: If sodium shifts into the cell (intracellular space) water will follow and leave the extracellular space (the vessel)

COLLOIDS & CRYSTALLOIDS

COLLOIDS

Large molecules

Colloids have **LARGE** molecules making it more efficient at increasing fluid volume in the blood.

PLASMA EXPANDERS!

EXAMPLES:

Albumin
Fresh frozen plasma (FFP)

USED FOR:

Shock
Pancreatitis
Burns
Excessive bleeding

CRYSTALLOIDS

Small molecules

Crystalloids have **SMALL** molecules. They are less expensive than colloids and provide immediate fluid resuscitation.

EXAMPLES:

Hypertonic solution
Isotonic solution
Hypotonic solution

IV THERAPY: COMPLICATIONS

SYMPTOMS

- Tachycardia
- Chest pain
- Hypotension
- ↓ LOC
- Cyanosis

AIR EMBOLISM

Air enters the vein through the IV tubing

TREATMENT

- Clamp the tubing
- Turn client on the left side & place in Trendelenburg position
- Notify the HCP

SYMPTOMS

- At the site...
 - Pain
 - Swelling
 - Coolness
 - Numbness
- No blood return

INFILTRATION

IV fluid leaks into surrounding tissue

TREATMENT

- Remove the IV
- Elevate the extremity
- Apply a warm or cool compress
- Do not rub the area

SYMPTOMS

- Tachycardia
- Redness
- Swelling
- Chills & Fever
- Malaise
- Nausea & vomiting

INFECTION

Entry of microorganism into the body via IV

TREATMENT

- Remove the IV
- Obtain cultures
- Possible antibiotics administration

SYMPTOMS

- ↑ blood pressure
- Distended neck veins
- Dyspnea
- Wet cough & crackles

CIRCULATORY OVERLOAD

Administration of fluids too rapidly (Fluid Volume Overload)

TREATMENT

- ↓ flow rate (keep-vein-open rate)
- Elevate the head of the bed
- Keep the client warm
- Notify the HCP

SYMPTOMS

- At the site
 - Heat
 - Redness
 - Tenderness
- ↓ Flow of IV

PHLEBITIS

Inflammation of the vein (can lead to a clot (thrombophlebitis))

TREATMENT

- Remove the IV
- Notify the HCP
- Restart the IV on the opposite side

SYMPTOMS

- Ecchymosis
- At the site
 - Blood
 - Hard & painful lump

HEMATOMA


Collection of blood in the tissues

TREATMENT

- **ELEVATE** the extremity
- Apply Pressure & Ice

BLOOD TRANSFUSIONS

ADMINISTRATION OF BLOOD TRANSFUSION



1 Insert an IV line using a 16g*, 18g, or 20g IV needle

**commonly used for trauma patients*

2 Run it with normal saline (keep-vein-open-rate)

3 Begin the transfusion slowly

A The first 15 min is the ***MOST CRITICAL***
The RN must stay at bedside

B Vital signs are monitored every 30 minutes - 1 hour

C After 15 minutes the flow can be increased (unless a transfusion reaction has occurred)

4 Document the patient's tolerance to the administration of the blood product

If you use too small of a needle, like a 24 gauge needle when administering blood products, it will cause the blood to **LYSIS**.

FACTS ABOUT BLOOD TRANSFUSION

- Administered by the RN
- Only Normal Saline (NS) can be used in conjunction with blood
- Type & screen and a cross match are good for **72 HOURS**

Blood must be hung (started) within **30 MINUTES** from the time the blood is picked up from the blood bank

All blood must be transfused within **4 HOURS** of the time the blood was hung (started)

TRANSFUSION REACTION

tachycardia, itching, skin rash, wheezing, dyspnea, anxiety, flushing, fever, back pain



STOP the transfusion if you suspect a **TRANSFUSION REACTION**

TRANSFUSION REACTION

A transfusion reaction is an adverse reaction that happens as a result of receiving blood transfusions

IMMEDIATE TRANSFUSION REACTION

Chills, diaphoresis, aches, chest pain, rash, hives, itching, swelling, dyspnea, cough, wheezing, or rapid, thready pulse

CIRCULATORY OVERLOAD

Infusion of blood too rapid for the client to tolerate
Cough, dyspnea, chest pain, headache, hypertension, tachycardia, bounding pulse, distended neck vein, wheezing

SEPTICEMIA

Blood that is contaminated with microorganisms
Rapid onset of chills, high fever, vomiting, diarrhea, hypotension & shock

IRON OVERLOAD

Complication that occurs in client's who receive multiple blood transfusions
Vomiting, diarrhea, hypotension, altered hematological values

**Always check with your hospital's protocol about IV and blood product administration*

SIGNS OF TRANSFUSION REACTIONS

- Fast heart rate
- Itching/urticaria/skin rash
- Wheezing/dyspnea/tachypnea
- Anxiety
- Flushing / fever
- Back pain

NURSING ACTIONS TO A TRANSFUSION REACTION

- STOP** the transfusion
- Change the IV tubing down to the IV site
- Keep the IV open w/ normal saline
- Notify the HCP & blood bank
- Do not leave the client alone (monitor the client's vital signs & continue to assess the client)

PHARMACOKINETICS



"ADME" some medications

PHARMACOKINETICS:
The study of how drugs are moved throughout the body

A ABSORPTION

Medication going from the location of administration to the bloodstream

ORAL
Takes the longest to absorb

SUBCUT & IM
Depends on the site of blood perfusion.
More blood perfusion = rapid absorption

IV
Quickest absorption time

D DISTRIBUTION

Transportation by bodily fluids of the medication to where it needs to go


INFLUENCING FACTORS:

- Circulation
- Permeability of the cell membrane
- Plasma protein binding

M METABOLISM

How is the medication going to be broken down?

MOST COMMON SITE: LIVER




INFLUENCING FACTORS:

- Age
(Infants & elderly have a limited med-metabolizing capacity)
- Medication type
- First-pass effect
A drug given orally gets metabolized and its effects are greatly reduced before it reaches the systemic circulation. It's generally related to the liver or gut. It may need to be administered via parenteral route (subQ, IM, or IV) because this route bypasses the liver and gut.
- Nutritional status

E EXCRETION

How is the medication going to be eliminated from the body?

MOST COMMONLY DONE BY: KIDNEYS



INFLUENCING FACTORS:

- Kidney dysfunction
Leads to an increase in the duration and intensity of a medication response

If the kidneys aren't working/excreting waste, the medication will stay in the body which leads to toxic levels

MEDICATION ADMINISTRATION

6 RIGHTS OF MED ADMIN

RIGHT **CLIENT** 

RIGHT **TIME** 


RIGHT **DOSE** 


RIGHT **MED** 


RIGHT **ROUTE** 


RIGHT **DOCUMENTATION** 

TYPES OF ORDERS

 **ROUTINE**
Given on a regular schedule with or without a termination date.

 **SINGLE "ONE-TIME"**
Used for a single case.
Not a routine medication.

 **STAT**
Only for administration once and given immediately.

 **PRN**
"As needed" must have an indication for use such as pain, nausea & vomiting.

⚠️ COMMON ⚠️ MEDICATION ERRORS

 Medication error kills, prevention is crucial!

- ✗ Wrong medication
- ✗ Incorrect dose
- ✗ Wrong...
 - Client
 - Route
 - Time
- ✗ Administer a medication the client is allergic to
- ✗ Incorrect D/C of Medication
- ✗ Inaccurate prescribing

SCOPE OF PRACTICE



RN

- * Post-op assessment
- * Initial client teaching
- * Starting blood products
- * Sterile procedures
- * IVs & IV medications
- * Discharge education
- * Clinical assessment

NOTE:

When a registered nurse delegates tasks to others, responsibility is transferred but accountability for patient care is not transferred. The RN is still responsible!

LPN/LVN

- * Stable client
- * Monitor RN's findings & gather data
- * Specific assessments
- * Reinforce teaching
- * Routine procedures (catheterization, ostomy care, wound care)
- * Monitors IVFs & blood products
- * Administer injections & narcotics (not IVs meds & 1st IV bag)
- * Tube potency & enteral feedings
- * Sterile procedures

SPECIFIC ASSESSMENTS

Lung sounds, bowel sounds, & neurovascular checks



UAP

- * Routine, stable vital signs
- * Documenting input and output
- * Can get blood from the blood bank
- * Activities of daily living (ADLs)

ADLs

 **Feeding**
(not with aspiration risk) 

 **Positioning**

 **Ambulation**

 **Cleaning**

 **Linen change**

 **Hygiene care**


RN = Registered Nurse LPN = Licensed Practical Nurse LVN = Licensed Vocational Nurse UAP = Unlicensed Assistive Personnel

PARENTERAL ADMINISTRATION

Any route of administration that does not involve drug absorption through the GI tract

SLOWEST ABSORPTION

10-15°
ANGLE



INTRADERMAL (ID)

USES:

- TB testing
- Allergy sensitivities

NEEDLE SIZE:

25 - 27 gauge


USUAL SITE:

Inner forearm

Should form a "BLEB"

Normal to overweight clients

90°
ANGLE



Thin clients

45°
ANGLE



SUBCUTANEOUS (SUBQ)

USES:

non-irritating, water-soluble medication (insulin & heparin)

NEEDLE SIZE:

23 - 25 gauge


USUAL SITE:

- Abdomen
- Posterior upper arm
- Thigh

Giving a malnourished/thin client a medication at a 90° angle could lead to accidental intramuscular injury!

QUICKEST ABSORPTION

90°
ANGLE



INTRAMUSCULAR (IM)

Do not inject more than 3 mL (2 mL for the deltoid)
Divide larger volumes into two syringes & use two different sites

USES:

Irritating, solutions in oils, and aqueous suspensions

NEEDLE SIZE:


22 - 25 gauge

USUAL SITE:

- Deltoid
- Vastus lateralis
- Ventrogluteal

Use the Z TRACK METHOD

25°
ANGLE



INTRAVENOUS (IV)

USES:

Administering medications, fluids, & blood products

NEEDLE SIZE:

16 gauge: clients who have trauma
18 gauge: surgery & blood administration
22 - 24 gauge: children, older adults, & clients who have medical issues or are stable post-op

USUAL SITE:

- Hand
- Wrist
- Cubital fossa
- Foot
- Scalp

25° angle used when starting an IV

The smaller the gauge number, the larger the IV bore

GAUGES & IV USES

16 G

Trauma, surgery, rapid fluid administration (bolus)

18 G

Administering blood, rapid infusions (bolus), CT scans with IV dye

20 G*

Medications, routine therapies, IV fluids

22 G

IV fluids, medications

24 G

Pediatric patients, elderly patients, very fragile/small veins

*Some hospitals allow blood to be administered with 20 G

Always check with your hospital's protocol about IV and blood product administration




LARGEST

SMALLEST

NONPARENTERAL ADMINISTRATION

Absorbed into the system through the digestive tract

ORAL OR ENTERAL




- **CONTRAINDICATIONS:** vomiting, aspiration precautions/absence of a gag reflex, decreased LOC, difficulty swallowing
- Have client sit at 90 angle to help with swallowing
- **NEVER** crush enteric-coated or time-release medications
- Break or cut scored tablets only!

TRANSDERMAL



- Place the patch on a dry and clean area of skin (free of hair)
- Rotate the sites of the patch to prevent skin irritation
- Always take off the old patch before placing a new one on

INHALATION



- Rinse mouth after the use of steroids
- 20 - 30 seconds between puffs
- 2 - 5 minutes between different medications
- Use a spacer if possible to prevent thrush

SUBLINGUAL & BUCCAL




SUBLINGUAL: Under the tongue
BUCCAL: Between the cheek & the gum

→ ★ **Do not swallow!**

Keep the medication under the tongue (sublingual) or in between the cheek and gum (buccal) until it has completely absorbed



SUPPOSITORIES



RECTAL

- Lateral or Sims' position
- Use lubrication
- Insert beyond the internal sphincter
- Leave it in for 5 minutes

VAGINAL

- Supine with knees bent & feet flat on the bed, close to hips
- Insert the suppository along the posterior wall of the vagina (3 - 4 inches deep)
- Stay supine for at least 5 minutes

INSTALLATION (DROPS, OINTMENTS, SPRAYS)

EYES

- If there is dried section use a moisten sterile gauze and wipe from inner to outer canthus to prevent bacterial from entering the eye
- Have the client tilt their head back slightly
- Pull lower eye lid down gently to expose the conjunctival sac
- Hold the dropper 1-2 cm above the conjunctiva sac & drop medication directly into the sac
- Close eye lid & apply gentle pressure on the nasolacrimal duct for 30 - 60 seconds



EARS

- Have client tilt their head
- Warm the solution before adm. to prevent vertigo & dizziness
- **ADULTS:** pull ear **UP**ward & outward
- **< 3 YEARS OF AGE:** pull ear **DOWN** & back



NOSE

- Have client lie supine
- Do not blow nose for 5 min after drop instillation




INTEGUMENTARY (SKIN) OVERVIEW

Color changes are more difficult to notice in dark-skinned clients

INSPECTION OF THE SKIN

	DESCRIPTION	INDICATION	LOCATIONS
PALLOR	Loss of color	Lack of blood flow, anemia, shock	Face, conjunctiva, nail beds, palm, lips, mucous membranes
ERYTHEMA	Redness <i>can be blanchable or non-blanchable</i>	Inflammation, localized vasodilation, sun exposure, rash, hyperthermia	Skin (areas of trauma or pressure)
JAUNDICE	Yellow to orange	Liver-dysfunction	Skin, sclera, mucous membranes
CYANOSIS	Bluish	Hypoxia (not enough oxygen) or impaired venous return	Lips, mucous membranes, nail beds, skin

The best way to assess for **JAUNDICE** is to press gently on the forehead or nose. If the skin looks yellow where you applied pressure, it indicates jaundice.



PERIPHERAL CYANOSIS

Cyanosis of the peripherals (fingertips, palms, toes)
Rarely a life-threatening medical emergency



CENTRAL CYANOSIS

Cyanosis around the mouth, tongue or mucous membranes



⚠️ Medical emergency!



EDEMA is accumulation of excess fluid in the body's tissues that causes swelling of the skin

WEeping EDEMA


Areas that have pitting edema can leak fluid out directly from the skin

EDEMA CAN BE:

→ Non-pitting

→ Pitting

GRADING PITTING EDEMA




PITTING is when you press the edematous area for a few seconds and it dimples or pits

TYPES OF WOUND DRAINAGE

SEROUS	Clear, watery plasma.	
SANGUINEOUS	Bright red blood.	Indicates active bleeding
SEROSANGUINEOUS	Pale, pink, watery. Mixture of clear and red fluid.	
PURULENT	Thick, yellowish-green. Foul odor.	May indicate infection


PRIMARY LESION

Develops as a result of a disease process




MACULE

Flat discoloration of the skin <1 cm
Example: freckles




PAPULE

Solid, slightly elevated lesion <1cm
Example: moles



NODULE

Solid & elevated lesion >1cm
Example: lipomas



PUSTULE

Enclosed pus-filled cavity
Example: acne



WHEEL

Superficial, raised lesion
Example: allergic reactions




VESICLE

Elevated cavity containing clear fluid
Example: chickenpox, shingles


SECONDARY LESION

Result from a primary lesion or due to a client's actions (scratching, picking)




FISSURE

Linear crack/tear with abrupt edge
Example: anal fissures, athlete's foot



SCAR

Normal tissue is lost & replaced with connective tissue causing a scar
Example: healed area after surgery/injury



EROSION

Scooped-out, shallow depression
Example: severe pressure injuries



SCALE


Compact, flaky skin (silvery or white)
Example: psoriasis

PRESSURE INJURIES (ULCERS)

"DECUBITUS ULCER" "BED SORES"


WHAT IS A PRESSURE INJURY?

The break down of skin integrity due to unrelieved pressure




TYPE 1

- Skin is intact (unbroken)
- Nonblanchable redness
- Swollen tissue
- Darker skin → may appear blue / purple




TYPE 2

- Skin is NOT intact
- Partial thickness loss
- No fatty tissue is visible
- Superficial ulcer




TYPE 3

- Skin is NOT intact
- Full thickness SKIN loss
 - Damage to or necrosis of subQ tissue
 - No bone, muscle, or tendon exposed
- Ulcer extend down to the underlying fascia, but not through it
- Deep crater with or without tunneling




TYPE 4

- Skin is NOT intact
- Full thickness TISSUE loss
 - Destruction of tissue
 - Bone, muscle, or tendon exposed
- Deep pockets of infection & tunneling



DEEP TISSUE INJURY (DTI)

- Skin is intact (unbroken)
- Tissue beneath the surface is damaged
- Appears purple or dark red



UNSTAGEABLE

Stage cannot be determined due to eschar or slough covering the visibility of the wound

RISK FACTORS

- | | |
|-----------------------------------|-------------------------------|
| A GING SKIN | P OOR NUTRITION |
| V ASCULAR DISORDERS | R EDUCED RBCs (ANEMIA) |
| O BESITY | E DEMA |
| I MMOBILITY & INCONTINENCE | S ENSORY DEFICITS |
| D IABETES | S EDATION |
| S KIN FRICTION | |



MNEMONIC

"AVOIDS PRESS"

BRADEN SCALE

Asses your client's skin **EVERY** shift for pressure injuries using the Braden Scale!

- Looks at 6 categories
- **SENSORY PERCEPTION**
 - **MOISTURE**
 - **ACTIVITY**
 - **MOBILITY**
 - **NUTRITION**
 - **FRICTION & SHEAR**
- Interpretation
- **LOW RISK: 22 - 23**
 - **LESS RISK: 19 - 21**
 - **HIGH RISK: <18**

PREVENTION

RELIEVE PRESSURE

- Apply pressure relieving devices (overlays, specialty beds, air cushions, foam-padded seat cushions, etc.)
- Do not use donut-type devices or synthetic sheepskins!

PROPER NUTRITION

- ↑ protein intake
- Adequate hydration
- Possible enteral nutrition

protein promotes wound healing

SKIN HYGIENE

- Clean skin with mild soap
- Clean incontinent clients
- Do not scrub or rub bony prominences
- Barrier for incontinence
- Moisturizer for hydration

REPOSITIONING

- Turn/reposition your client every 2 hours while in the bed
- Lift, do not PULL
 - Pulling could cause shearing & friction from force

HYPOVOLEMIA VS. HYPERVOLEMIA

HYPOVOLEMIA


"LOW" "VOLUME" "IN THE BLOOD"

ALSO CALLED Dehydration • Fluid volume deficit (FVD) • Hypovolemic shock!

CAUSES

- Loss of fluid from ANYWHERE
 - Thoracentesis
 - Paracentesis
 - Hemorrhage
 - NG tube
 - Trauma
 - GI losses
 - Vomiting
 - Diarrhea
- Third spacing
 - Burns
 - Ascites
- Polyuria (peeing a lot)
 - Diabetes
 - Diuretics
 - Diabetes insipidus

Third spacing shifts the fluids from the **INTRAVASCULAR SPACE** (the vein) into the **INTERSTITIAL SPACE** (third space).

This causes a drop in the circulating blood volume 

SIGNS & SYMPTOMS

- Flat neck veins
- ↑ HR (weak & thready)
- ↑ Respirations
- ↑ Urine specific gravity
- ↓ BP
- ↓ CVP
- ↓ Weight
- ↓ Skin turgor
- ↓ Urine output
- Dry mucous membranes
- Thirst

LESS VOLUME = LESS PRESSURE

LABS **CONCENTRATED** (DEHYDRATED) MAKES THE # GO UP

- ↑ Urine specific gravity
- ↑ Serum sodium
- ↑ Hematocrit (%)
- ↑ BUN



NURSING CONSIDERATIONS / TREATMENT

- Fluid replacement
 - Fluids (PO or IV)
- Safety precautions
 - Risk for fall due to orthostatic hypotension
- Daily I&O + weights

monitor for fluid volume overload



HYPERVOLEMIA

"HIGH" "VOLUME" "IN THE BLOOD"

ALSO CALLED Over-hydration • Fluid volume excess


CAUSES

- Heart failure
 - Kidney dysfunction
 - Can't filter the blood = back up of fluids
- Cirrhosis
- ↑ Sodium intake



WHERE SODIUM GOES WATER FLOWS!


Sodium is the cool kid, so water wants to be his friend.



SIGNS & SYMPTOMS

- Distended neck vein (JVD)
- ↑ HR (bounding)
- ↑ BP
- ↑ Weight
- ↑ CVP
- Edema
- Wet lung sounds
 - Crackles / dyspnea
 - Due to back flow of fluid from the heart
- Polyuria
 - Kidneys are trying to get rid of the excess fluid

MORE VOLUME = MORE PRESSURE



LABS **DILUTED** (OVER-HYDRATED) MAKES THE # GO DOWN

- ↓ Urine specific gravity
- ↓ Serum sodium
- ↓ Hematocrit (%)
- ↓ BUN



NURSING CONSIDERATIONS / TREATMENT

- Low sodium diet
- Daily I&O + weights
- Diuretics
- High-Fowler's or Semi-Fowler's position
 - Easier to breathe



WHERE SODIUM GOES WATER FLOWS!



MENTAL HEALTH

BROUGHT TO YOU BY



THERAPEUTIC COMMUNICATION TECHNIQUES

Client-centered type of communication to build and help relationships with clients, families, and all relationships.

✓ DO

- Allow client to control the discussion
- Give recognition/validation
- Active listening!
- Use open-ended questions

Don't be a **LOSER**, be an active listener!

- L** Lean forward toward the client
- O** Open posture
- S** Sit squarely facing the client
- E** Establish eye contact
- R** Relax & listen

EXAMPLES

"Is there something you would like to talk about?"

"Tell me more about that"

"So you are saying you haven't been sleeping well?"

"Tell me more about ____"

✗ DON'T

- Ask "why"
- Ask too many questions
- Give advice
- Give false reassurance
- Change the conversation topic
- Give approval or disapproval
- Use close-ended questions/statements

EXAMPLES

"Don't worry!"

"I think you should ____"

"Don't be silly"

"That's great!"

THERAPEUTIC COMMUNICATION CAN BE BOTH...

VERBAL COMMUNICATIONS




Words a person speaks



NON-VERBAL COMMUNICATIONS



You may say all the "right" things but deliver it poorly.



Facial expressions

Eye contact

Posture

Vocal cues


(yawning, tone of voice, pitch of voice)

Movement

Appearance

Body language

PERSONALITY DISORDERS



NURSING CARE

- * **Safety** is a priority
- * Develop a therapeutic relationship
- * Respect the client's needs while still setting limits and consistency
- * Give the client choices to improve their feeling of control

Clients with a personality disorder are at a ↑ risk for violence & self-harm

TREATMENT

Medications such as:



- ➔ Antidepressants
- ➔ Anxiolytics
- ➔ Antipsychotics
- ➔ Mood stabilizers

Therapies such as:

- ➔ Psycho
- ➔ Group
- ➔ Cognitive
- ➔ Behavioral

*For more information about psychiatric medications, see the Pharmacology Bundle

EATING DISORDERS



ANOREXIA NERVOSA

- * ↓ Weight (BMI <18.5)
- * ↓ Blood pressure
- * ↓ Heart rate from dehydration & electrolyte imbalance
- * ↓ Sexual development
- * ↓ Subcutaneous tissue = Hypothermia
- * ↓ Period regularity
- * Amenorrhea (*period may stop*)
- * Refuses to eat
- * Lanugo (*thin hair to keep the body warm*)
- * Typically does not purge
- * Restricts self from eating
- * Fear of gaining weight
- * Constipation (*from dehydration*)

TREATMENT

- 👉 ↑ **WEIGHT SLOWLY**
(2 -3 lbs a week)
- 👉 **MONITOR EXERCISE**



BULIMIA NERVOSA

- * Binge eating followed by purging
- * Normal weight to overweight (BMI 18.5 - 30)
- * Teeth erosion
- * Bad breath
- * May use laxatives and/or diuretics

TREATMENT

Monitor client during and after meals for acts of purging



BINGE EATING

- * Binge eating not followed by purging
- * Tend to be overweight
- * Binging causes:
 - Depression
 - Hatred
 - Shame



REEFEEDING SYNDROME

Potential complications when fluids, electrolytes, and carbohydrates are introduced too quickly to a malnourished client. Treatment should be done **slowly** to avoid this syndrome.

TREATMENT FOR ALL EATING DISORDERS

Teach coping skills

Maintain trust


Have the client be a part of the decision making & the plan of care!

Therapy group, individual or family

BIPOLAR DISORDER

MOOD SWINGS:

Depression to mania with periods of normalcy



TREATMENT	
<p style="font-weight: bold; margin-top: 10px;">NURSING CONSIDERATIONS FOR THE ACUTE PHASE</p>	<ul style="list-style-type: none"> • Provide a safe environment <i>Remove harmful objects from the room</i> • Set limits on manipulative behavior • Provide finger foods & fluids • Re-channel energy for physical activity • ↓ Stimuli <ul style="list-style-type: none"> - Turn off or turn down the TV & music - Keep away from other clients if they are bothersome
<p style="font-weight: bold; margin-top: 10px;">PHARMACOLOGY</p>	<ul style="list-style-type: none"> • Lithium carbonate • Anticonvulsants • Antidepressants • Antipsychotics • Antianxiety <p style="text-align: right; font-size: small; margin-top: 10px;"><i>See pharmacology section for more details</i></p>

SCHIZOPHRENIA SPECTRUM DISORDER OVERVIEW

PHASES

1	PRE-MORBID	Normal functioning. Symptoms have not become apparent yet.
2	PRODROMAL	More tempered form of the disorder. Can be months to years for the disorder to become obvious.
3	SCHIZOPHRENIA	Positive symptoms are noticeable and apparent.
4	RESIDUAL	Periods of remission. Negative symptoms may remain, but S&S of the acute stage (positive symptoms) are gone.

POSSIBLE CAUSES

(not fully known)



↑ in the neurotransmitter **DOPAMINE**



ILLICIT SUBSTANCE
(LSD & Marijuana)



ENVIRONMENTAL
(malnutrition, toxins, viruses during pregnancy)



GENETICS
(family history)

SIGNS & SYMPTOMS

POSITIVE

- Delusions
- Anxiety/agitation
- Hallucinations
- Auditory** *most common
- Jumbled speech
- Disorganized behavior

NEGATIVE

- Flattened/bland effect
- Lack of energy
- Reduced speech
- Avolition**
Lack of motivation
- Anhedonia**
Not capable of feeling joy or pleasure
- Lack of social interaction

TREATMENT

- **Medication**
 - Antipsychotic medications
 - Antidepressants
 - Mood stabilizers (lithium)
 - Benzodiazepines

**For more information about psychiatric medications, see the Pharmacology Bundle*
- **Therapy**
- **Exercise**

NURSING CONSIDERATIONS

- 👉 Try to establish trust with the client
- 👉 Encourage compliance with the medications
- 👉 Promote self-care
- 👉 Encourage group activities
- 👉 Offer therapeutic communication

HOW TO ADDRESS HALLUCINATIONS?

- Don't address the hallucinations
- Be compassionate
- Bring the conversation back to reality
- Do not argue with the client
- Provide safety for the client & the staff!

EXAMPLE:

"I don't see spiders on the wall but I see you are scared"

TYPES OF DEPRESSION

MAJOR DEPRESSIVE DISORDER (MDD)

Has at least 5 of these symptoms every day for at least 2 weeks:

- Depressed mood
- Too much or too little sleep
- Indecisiveness
- Thoughts of death (suicide)
- ↓ ability to think/concentrate
- Not able to feel pleasure
- ↑ or ↓ motor activity
- Weight fluctuations
(5% change within a month)

FACTS

- MDD impairs the client's normal functioning
- MDD is not the same depression seen in bipolar disorder
- MDD is not a mood swing, it's constant

TREATMENT PHASES FOR MDD

ACUTE: 6 - 12 weeks

Hospitalization & medications may be prescribed

GOALS:

- ↓ Depressive symptoms
- ↑ Functionality

CONTINUATION: 4 - 9 months

Medication is continued

GOALS:

- Prevent relapse

MAINTENANCE: 1+ year

Medication may be continued or be phased out


GOALS:

- Prevent relapse & further depressive episodes

Treatment for the client will reflect what phase they are in!

PREMENSTRUAL DYSPHORIC DISORDER (PMDD)

Depression that occurs during the luteal phase of the menstrual cycle.



SYMPTOMS

- Emotional
- ↑ Eating
- ↓ Energy
- ↓ Concentration

SUBSTANCE INDUCED DEPRESSIVE DISORDER

Depression associated with withdrawal or the use of alcohol and drugs.



PERSISTENT DEPRESSIVE DISORDER (DYSTHYMIA)

A more mild form of depression compared to MDD, although it can turn into MDD later in life.



SEASONAL AFFECTIVE DISORDER (SAD)

Depression that occurs seasonally. Often occurs during the winter months when there is less sunshine.



TREATMENT: Light therapy

POSTPARTUM

Depression that happens after a woman goes through childbirth. The woman may feel disconnected from the world. She may have a fear of harming her newborn.



NURSING CONSIDERATIONS

- Safety is a priority. Those struggling with depression have a higher suicide risk.
 - INITIATE SUICIDE PRECAUTIONS:**
 - Remove sharp things
 - Keep medications out of reach
 - Remove objects that may be used for strangulation (wires)
- Help the client identify coping methods & teach alternatives if needed
- Provide local resources such as churches, local programs, community resources, etc.
- **ENCOURAGE:**
 - Physical activity
 - Self-care
 - Supportive relationships
 - Individual therapy, support groups, & peer support*

TREATMENT

ANTIDEPRESSANTS

- SSRIs
- SNRIs
- TCAs
- MAOIs

NON-PHARMACOLOGICAL THERAPIES

- Light therapy
- St. John's wort

ELECTROCONVULSIVE THERAPY (ECT)

Used for clients who are unresponsive to other treatments. Transmits a brief electrical stimulation to the patient's brain.

THE PROCEDURE

- The client is asleep under anesthesia
- The client will not remember and is unaware of the procedure
- Muscle relaxants may be given to ↓ seizure activity & ↓ risk for injury
- Client may have memory loss, confusion, & headache post-procedure



*For more information about antidepressants, see the psychiatric section in the Pharmacology Bundle

DIFFERENT TYPES OF ANXIETY DISORDERS

NORMAL

WORST

	MILD	MODERATE	SEVERE	PANIC
LEVELS OF ANXIETY	Normal/healthy amount of anxiety. Allows one to have sharp focus & problem solve.	Thinking ability is impaired. Sharp focus & problem-solving can still happen just at a lower level.	Focus & problem solving are not possible. Feelings of doom may be felt.	Most extreme anxiety. Unstable & not in touch with reality.
SYMPTOMS	Nail-biting Tapping Foot jitters	GI upset Headache Voice is shaky	Dizziness Headache Nausea Sleeplessness Hyperventilation	Pacing Yelling Running Hallucinations

ANXIETY DISORDERS	
Separation Anxiety Disorder	Experiences extreme fear of anxiety when separated from someone they are emotionally connected to. This is a normal part of infancy, but not a normal part of adulthood.
Specific Phobia	Irrational fear of a particular object or situation. SOME EXAMPLES: <ul style="list-style-type: none"> • Monophobia - Fear of being alone • Zoophobia - Fear of animals • Acrophobia - Fear of heights
Social Anxiety Disorder (Social Phobia)	Fear of social situations or presenting in front of groups. They fear embarrassment. They may have symptoms (real or fake) to escape the situation.
Panic Disorder	Reoccurring panic attacks that last 15 - 30 minutes with physical manifestations.
Agoraphobia	Extreme fear of certain places where the client feels unsafe or defenseless. May even be too fearful of places to maintain employment.
Generalized Anxiety Disorder (GAD)	Uncontrolled extreme worry for at least 6 months that causes impairment of functionality.

AGORA MEANS "open space"

OBSESSIVE COMPULSIVE DISORDERS	
Obsessive Compulsive Disorder (OCD)	<p>OBSSESSION: Recurrent thoughts</p> <p>COMPULSION: Recurrent acts or behaviors</p> <p>This obsessiveness is usually because it decreases stress & helps deal with anxiety.</p>
Hoarding Disorder	Compulsive desire to save items even if they have no value to the person. It may even lead to unsafe living environments.
Body Dysmorphic Disorder	Preoccupied with perceived flaws or imperfections in physical appearance that the client thinks they have.

SOMATIC SYMPTOM & RELATED DISORDERS (Somatoform Disorders)

SOMATIC SYMPTOM DISORDER

Somatization is psychological stress that presents through physical symptoms that can not be explained by any pathology or diagnosis.

NURSING CONSIDERATIONS

- SAFETY is a priority Asses for symptoms or thoughts of self-harm or suicide
- Understand the somatic symptoms are real to the client even though they are not real
- Help the client verbalize their feelings while limiting the amount of time talking about their somatic symptoms
- Assess coping mechanism & educate on alternative ways of coping

MANIFESTATIONS

- Consumed by physical manifestations to the point it disrupts daily life
- Seeks medical help from multiple places
- Remission & exacerbations
- Overmedicates with analgesic and antianxiety medications
- ↑ Stress = ↑ somatic symptoms



PHQ-15: PATIENT HEALTH QUESTIONNAIRE 15

An assessment tool used to identify 15 of the most common somatic symptoms

CONVERSION DISORDER

Sudden onset of neurological manifestations & physical symptoms without a known neurological diagnosis. It can be related to a psychological conflict/need beyond their conscious control.

NURSING CONSIDERATIONS

- Ensure SAFETY
- Gain trust & rapport with the client
- Assess coping mechanism & educate on alternative ways of coping
- Assess stress management methods
- Encourage therapy such as:
 - Individual therapies
 - Group therapies
 - Support groups

MANIFESTATIONS

MOTOR

Paralysis pseudoseizures

Pseudocyesis:

Signs & symptoms of pregnancy without the presence of a fetus AKA false pregnancy. This may be present in a client who desires to become pregnant


SENSORY

Blindness

Deafness

Sensations (burning/tingling)

Inability to smell/speak



MEDICATIONS

The client may be prescribed **antidepressants** or **anxiolytics**

POSTTRAUMATIC STRESS DISORDER (PTSD)

Mental health condition where exposure to a traumatic event has occurred.

NURSING CONSIDERATIONS

- Teach relaxation techniques
- Teach ways to ↓ anxiety
- Support groups

MANIFESTATIONS

Lasting longer than 1 month:

- Anxiety
- Detachment
- Nightmares of the event




MEDICATIONS

Antidepressants may be prescribed

*For more information about antidepressants, see the psychiatric section in the Pharmacology Bundle

NEUROCOGNITIVE DISORDERS



Dementia & Alzheimer's are NOT the same.

Dementia is a general term that refers to a group of symptoms, not a specific disease. Dementia may advance to a major neurocognitive disorder such as Alzheimer's disease.

DELIRIUM

SHORT TERM / SUDDEN CHANGE

Impairment (hours - days)

There is always an underlying cause... something is causing the delirium!

- Hospitalization
- ICU delirium
- Polypharmacy
- Old age
- Stroke
- Surgery
- Restraints
- Secondary to a medical condition (infection, electrolyte imbalance, substance abuse...etc)

Delirium is a medical emergency and requires prompt diagnosis & treatment

- Disorganization
 - Most common to time & place
 - Happens mostly at night
- ↓ Memory
- Anxiety & agitation
- Delusional thinking
- Ranges from lethargic to hypervigilance!

- Safety: prevent physical harm
- Avoid restrains when possible
- Remember physical needs (Hygiene, food, water, sleep, etc)
- May be prescribed anti-anxiety/antipsychotic medications

ONSET

RISK FACTORS

MANIFESTATIONS

INTERVENTIONS

CURE?

ALZHEIMER'S

CONTINUOUS

Decline of function (months - years)

Genetics

Family history (immediate family)

Head Injury

Traumatic brain injuries (TBI) & head trauma

Advanced Age

>65 have the highest risk

Cardiovascular Disease & Lifestyle Factors

Inactivity, unhealthy diet, high cholesterol, obesity, & diabetes

STAGES OF ALZHEIMER'S DISEASE

SEVERE	LATE STAGE <i>Requires full assistance</i>	• Needs assistant with all ADL's	• Losing the capability to have discussions	• Losing physical skills (walking, sitting, swallowing)	• May result in death or coma
		MODERATE	MIDDLE STAGE <i>noticeable to others</i>	• Forgets own history	• Gets lost & wanders often
				• Difficulty completing tasks	• Gets angry & frustrated
MILD	EARLY STAGE <i>not noticeable to others</i>	• Memory lapse	• Misplacing things	• Short term memory	• Difficulty focusing
		• Can still accomplish own ADL's			

Caring for a client with Alzheimer's is very complex!

- Help families in planning for extended care
- Monitor nutrition, weight, & fluids status
- Maintain a quiet environment to ↓ stimuli
- **Cholinesterase inhibitor** may be prescribed to improve quality of life but does NOT cure the disease.

Communication

- Speak slowly
- Give one direction at a time
- Don't ask complex or open-ended questions
- Ask simple, direct questions
- Face the client directly when speaking

USES



Used in early & moderate stages of dementia & Alzheimer's disease. May also be used for Parkinson's dementia.

GENERIC	TRADE NAME
donepezil	Aricept
galantamine	Razadyne
rivastigmine	Exelon

Reversible if prompt treatment is initiated

Irreversible



MOTHER BABY

BROUGHT TO YOU BY



ABBREVIATIONS

IUP/IUFD Intrauterine pregnancy / intrauterine fetal demise
SAB Spontaneous abortion
TAB Therapeutic abortion
LMP Last menstrual period
ROM Rupture of membranes
SROM Spontaneous rupture of membranes
AROM Artificial rupture of membranes
PROM Prolonged rupture of membranes (>24 hours)
PPROM Preterm premature rupture of membranes
SVD Spontaneous vaginal delivery
FHR Fetal heart rate
EFM Electronic fetal monitoring
US Ultrasound transducer (detects FHR)
FSE Fetal scalp electrode (precise reading of FHR)
IUPC Intrauterine pressure catheter (strength of contractions)
LTV Long term variability
SVE Sterile vaginal exam
MLE Midline episiotomy

NST Non-stress test
CST Contraction stress test
BPP Biophysical profile
VBAC Vaginal birth after cesarean
AFI Amniotic fluid index
BUFA Baby up for adoption
NPNC No prenatal care
PTL Preterm labor
BOA Born on arrival
BTL Bilateral tubal ligation
D&C / D&E ... Dilation & curettage / dilation & evacuation
LPNC Late prenatal care
TIUP Term intrauterine pregnancy
VMI / VFI Viable male infant / viable female infant
EDB Estimated date of birth
EDC Estimated date of confinement
EDD Estimated date of delivery

PREGNANCY DURATION

40 WEEKS GESTATIONAL AGE

The number of completed weeks counting from the 1st day of the last normal menstrual cycle (LMP).

38 WEEKS FETAL AGE

This refers to the age of the developing baby, counting from the estimated date of conception. The fetal age is usually 2 weeks less than the gestational age.

TRIMESTERS

FIRST TRIMESTER

0 – 13 WEEKS

SECOND TRIMESTER

14 – 26 WEEKS

THIRD TRIMESTER

27 – 40 WEEKS

PRENATAL TERMS

Gravida / Gravidity

A woman who is pregnant / the number of pregnancies

NULLIGRAVIDA

Never been pregnant

PRIMIGRAVIDA

Pregnant for the first time

MULTIGRAVIDA

A woman who has had 2+ pregnancies

Parity

The number of pregnancies that have reach viability (20 weeks of gestation) whether the fetus was born alive or not

NULLIPARA

0

Zero pregnancies beyond viability (20 weeks)

PRIMIPARA

1

One pregnancy that has reached viability (20 weeks)

MULTIPARA

2+

Two or more pregnancies that have reached viability (20 weeks)

PRETERM

Pregnancies that have reached 20 weeks but ended before 37 weeks

TERM

Pregnancies that have lasted between week 37 and week 42

EARLY TERM: 37 – 38 6/7

FULL TERM: 39 – 40 6/7

LATE TERM: 41 – 41 6/7

POSTDATE/POSTTERM

A pregnancy that goes beyond 42 weeks

GTPAL

An acronym used to assess pregnancy outcomes

G

GRAVIDITY



The number of pregnancies

- Includes the present pregnancy
- Includes miscarriages / abortions
- Twins / triplets count as one

T

TERM BIRTHS



The number born at term

- > 37th week of gestation
- Includes alive or stillborn
- Twins / triplets count as one

P

PRE-TERM BIRTHS



The number of pregnancies delivered beginning with the 20th - 36 ⁶/₇th weeks of gestation

- Includes alive or stillborn
- Twins / triplets count as one

A

ABORTIONS / MISCARRIAGES



The number of pregnancies delivered before 20 weeks gestation

- Counts with gravidity
- Twins / triplets count as one

L

LIVING CHILDREN



The number of current living children

- Twin / triplets count individually

ANSWER KEY

Q#2 is (C) 4-2-1-0-4
Q#1 is (D) 3-2-0-1-2

PRACTICE QUESTION 1

You are admitting a client to the mother-baby unit. Two hours ago she delivered a boy on her due date. She gives her obstetric history as follows: she has a three-year-old daughter who was delivered a week past her due date and last year she had a miscarriage at 8 weeks gestation. How would you note this history using the GTPAL system?

- A. 2-2-1-0-2
- B. 3-2-1-0-1
- C. 3-2-1-0-2
- D. 3-2-0-1-2

PRACTICE QUESTION 2

A prenatal client's obstetric history indicates that she has been pregnant 3 times previously and that all her children from previous pregnancies are living. One was born at 39 weeks gestation, twins were born at 34 weeks gestation, & another child was born at 38 weeks gestation. She is currently 38 weeks pregnant. What is her gravidity & parity using the GTPAL system?

- A. 4-1-3-0-4
- B. 4-1-2-0-3
- C. 4-2-1-0-4
- D. 4-2-2-0-4

PREGNANCY SIGNS & SYMPTOMS

PRESUMPTIVE

SUBJECTIVE

Think
"Mom"

These are changes felt by the women that are subjective.
Can be associated with other things.

NOT a definite diagnosis for pregnancy!

- P** Period Absent (Amenorrhea)
- R** Really tired
- E** Enlarged breasts
- S** Sore breasts
- U** Urination increased (urinary frequency)
- M** Movement perceived (quickening)
- E** Emesis & nausea

Why is quickening not a positive sign?

Quickening can be difficult to distinguish from peristalsis or gas so it can not be a positive sign.

PROBABLE

OBJECTIVE

Think
"Doctor"

Pregnancy signs that the nurse or doctor can observe

- P** Positive (+) pregnancy test (high levels of the hormone: hCG)
- R** Returning of the fetus when uterus is pushed w/ fingers (ballottement)
- O** Objective
- B** Braxton hicks contractions
- A** A softened cervix (Goodell's sign)
- B** Bluish color of the vulva, vagina, or cervix (Chadwick's sign)
- L** Lower uterine segment soft (Hegar's sign)
- E** Enlarged uterus

Why is a positive pregnancy test not a positive sign?

High levels of hCG can be associated with other conditions such as certain medications or hydatidiform mole (molar pregnancy).



POSITIVE

OBJECTIVE

Think
"Baby"

Can only be attributed to a fetus

Definite diagnosis for pregnancy!

- F** Fetal movement palpated by a doctor or nurse
- E** Electronic device detects heart tones ♥
- T** The delivery of the baby
- U** Ultrasound detects baby
- S** Seeing visible movements

PREGNANCY PHYSIOLOGY

HORMONES

- Prolactin:** Allows for breast milk production
- Estrogen:** Growth of fetal organs & maternal tissues
- Progesterone & Relaxin:** Relaxes smooth muscles
- hCG:** Produced by placenta, prevents menstruation
- Oxytocin:** Stimulates contractions at the start of labor

RESPIRATORY

- ↑ Basal metabolic rate (BMR)
- ↑ O₂ needs
- Respiratory alkalosis (MILD)

CARDIOVASCULAR

- ↑ Cardiac output (↑ Heart rate + ↑ stroke volume)
- Blood pressure stays the same or a slight decrease
- ↑ in plasma volume
- ♥ Enlarges (May develop systolic murmurs)

Blood pressure should not be increased! This could indicate preeclampsia

RENAL

- ↑ GFR from ↑ plasma volume
- Smooth muscle relaxation of the uterus = ↑ risk of UTI's!
- ↑ Urgency, frequency & nocturia
- EDEMA!!

SKIN

- **Striae**
Stretch marks (abdomen, breasts, hips, etc)
- **Chloasma**
Mask of pregnancy
Brownish hyperpigmentation of the skin
- **Linea Nigra**
"Pregnancy line" dark line that develops across your belly during pregnancy
- **Montgomery glands / Tubercles**
Small rough / nodular / pimple-like appearance of the areola (nipple)

MUSCULOSKELETAL

- **Lordosis:** center of gravity shifts forward leading to inward curve of spine
- Low back pain
- Carpal tunnel syndrome
- Calf cramps

PITUITARY

- ↓ FSH/LH due to ↑ Progesterone
- ↑ Prolactin
- ↑ Oxytocin

THYROID

- ↑ Thyroxine
- May have moderate enlargement of the thyroid gland (goiter)
- ↑ Metabolism & ↑ appetite

GASTROINTESTINAL

- **Pyrosis**
↑ Progesterone = LOS to relax = ↑ heartburn
- **Constipation & hemorrhoids**
↑ Progesterone = ↓ gut motility
- **Pica**
Non-food cravings such as ice, clay, and laundry starch

HEMATOLOGICAL


FIBRINOGEN < Non-pregnant levels: 200-400 mg/dL
Pregnant levels: up to 600 mg/dL

Pregnant women are **HYPERCOAGULABLE** (increased risk for DVTs)

- ↑ White blood cells
- ↓ Platelets

ANEMIA

Plasma volume is greater than the amount of red blood cell (RBC) = hemodilution = **physiological anemia**



NAEGELE'S RULE

Used for estimating the expected date of delivery (EDD) based on LMP (last menstrual period)

DATE OF LAST MENSTRUAL PERIOD — 3 CALENDAR MONTHS + 7 DAYS + 1 YEAR



REMEMBER:
How many days are
in each month?

"30 days hath September,
April, June & November.
All the rest have 31, except
February alone (28 days) "

EXAMPLE

1st day of last period: September 2, 2015
Minus 3 calendar months: June 2, 2015
Plus 7 days: June 9, 2015
Plus 1 year: June 9, 2016 (EDD)

FACTS ABOUT NAEGELE'S RULE

- ♥ Bases calculation on a woman who has a 28-day cycle (most women vary)
- ♥ The typical gestation period is 280 days (40 weeks)
- ♥ First-time mothers usually have a slightly longer gestation period

WHAT TO AVOID DURING PREGNANCY

TERATOGENIC DRUGS



MNEMONIC

TERA-TOWAS

- T** Thalidomide
- E** Epileptic medications (valproic acid, phenytoin)
- R** Retinoid (vit A)
- A** Ace inhibitors, ARBS
- T** Third element (lithium)
- O** Oral contraceptives
- W** Warfarin (coumadin)
- A** Alcohol
- S** Sulfonamides & sulfones

TORCH INFECTIONS

TORCH infections are a group of infections that cause fetal abnormalities. Pregnant women should avoid these infections!



MNEMONIC

TORCH



- T** Toxoplasmosis
- Parv **O** Virus-B19 (fifth disease)
- R** Rubella
- C** Cytomegalovirus
- H** Herpes simplex virus

STAGES OF LABOR

STAGE 1

CERVIX **DILATES** FROM 0-10 CM

LONGEST STAGE

Latent (early)

- ♥ Cervix dilates: 1 - 3 cm
- ♥ Intensity: Mild
- ♥ Contractions: 15 - 30 mins

Active

- ♥ Cervix dilates: 4 - 7 cm
- ♥ Intensity: Moderate
- ♥ Contractions: 3 - 5 min (30-60 sec in duration)

Transition

- ♥ Cervix dilates: 8 - 10 cm
- ♥ Intensity: Strong
- ♥ Contractions: Every 2-3 min (60-90 sec in duration)

INTERVENTIONS

- ♥ Promote comfort
 - Warm shower, massage, or epidural
- ♥ Offer fluids & ice chips
- ♥ Provide a quiet environment
- ♥ Encourage voiding every 1 - 2 hours
- ♥ Encourage participation in care & keep informed
- ♥ Instruct partner in **effleurage** (light stroking of the abdomen)
- ♥ Encourage effective breathing patterns & rest between contractions



Labor Actively Transitioning

>30 min = Retained placenta

STAGE 2

THE **BABY** IS DELIVERED

- Starts when cervix is fully dilated & effaced
- Ends after the baby is delivered

PUSHING!

INTERVENTIONS

- ♥ Provide ice chips & ointment for dry lips
- ♥ Provide praise & encouragement to the mother
- ♥ Monitor uterine contractions & mother's vital signs
- ♥ Maintain privacy & encourage rest between contractions
- ♥ Encourage effective breathing patterns & rest between contractions
- ♥ Monitor for signs of birth (perineal bulging or visualization of fetal head)

STAGE 3

THE **PLACENTA** IS DELIVERED

The **PLACENTA** is expelled (5 - 30 min after birth)

SIGNS OF A PLACENTA DELIVERY


DELIVERY MECHANICS

- ♥ Lengthening umbilical cord
- ♥ Gush of blood
- ♥ Uterus changes from oval to globular shape

- "**Shiny Schultz**"
Side of *baby* delivered 1st
- "**Dirty Duncan**"
Side of *mother* delivered 1st

INTERVENTIONS

- ♥ Assessing mother's vital signs
- ♥ Uterine status (fundal rubs every 15 minutes)
- ♥ Provide warmth to the mother
- ♥ Promote parental-neonatal attachment
- ♥ Examine placenta & verify it's intact
 - Should have 2 arteries & 1 vein



STAGE 4

RECOVERY!

RECOVERY: first 1-4 hours after delivery of the placenta

- ♥ Assessing the fundus
- ♥ Continue to monitor vital signs & temperature for infection
- ♥ Administer IV fluids
- ♥ Monitor lochia discharge (lochia may be moderate in amount & red).
- ♥ Monitor for respiratory depression, vomiting, & aspiration if general anesthesia was used
- ♥ **Great time to watch for complications such as bleeding (postpartum hemorrhage)**



- ♥ FIRM
- ♥ Midline





- ♥ Soft
- ♥ Boggy
- ♥ Displaced



looks like a smiley face!

2 "A" for **ARTERIES**
1 "V" for **VEIN**

TRUE VS. FALSE LABOR

	FALSE LABOR	TRUE LABOR
CONTRACTIONS	<ul style="list-style-type: none"> • Irregular • Stops with walking / position change • Felt in the back or the abdomen above the umbilicus • Often stops with comfort measures 	<ul style="list-style-type: none"> • Occur regularly <ul style="list-style-type: none"> ▪ Stronger ▪ Longer ▪ Closer together • More intense with walking • Felt in lower back → radiating to the lower portion of the abdomen • Continue despite the use of comfort measures
CERVIX	<ul style="list-style-type: none"> • May be soft • NO significant change in.... <ul style="list-style-type: none"> ▪ Effacement ▪ Dilation • No bloody show • In posterior position (baby's head facing mom's front of belly) 	<ul style="list-style-type: none"> • Progressive change <ul style="list-style-type: none"> ▪ Softening ▪ Effacement ▪ Dilation signaled by the appearance of bloody show ▪ Moves to an increasingly anterior position (baby's head facing mom's back) 
FETUS	<ul style="list-style-type: none"> • Presenting part is usually not engaged in the pelvis 	<ul style="list-style-type: none"> • Presenting parts become engaged in the pelvis • Increased ease of breathing (more room to breathe) • Presenting part presses downward & compresses the bladder = urinary frequency

SIGNS OF LABOR

LABOR

Moving the fetus, placenta, & the membranes out of the uterus through the birth canal

Signs of Preceding Labor

- ☞ Lightening
- ☞ Increased vaginal discharge (bloody show)
- ☞ Return of urinary frequency
- ☞ Cervical ripening
- ☞ Rupture of membranes "water breaking"
- ☞ Persistent backache
- ☞ Stronger Braxton Hicks contractions
- ☞ Days preceding labor
 - Surge of energy
 - Weight loss (1- 3.5 pounds) from a fluid shift

FETAL HEART TONES


EARLY DECELERATIONS



"Mirror" image of mom's contractions
(They don't technically come early)

NORMAL FETAL HEART RATE
120 - 160 BPM

NORMAL! ✓




CAUSE:

- ♥ From head compression

INTERVENTION:

- ♥ Continue to monitor
- ♥ No intervention needed




LATE DECELERATIONS



Literally comes late after mom's contraction

NON-REASSURING ✗




CAUSE:

- ♥ Uteroplacental insufficiency

INTERVENTION:

- ♥ D/C oxytocin
- ♥ Position change
- ♥ Oxygen (nonrebreather)
- ♥ Hydration (IV fluids)
- ♥ Elevate legs to correct the hypotension




VARIABLE DECELERATIONS



*Variable: Looks "V" shaped

NON-REASSURING ✗




CAUSE:

- ♥ Cord compression

INTERVENTION:

- ♥ D/C Oxytocin
- ♥ Amnioinfusion
- ♥ Position change
- ♥ Breathing techniques
- ♥ Oxygen (nonrebreather)

Side-lying or knee chest will relieve pressure on cord




PREECLAMPSIA OVERVIEW

Overview of Hypertensive disorders during pregnancy

WHAT IS HYPERTENSION?

SYSTOLIC > 140
OR
DIASTOLIC > 90

Hypertension may be abbreviated "HTN"



SIGNS & SYMPTOMS

"PRE" eclampsia

- P** Proteinuria
 - R** Rising BP
 - E** Edema
- Triad signs

- ♥ Severe headache
- ♥ RUQ or epigastric pain
- ♥ Visual disturbances
- ♥ ↓ Urine output
- ♥ Hyperreflexia
- ♥ Rapid weight gain

PATHOLOGY

Pathology isn't completely known

PLACENTA is the root cause

- ♥ Defective spiral artery remodeling
- ♥ Systemic vasoconstriction & endothelial dysfunction

RISK FACTORS

- ♥ HX of preeclampsia in previous pregnancies
- ♥ Family history of preeclampsia
- ♥ 1st pregnancy AMA (advanced maternal age)
- ♥ Obesity
- ♥ Very young (<18) or very old (>35)
- ♥ Medical conditions (Chronic HTN, renal disease, diabetes, autoimmune disease)

HELLP SYNDROME

Variant of preeclampsia
Life-threatening complication

- H** Hemolysis
- EL** Elevated liver enzymes
- LP** Low platelet count

ECLAMPSIA

(seizures activity or a coma)

IMMEDIATE CARE:

- Side-lying
- Padded side rails with pillows/blankets
- O₂
- Suction if needed
- Do not restrain
- Do not leave



MAGNESIUM SULFATE

RX given to prevent seizures during & after labor.

***Remember:** magnesium acts like a depressant

THERAPEUTIC RANGE: 4 – 7 mg/dL

TOXICITY!

- RR <12
- ↓ DTR's
- UOP <30 mL/hr
- EKG Changes


*Mag is excreted in urine
↓UOP → ↑Mag levels

ANTIDOTE: CALCIUM GLUCONATE

*because magnesium sulfate can cause respiratory depression

VEAL CHOP

A tool to help interpret fetal strips



ASSESSMENT OF UTERINE CONTRACTIONS

DURATION	<p>BEGINNING of the contraction to the END of that same contraction</p> <ul style="list-style-type: none"> • Lasts 45 - 80 seconds • Should not exceed 90 seconds <p><i>Only measured through external monitoring</i></p>
FREQUENCY	<p>Number of contractions from the BEGINNING of one contraction to the BEGINNING of the next</p> <ul style="list-style-type: none"> • 2 - 5 contractions every 20 minutes • Should not be more FREQUENT than every 2 minutes <p><i>Only measured through external monitoring</i></p>
INTENSITY	<p>Strength of a contraction at its PEAK</p> <ul style="list-style-type: none"> • 25 - 50 mm Hg • Should not exceed 80 mm HG <p><i>Can be palpated</i></p> <p>MILD - nose MODERATE - chin STRONG - forehead</p>
RESTING TONE	<p>TENSION in the uterine muscle between contractions (relaxation of the uterus = fetal oxygenation between contractions)</p> <ul style="list-style-type: none"> • Average: 10 mm HG • Should not exceed 20 mm HG <p><i>Can be palpated</i></p> <p>SOFT = good FIRM = not resting enough</p>

LABOR & BIRTH PROCESSES

5 P's 5 factors that affect the process of labor & birth

PASSENGER

FETUS & PLACENTA

PASSAGEWAY

THE BIRTH CANAL

POSITION

POSITION OF THE MOTHER

POWERS

CONTRACTIONS

PSYCHOLOGY

EMOTIONAL RESPONSE

PASSENGER

FETUS & PLACENTA


SIZE OF THE FETAL HEAD

FONTANELS

- Space between the bones of the skull allows for molding
- Anterior (larger)
 - Diamond-shaped
 - Ossifies in 12-18 months
- Posterior
 - Triangle shaped
 - Closes 8 - 12 weeks

MOLDING


- Change in the shape of the fetal skull to "mold" & fit through the birth canal



FETAL PRESENTATION

Refers to the part of the fetus that enters the pelvic inlet first through the birth canal during labor

- 1 CEPHALIC** MOST COMMON
 - Head first
 - Presenting part: Occipital (back of head/skull)



- 2 BREECH**
 - Buttocks, feet, or both first
 - Presenting part: Sacrum



- 3 SHOULDER**
 - Shoulders first
 - Presenting part: Scapula



FETAL LIE

Relation of the long axis (spine) of the fetus to the long axis (spine) of the mother

LONGITUDINAL OR VERTICAL

- The long axis of the fetus is parallel with the long axis of the mother
- Longitudinal: cephalic or breech

TRANSVERSE, HORIZONTAL, OR OBLIQUE

- Long axis of the fetus is at a right angle to the long axis of the mother
- Transverse: vaginal birth **CANNOT** occur in this position
- Oblique: usually converts to a longitudinal or transverse lie during labor

CONTINUED →

LABOR & BIRTH PROCESSES

PASSENGER

CONTINUED

FETAL ATTITUDE

GENERAL FLEXION

- Back of the fetus is rounded so that the chin is flexed on the chest, thighs are flexed on the abdomen, legs are flexed at the knees

BIPARIETAL DIAMETER

- 9.25 cm at term, the largest transverse diameter and an important indicator of fetal head size

SUBOCCIPITOBREGMATIC DIAMETER

- Most critical & smallest of the anteroposterior diameters

LIGHTENING
When the baby "drops" into the mother's pelvis

FETAL POSITION


FETAL STATION

- Where the baby's **PRESENTING PART** is located in the pelvis
 - Measured in centimeters (cm)
 - Find the ischial **spine = ZERO**
 - **Above the ischial spine is (-)**
 - **Below the ischial spine is (+)**
- +4 / +5 = Birth is about to happen**

Head, foot, butt (closest to exit of uterus)



I'm (+) that I'm getting this baby out




ENGAGEMENT

- Fetal station **ZERO** = baby is "engaged"
- Presenting parts have entered down into the pelvis inlet & is at the ischial spine line (0)

When does this happen?

- **FIRST-TIME MOMS:** 38 weeks
- **ALREADY HAD BABIES:** can happen when labor starts



PASSAGEWAY

THE BIRTH CANAL: Rigid bony pelvis, soft tissue of cervix, pelvic floor, vagina & introitus

TYPES OF PELVIS



GYNECOID

- Classic female type
- Most common



ANDROID

- Resembling the male pelvis



ANTHROPOID

- Oval-shaped
- Wider anteroposterior diameter



PLATYPELLOID

- The flat pelvis
- Least common

SOFT TISSUE

LOWER UTERINE SEGMENT

- Stretchy

CERVIX

- Effaces (thins) & dilates (opens)
- After fetus descends into the vagina, the cervix is drawn upward and over the first portion

PELVIC FLOOR MUSCLES

- Helps the fetus rotate anteriorly

VAGINA

INTROITUS

- External opening of the vagina

LABOR & BIRTH PROCESSES

POSITION

POSITION OF THE MOTHER DURING BIRTH

UPRIGHT POSITION

Sitting on a birthing stool or cushion

LITHOTOMY POSITION

Supine position with buttocks on the table

MOST COMMON

"ALL FOURS" POSITION

On all fours: putting your weight on your hands & feet

LATERAL POSITION

Laying on a side

Frequent changes in position helps with:

- Relieving fatigue
- Increasing comfort
- Improving circulation

POWERS

CONTRACTIONS: PRIMARY & SECONDARY

PRIMARY POWERS

INVOLUNTARY uterine contractions
Signals the beginning of labor

DILATION

- Dilation of the cervix is the gradual enlargement or widening of the cervical opening & canal once labor has begun
- Pressure from amniotic fluid can also apply force to dilate

0 - 10 cm
closed full dilation

MEASURED IN CM

EFFACEMENT

- Shortening & thinning of the cervix during the first stage of labor
- Cervix normally:
 - 2 - 3 CM** long
 - 1 CM** thick
- The cervix is "pulled back / thinned out" by a shortening of the uterine muscles

Degree of **EFFACEMENT** is **EXPRESSED** in % (0-100%)

SECONDARY POWERS

VOLUNTARY bearing-down efforts by the women once the cervix has dilated

- Does not affect cervical dilation but helps with expulsion of infant once the cervix is fully dilated
- When the presenting part reaches the pelvic floor, the contractions change in character & become expulsive.
- Laboring women start to feel an involuntary urge to push & she uses secondary powers to aid in the expulsion of the fetus

FERGUSON REFLEX

- When the stretch receptors release oxytocin, it triggers the maternal urge to bear down

PSYCHOLOGY

EMOTIONAL RESPONSE

Anxiety can increase pain perception & the need for more medications (analgesia & anesthesia)

THINGS TO CONSIDER:

SOCIAL SUPPORT




PAST EXPERIENCE

KNOWLEDGE

NEWBORN ASSESSMENT

APGAR

7 - 10 supportive care
4 - 6 moderate depression
< 4 aggressive resuscitation

SCORE	0 POINTS	1 POINT	2 POINTS
A Activity (Muscle tone)	Absent	Flexed arms & legs	Active
P Pulse			
G Grimace (Reflex irritability)	Floppy	Minimal response to stimulation	Prompt response to stimulation
A Appearance (Skin color)	Blue / pale all over	Pink body, Blue extremities (acrocyanosis)	Pink all over
R Respiration (Effort)	No Breathing	Slow & irregular	Vigorous cry

VITAL SIGNS

RESPIRATORY RATE:

30 - 60 breaths/min

HEART RATE: 110 - 160 BPM

Can be 180 if crying
Can be 100 if sleeping
Take apical pulse for 1 full min

TEMPERATURE (AUXILIARY):

97.7° - 99.5° F
36.5° - 37.5° C

BLOOD PRESSURE:

Not done routinely
Systolic 60 - 80 mm Hg
Diastolic 40 - 50 mm Hg

MAP

Equal to the # of weeks gestation or higher

SIGNS OF RESPIRATORY DISTRESS
Retractions
Nasal flaring
Grunting

Breathing pattern is **IRREGULAR**.
Newborns are **ABDOMINAL** breathers.

To count breaths, place your hand on their abdomen  Count for a full minute!



GENERAL CHARACTERISTICS

Head & Chest Circumference

HEAD CIRCUMFERENCE

32 - 39 cm
14 - 15 inches
*measure above eyebrows

CHEST CIRCUMFERENCE

30 - 36 cm
12 - 14 inches
*measure above nipple line

Length & Weight

EXPECTED LENGTH

44 - 55 cm
17 - 22 in

EXPECTED WEIGHT

2,500 - 4,000 g
5 lb, 8 oz - 8 lb, 14 oz

INITIAL GOALS

1ST PRIORITY = AIRWAY

Suction with bulb syringe / deep suction
*Newborns are obligatory nose breathers

2ND PRIORITY = WARMTH

Dry with a blanket or place in warmer

CIRCULATORY SYSTEM

- Blood flow from umbilical vessels & placenta stop at birth
- Acrocyanosis:**
Blueness of hands & feet (normal during the first 24 hours of life)
- Closure of:
 - Ductus arteriosus
 - Foramen ovale
 - Ductus venosus
- Transient murmurs are normal

HEAD



CAPUT SUCCEDANEUM:

- Edema (collection of fluid)
- Crosses the suture lines




Like a baseball **CAP**



CEPHALOHEMATOMA:


- Birth trauma (collection of blood)
- Does not cross the suture lines



MOLDING:

abnormal head shape that results from pressure (normal)

Fontanelles may be bulging when the newborn cries, vomits, or is lying down. This is normal.



FONTANELLES:

Bulging = increase ICP or hydrocephalus
Sunken = dehydration

UMBILICAL CORD

Should have **2 ARTERIES** & **1 VEIN**



Should be dry, no odor, & no drainage



looks like a smiley face!


↓ TEMP → HEAT LOSS DUE TO:

EVAPORATION: Moisture from skin & lungs

CONVECTION: Body heat to cooler air

CONDUCTION: Body heat to a cooler surface in direct contact

RADIATION: Body heat to a cooler object nearby





POSTPARTUM ASSESSMENT

"BUBBLES"

B

BREASTS

- May be sore after breastfeeding
- Breastfeed every 2 - 3 hours (15 - 20 minutes each breast)
- Position newborn "tummy to mummy"
- Latch should be completely around the areola

MASTITIS

Infection & inflammation of breast tissue

- Continue breastfeeding
- Warm compress
- Hydration
- Rest
- Analgesics
- Wash hands!

U

UTERUS

UTERINE ATONY

RISK FACTORS

- Retained placenta
- Chorioamnionitis (infection)
- Uterine fatigue
- Full bladder

SYMPTOMS

- Enlarged
- Soft
- Boggy
- Not midline
- Poorly contracted uterus

INTERVENTIONS

- Fundal massage
- Assist to void or use in-and-out catheter

B

BOWELS

Constipation is common after birth. Increasing **FLUIDS & FIBER** may help!

HEMORRHOIDS

- May see blood in the stool
- Should begin to shrink following birth

INTERVENTIONS

- Tucks / witch hazel
- Ice pack
- Squeeze bottle
- Sitz Bath

B

BLADDER

- Postpartum urinary retention is common
 - In-and-out catheterization may be needed
 - Bladder distention can cause a displaced & boggy uterus!

SIGNS OF INFECTION



- Foul smelling or purulent lochia
- Fever (>100.4 F)
- Abdominal tenderness
- Tachycardia

L

LOCHIA

"Really Sore After"



RUBRA

bright red
1 - 3 days

SEROSA

pinkish/brown
4 - 10 days

ALBA

whitish-yellow
10 - 14 days

**Can last up to 6 weeks*

E

EMOTIONAL STATUS

- Postpartum depression (PPD) is common for women following childbirth
- As the nurse ask about feelings of...
 - depression • hopelessness • self-harm • harm to the newborn*
- Crying
- Irritable
- Sleep disturbances
- Anxiety
- Feelings of guilt

S

SECTION (c-section incisions) / Episiotomy

- Promote proper wound healing
- Report to the health care provider: *pain • inflammation • surrounding skin is warm to touch*

POSTPARTUM HEMORRHAGE

POSTPARTUM HEMORRHAGE is defined as:


VAGINAL BIRTH: loss of >500 ml of blood
CESAREAN BIRTH: loss of >1,000 ml of blood
A CHANGE IN HEMATOCRIT BY 10%

PATHOLOGY

The uterus is like a **BASKET WEAVE OF MUSCLE FIBERS** that crimps off vessels **protecting mom from hemorrhage.**

If the uterus is not doing this crimping off, it causes bleeding!

The uterus is often called the **LIVING LIGATURE**



SIGNS & SYMPTOMS

- ♥ Hypotonia of the uterus
- ♥ Atony / boggy uterus
- ♥ Deviated to the right
- ♥ Uncontrolled bleeding

#1 cause of uterine atony is **A FULL BLADDER**

RISK FACTORS

- ♥ Multiple gestations
- ♥ Polyhydramnios
- ♥ Macrosomic fetus (> 8 lbs)
- ♥ Multifetal gestation

overdistended uterus

DRUGS



MEMORY TRICK

"OH MY HEMORRHAGE"

This is a way to remember the order in which the drugs are used

#1

OXYTOCIN

"Pitocin"

ACTION

Stimulates contraction of the uterine smooth muscle

#2

METHERGINE

"Methylergonovine"

ACTION

Vasoconstriction

CONTRAINDICATIONS

Contraindicated in people with hypertension

*Remember vasoconstriction causes blood pressure to rise

#3

HEMABATE

ACTION

Hemabate is a prostaglandin! Hemabate helps control blood pressure and muscle contractions (uterine contractions).

CONTRAINDICATIONS

Contraindicated in people with asthma

Another medication that can be used:

MISOPROSTOL

given rectally

ACTION

Stimulates contraction of the uterine smooth muscle



PEDIATRICS

BROUGHT TO YOU BY



PEDIATRIC MILESTONES

INFANT BIRTH - 12 MONTHS

	GROSS MOTOR	FINE MOTOR	LANGUAGE
1 MONTH	<ul style="list-style-type: none"> • Head lag • Rounded back while sitting • Lifts and turns head to the side in prone position 	<ul style="list-style-type: none"> • Fists mostly clenched • Involuntary hand movements 	
2 MONTHS	<ul style="list-style-type: none"> • Raises head & chest • Head control improving 		<ul style="list-style-type: none"> • Makes verbal noise (coos)
3 MONTHS	<ul style="list-style-type: none"> • Raises head 45 degrees in prone • Tiny head lag in pull-to-sit 	<ul style="list-style-type: none"> • Holds hand in front of face with hands open 	
4 MONTHS	<ul style="list-style-type: none"> • Lifts head & looks around • Rolls from prone to supine • Head leads body when pulled to sit 	<ul style="list-style-type: none"> • Bats at objects 	<ul style="list-style-type: none"> • Babbling (copies noises)
5 MONTHS	<ul style="list-style-type: none"> • Rolls from supine to prone & back again • Sits with back upright when supported 	<ul style="list-style-type: none"> • Grasps rattle 	
6 MONTHS	<ul style="list-style-type: none"> • Tripod sit 	<ul style="list-style-type: none"> • Releases objects in hand to take another 	<ul style="list-style-type: none"> • Babbles (nonspecific)
7 MONTHS	<ul style="list-style-type: none"> • Sits alone with some use of hands for support 	<ul style="list-style-type: none"> • Transfers objects from one hand to the other 	
8 MONTHS	<ul style="list-style-type: none"> • Sits unsupported 	<ul style="list-style-type: none"> • Gross pincer grasp (rakes) 	
9 MONTHS	<ul style="list-style-type: none"> • Crawls with abdomen off the floor 	<ul style="list-style-type: none"> • Bangs objects together 	
10 MONTHS	<ul style="list-style-type: none"> • Pull to stand • Able to cruise on objects (furniture) 	<ul style="list-style-type: none"> • Fine pincer grasp • Puts objects into containers & takes them out 	
11 MONTHS		<ul style="list-style-type: none"> • Offers objects to others & releases them 	
12 MONTHS	<ul style="list-style-type: none"> • Walks independently • Sits down from standing position without assistance 	<ul style="list-style-type: none"> • Feeds self finger-foods • Draws simple marks on paper • Turns pages in a book 	<ul style="list-style-type: none"> • RECEPTIVE LANGUAGE • EXPRESSIVE LANGUAGE • SIGNS OF DELAY

MEMORY TRICK
Rolls on the floor rhymes with four!

MEMORY TRICK
You grasp something with 5 fingers



RECEPTIVE LANGUAGE

- Understands common words independent of context
- Follows a one-step gestured command

EXPRESSIVE LANGUAGE

- First word (example: "mama")
- Uses a finger to point to things
- Imitates: gestures & vocal



SIGNS OF DELAY



- After independent walking for several months
 - Persistent tiptoe walking
 - Failure to develop a mature walking pattern

PEDIATRIC MILESTONES

TODDLER 1-3 YEARS

15 MONTHS

18 MONTHS

24 MONTHS

30 MONTHS

GROSS MOTOR

- Walks independently



- Climbs stairs
- Pulls toys

- Kicks a ball
- Able to stand on tiptoes
- Climbs on & off furniture



Think Terrible Two's!

FINE MOTOR

- Feeds self finger foods
- Uses index finger to point
- Full pincer grasp developed

- Uses their hands a lot for: reaching, grabbing, releasing, stacking blocks
- Turns book pages
- Removes shoes and socks
- Stacks four cubes

- Builds tower of 6-7 cubes
- Right/left-handed
- Scribbles, paints, & imitates strokes
- Turns doorknobs
- Puts round pegs into holes

RECEPTIVE LANGUAGE

- Understands 100-150 words
- Follows commands without gestures
- Looks at adults when communicating

- Understands "no"
- Understands 200 words
- Says: "what's this?"

- Points to named body parts/pictures in books
- Listens to simple stories
- Says: "my" & "mine"

- Follows a series of 2 independent commands

EXPRESSIVE LANGUAGE

- Repeats words
- Babbles sentences

- Vocab: 15-20 words
- Uses names of familiar objects

- Vocab: 40-50 words
- Sentences of 2-3 words (ex. "want cookie")
- Use descriptive words: hungry, hot, cold

- Vocab: 150-300 words

SIGNS OF DELAY



- Persistent tiptoe walking
- Does not develop a mature walking pattern




- Not walking
- Not speaking 15 words
- Does not understand the function of common household items

- Does not: use two-word sentences, imitate actions, or follow basic instructions
- Cannot push a toy with wheels



PEDIATRIC MILESTONES

PRESCHOOL 3-6 YEARS

	3 YEARS	4 YEARS	5 YEARS
GROSS MOTOR	<ul style="list-style-type: none"> • Climbs well and runs easily • Pedals tricycle • Walks up & down stairs with alternating feet • Bends over without falling 	<ul style="list-style-type: none"> • Throws ball overhead • Kicks ball forward • Can bounce a ball back • Hops on one foot • Alternating feet going up & down steps 	<ul style="list-style-type: none"> • May be able to: <ul style="list-style-type: none"> • Skip • Swim • Skate • Climb • Swing
FINE MOTOR	<ul style="list-style-type: none"> • Undresses self • Copies circles • Tower of 9-10 • Holds a pencil • Screws and unscrews lids • Turns book pages one at a time 	<ul style="list-style-type: none"> • Uses scissors • Copies capital letter • Draws circles, squares, & traces a cross or diamond • Draws a person with 2-4 body parts • Laces shoes 	<ul style="list-style-type: none"> • Can draw a person and some letters • May dress/undress themselves • Can use a fork, spoon, & knife • Mostly cares for own toileting needs
COMMUNICATION	<ul style="list-style-type: none"> • Understands most sentences • Understands physical relation (in, on, under) • Follows a 3-part command • Half of the conversation understood by outside family • Says: "why?" • 3 or 4-word sentences • Talks about past • Vocab: 1,000 words • Says their name, age, & gender • Uses pronouns and plurals 	<ul style="list-style-type: none"> • Speaks in complete sentences • Tells a story • 75% of speech understood by outside observers • Stays on topic in conversation • Knows the name of familiar animals • Knows at least one color • Uses language to engage in make-believe • Can count a few numbers • Vocab: 1,500 words 	<ul style="list-style-type: none"> • Most of the child's speech can be understood • Explains how an item is used • Participates in long & detailed conversations • Talks about past, future, and imaginary events • Answers questions that use "why" and "when" • Can count to 10 • Says name & address • Recalls part of a story • Speech should be completely intelligible, even if the child has articulation difficulties • Speech is generally grammatical correct • Vocab: 2,000 words 
SIGNS OF DELAY	<ul style="list-style-type: none"> • Difficulty with stairs • Falls a lot while walking • Can't build a 4+ block tower • Extreme difficulty separating from parents • No make-believe play • Can't copy a circle • No short paragraphs • Doesn't understand simple instructions • Unclear speech & drooling • Little interest in other kids 	<ul style="list-style-type: none"> • Can't jump in place or ride a tricycle • Can't stack 4 blocks • Can't throw a ball overhead • Does not grasp crayon with thumb and fingers • Difficulty with scribbling • Can't copy a circle • Doesn't say 3+ word sentences • Can't use the words "me" & "you" • Ignores other children or doesn't show interest in interactive games • Still clings or cries if parents leave 	<ul style="list-style-type: none"> • Sad often • Little interest in playing with other kids • Unable to separate from their parents • Is extremely aggressive, fearful, passive, or timid. • Easy distracted (can't concentrate for 5 minutes) • Can not do ADL's by themselves (brush teeth, undress, wash & dry hands, etc) • Rarely engages in fantasy play

PEDIATRIC MILESTONES

PHYSIOLOGICAL CHANGES

10-13
YEARS

EARLY ADOLESCENCE

14-16
YEARS

MIDDLE ADOLESCENCE

17-20
YEARS

LATE ADOLESCENCE

MALE

- Pubic hair spread laterally, begins to curl, pigmentation increases
- Growth & enlargement of testes & lengthening of the penis
- Lengthy look due to extremities growing faster than the trunk

- Pubic hair becomes more coarse in texture & takes on adult distribution
- Testes, scrotum, & penis continue to grow
- The skin around the scrotum darkens
- Glands penis develops
- May experience breast enlargement
- Voice changes

- Mature pubic hair distribution & coarseness
- Breast enlargement disappears
- Adult size & shape of testes, scrotum, and penis
- Scrotum skin darkening

FEMALE

- First menstrual period (average age is 12 years)
- Breasts bud and areola continue to enlarge (no separation of the breasts)
- Pubic hair begins to curl & spread over the mons pubis

- Pubic hair becomes coarse in texture
- Amount of hair increases
- Areola & papilla separate from the contour of the breasts to form a secondary mound

- Mature pubic hair distribution and coarseness

PEDIATRIC CPR (<12 MONTHS)

Cardiac arrest in infants usually stems from **RESPIRATORY ETIOLOGY**

PEDIATRIC VITAL SIGNS

AGE	RESPIRATIONS	PULSE	SYSTOLIC BP
NEWBORN	30 - 50	120 - 160	60 - 80
6 MO - 1 YR	30 - 40	120 - 140	70 - 80
2 - 4 YR	20 - 30	100 - 110	80 - 95
5 - 8 YR	14 - 20	90 - 100	90 - 100
8 - 12 YR	12 - 20	80 - 100	100 - 110
> 12 YR	12 - 20	60 - 90	100 - 120

BREATHS/MIN

BEATS/MIN

ORDER OF EVENTS

1 PULSE

- * Check pulse no longer than 10 seconds

INFANT: Check **BRACHIAL** pulse

CHILD: Check **CAROTID** pulse

2 CALL FOR HELP

- * Active the emergency response system / shout for nearby **HELP**
- * Delegate someone else to call 911 / get the AED

3 CHEST COMPRESSIONS

- * 2 minutes of CPR before retrieving the AED
- * Rate of 100 - 120 compression/min
- * Using either 2 fingers or 2 thumbs on the sternum
- * Depth: **INFANT:** Equal to one-third of chest's anterior-posterior diameter
CHILD: 2 inches
- * Allow for recoil between compressions


SINGLE RESCUER

30:2 compression-to-breath ratio


TWO RESCUERS

15:2 compression-to-breath ratio

**2 - FINGER
COMPRESSION
TECHNIQUE**



**2 - THUMB
ENCIRCLING HAND
TECHNIQUE**



4 CONTINUE UNTIL SIGNS OF HELP ARRIVE OR AED BECOMES AVAILABLE

PIAGET'S STAGES OF COGNITIVE DEVELOPMENT



Saying Piaget's Cognitive Stages is Fun


SENSORIMOTOR STAGE

0 - 2 YEARS



- * Development through our 5 senses
- * Development through motor response
- * **OBJECT PERMANENCE** is developed
- * Egocentric

➔ Can only see the world from one's own point of view



Realizing that objects that are out of sight still exist

PREOPERATIONAL STAGE

2 - 7 YEARS



- * Symbolic thinking
- * Imagination
- * Abstract thinking is still difficult
- * Asks a lot of questions (intuition)


- Magical thinking
- **ANIMISM** - thinks objects are alive
- Plays pretend

CONCRETE OPERATIONAL STAGE

7 - 11 YEARS



- * Develop concrete cognitive operations
 - ➔ Sorting blocks in a certain order
- * **CONSERVATION** is developed
- * Conductive reasoning (Mathematical advancements)



CONSERVATION
Understanding that something stays the same in volume even though its shape changes.

FORMAL OPERATIONAL STAGE

> 11 YEARS




- * More rational, logical, organized, moral, and consistent thinking
- * **HYPOTHETICAL THINKING** - Can think outside the present
- * Abstract concepts
 - ➔ Love, hate, failures, successes
- * Deductive reasoning

VARIATIONS IN PEDIATRIC ANATOMY & PHYSIOLOGY

RESPIRATORY

- Narrow airways
- Newborns have ↓ alveoli than an adult
 - ➔ Thousands of alveoli grow each day for the first few months of life!
- Floppy airways from less cartilage
- Obligatory nose breathers
- ↑ metabolic rate
- ↑ O₂ requirements



HEAD SIZE

- Head is the fastest growing part of an infant (large in proportion to the body!)
- Head & neck muscles are not well developed

BRAIN & SPINAL CORD

- Cranial bones not completely fused
- The brain is highly vascular = ↑ risk for hemorrhage
- Sutures & fontanelles makes the skull flexible and allows for growth of the brain
- The spine is very mobile = ↑ risk for cervical spine injury

EARS

- ↑ RISK FOR EAR INFECTION
- Eustachian tubes are short, wide, & flat = making drainage difficult = harbors microorganisms

CARDIOVASCULAR


- The transition from fetal circulation → normal circulation at birth
- Infants hearts are thinner and less compliant

SKIN

- Epidermis is thinner
- Blood vessels are closer to the surface - loses heat very easily!

KIDNEYS

- Kidneys are larger in relation to abdomen = less protection
- GFR is slower
- ↓ ability to concentrate urine & reabsorb = ↑ risk for dehydration



IMMUNE SYSTEM

- ↑ RISK FOR INFECTION
- Immature immune systems
- ↓ inflammatory response
- Limited exposure to disease (losing immunity from maternal antibodies)

NERVOUS SYSTEM


- Myelination is incomplete at birth
- Myelination happens in **CEPHALOCAUDAL DIRECTION** (head to tail)

CEPHALOCAUDAL DIRECTION (HEAD TO TAIL)

HEAD CONTROL BEFORE WALKING!



PROXIMODISTAL (INWARD OUTWARD)



SUDDEN INFANT DEATH SYNDROME (SIDS)

Sudden death of a previously healthy infant younger than 1 year of age



RISK FACTORS


- AGE: 1 - 6 months (↑ risk)
- Preterm
- Sleep position
- Sibling death
- Nicotine exposure
- Socioeconomic status
- Lack of prenatal care
- Genetic
- Bedding (can be smothered)
- Room temp (cooler is better)

THERE ARE NO SIGNS OR SYMPTOMS!

Sudden death
Leading cause of death in infants

EDUCATION / PREVENTION


- Sleep in **SUPINE POSITION**
- Bedding
 - **Firm** mattress
 - No toys, blankets, pillows, or stuffed animals
- Avoid over bundling or overdressing the infant
- Avoid smoking
- **No co-bedding**
(Infant should sleep separate from the parents)
- Normal room temp
- Encourage pacifier use



ABCs OF SAFE SLEEPING

- A** Alone
- B** On their **B**ack
- C** In a **C**rib

NEURAL TUBE DEFECTS



NORMAL SPINE

The neural tube closes:
3rd - 4th week of gestation

SPINA BIFIDA

is a general term for a birth defect typically diagnosed during pregnancy where the spinal column fails to close.

Spina bifida means **"SPLIT SPINE"**

CAUSES

NOT KNOWN...

BUT MANY FACTORS HINDER NORMAL CNS DEVELOPMENT

- Drugs
- Chemicals
- Folic acid deficiency (Vitamin B9)
- Malnutrition
- Genetics
- Diabetes
- Obesity



SPINA BIFIDA OCCULTA

MILDEST FORM

Defect of the vertebral body **WITHOUT** protrusion of the spinal cord or meninges.

Typically asymptomatic

May have dimpling, abnormal patches of hair, or discoloration near the spine.

Does not need immediate medical care if asymptomatic.

If symptoms are present, the client may get an MRI.




MENINGOCELE

Sac protruding from the spinal area.
Most are covered with skin.

Meninges herniate through a defect in the vertebrae.

Usually minor or no neurological deficits.

Surgical correction of the lesion



MYELOMENINGOCELE

MOST SEVERE FORM

Protrusion of the meninges, cerebrospinal fluid, and spine.
Skin may be exposed as well.


The spinal cord often ends at the point of the defect.

=

Absent motor & sensory function beyond that point.

- Multiple surgical procedures
- Paralysis
- Bladder / bowel incontinence
- Neurogenic bladder
- Meningitis (infection)
- Hypoxia
- Hemorrhage
- Freq. catheterization causes...
 - ➔ Latex allergy
 - ➔ UTIs / pyelonephritis
 - ➔ Renal damage

BRONCHIOLITIS (RSV)



PATHO

small airways in the lungs

BRONCHIOLITIS
inflammation

- * Viral illness usually caused by **Respiratory syncytial virus (RSV)**
- * Very contagious
- * Starts as an upper respiratory infection & moves into the chest

SIGNS & SYMPTOMS

INITIAL

- * Upper respiratory symptoms
 - ➔ Nasal congestion
 - ➔ Runny nose
 - ➔ Cough
 - ➔ Sneezing
- * Fever

CONTINUED

- * Lower respiratory tract symptoms
 - ➔ Tachypnea
 - ➔ Cough
 - ➔ Wheezing

EMERGENT

- * Grunting
- * Nasal flaring
- * Cyanosis
- * Hypoxia
- * Respiratory failure
- * Apneic episodes

TREATMENT

- * Self-limited illness & supportive care
- * Airway maintenance
 - ➔ Oxygen
 - ➔ Suctioning
Saline nose drops & then suction the nares with a bulb syringe to remove the secretions before feeding or at bedtime
 - ➔ Position the child at a 30 - 40 degree angle
- * Hydration
Increase fluid intake (oral or IV) (risk for dehydration)
- * Hospitalization
Only necessary if the child has severe symptoms
- * Use contact & standard precautions during care

**MOST
CHILDREN
CAN BE
MANAGED
AT HOME**

REYES SYNDROME

RARE DISEASE AFFECTING YOUNG CHILDREN
RECOVERING FROM A VIRAL ILLNESS
(FLU OR CHICKEN POX)




CAUSE

EXACT CAUSE UNKNOWN

Triggered due to the intake of salicylates or salicylate-containing products such as **aspirin** to treat a viral illness (Flu / Chickenpox)



SIGNS & SYMPTOMS



ENCEPHALOPATHY / CEREBRAL EDEMA

ACUTE FATTY LIVER FAILURE

LABS

- ↑ LIVER ENZYMES
- ↑ AST
- ↑ ALT



“CHILD(S)”

- C Confusion (changes in mental status)
- H Hyperreflexia
- I Irritability
- L Lethargy
- D Diarrhea & vomiting
- S Seizures

TREATMENT


- * Early recognition & treatment
- * Education on prevention!
- * Monitor fluid status
- * Swelling of the brain occurs
 - ➔ Maintaining cerebral perfusion
 - ➔ Managing & preventing increased ICP
 - ➔ Seizure precautions

Educate on products that contain **SALICYLATES:**

ASPIRIN
ALKA-SELTZER
PEPTO-BISMOL
KAOPECTATE

INTUSSUSCEPTION

PATHO



ILEUM TELESOPES INTO THE CECUM

↓
OBSTRUCTION = PAIN

↓
COMPRESSION OF BLOOD VESSELS

↓
BLOOD FLOW DECREASES

↓
BOWEL ISCHEMIA

↓
RECTAL BLEEDING (**CURRENT JELLY STOOLS!**)

SIGNS & SYMPTOMS

- * Intermittent pain / cramping
- * Child draws up their legs toward the abdomen in severe pain while crying
- * Vomiting & diarrhea
- * **Currant-jelly stools (bloody)**
- * Lethargy
- * Sausage-shaped mass in the upper mid-abdomen

THIS IS BECAUSE
TELESCOPING IS
INTERMITTENT

CAUSES

- * **NOT COMPLETELY KNOWN**
- * May be due to a virus that causes swelling
- * Condition child is born with
 - ↳ Diverticulum
 - ↳ Polyps

TREATMENT

- * May spontaneously be reduced (Passage of normal, brown stools)
- * IV fluids
- * Antibiotics
- * Decompression via NG tube
- * Provide comfort & emotional support to the parents
- * Monitor for signs of perforation & shock
- * May need air or barium enema
 - ↳ Provide education to child & family about pre-op & post-op

DIAGNOSTIC / TREATMENT

AIR or BARIUM ENEMA works to diagnose & also helps reduce the intussusception


HYPERTROPHIC PYLORIC STENOSIS

PATHO

A HYPERTROPHIED PYLORIC MUSCLE CAUSES NARROWING OF THE PYLORIC CANAL



THICKNESS CREATES A NARROW STOMACH OUTLET



NORMAL

HYPERTROPHIED PYLORUS MUSCLE

HYPERTROPHIC



INCREASE IN SIZE

PYLORIC



PYLORUS

STENOSIS



NARROWING

Opening from the stomach into the small intestines

SIGNS & SYMPTOMS

- * Projectile vomiting
- * Non-bilious emesis
- * Olive-shape mass palpable in the right upper quadrant
- * Infants will be hungry constantly despite regular feedings
- * Weight loss
- * **DEHYDRATION!**

↑ Hematocrit from hemoconcentration
↑ BUN

STOMACH CONTAINS ACID WHICH BECOMES DEPLETED WHEN VOMITING WHICH LEADS TO **METABOLIC ALKALOSIS**
↑PH & ↑HCO₃

TREATMENT

- * Monitor ...
 - ➔ I&O's
 - ➔ Vomiting episodes & stools
 - ➔ Signs of dehydration & electrolyte imbalances
- * Obtain daily weights
- * Provide comfort & emotional support to the parents
- * Educate about surgery

PYLOROMYOTOMY

Cut the muscle of the pylorus
↓
Relieving the gastric outlet obstruction

EPIGLOTTITIS

PATHO


Inflammation of the **EPIGLOTTIS** leading to an **UPPER AIRWAY OBSTRUCTION**

WHAT IS THE EPIGLOTTIS?

Piece of cartilage at the back of the tongue

FUNCTION:

Closes the entry to the trachea during swallowing....
AKA prevents aspiration




CAUSES

- ✦ Most common cause: **HAEMOPHILUS INFLUENZA TYPE B**
- ✦ Streptococcus pneumonia

PEDS incident falling due to Hib vaccination

SIGNS & SYMPTOMS

- ✦ Tachycardia
- ✦ Sore throat
- ✦ High fever
- ✦ Anxious / apprehensive / agitation
- ✦ Difficulty speaking
- ✦ Nasal flaring
- ✦ Stridor (Frog-like croak on inspiration)
- ✦ Drooling / dysphagia
- ✦ **Tripod position**
- ✦ Sitting forward with the neck extended to breath - mouth open
- ✦ Retractions (chest)
- ✦ Nasal flaring
- ✦ Absent cough!



NURSING MANAGEMENT

- ✦ Never leave the client
- ✦ Asses oxygen status
- ✦ IV access
- ✦ May need emergency intubation
- ✦ Calm environment
 - Stay with parents
 - Don't restrain the child
 - Help to avoid crying
 - Most comfortable position (usually tripod position)
- ✦ Do not place them in supine position. It becomes harder to breathe.

Do not visualize the throat with a tongue blade.
Take oral temperature or take throat culture...

WHY? It can cause **REFLEX LARYNGOSPASMS** (cutting off the airway)

- ✦ NPO
- ✦ Medications
 - Antibiotics
 - Antipyretics
 - Corticosteroids (decrease inflammation)
 - IV Fluids

LARYNGOTRACHEOBRONCHITIS “CROUP”

PATHO


Inflammation of the **LARYNX, TRACHEA, & BRONCHI** occur as a result of viral infection


Most commonly caused by the **PARAINFLUENZA VIRUS**



SIGNS & SYMPTOMS

* Inflammation & edema obstructs the airway

*  Symptoms occur at night

- Stridor
- Subglottic swelling (causes hoarseness in the voice)
- Seal-bark cough 

CROUP VS. EPIGLOTTITIS

ONSET	Sudden (at night)	Rapid (within hours)
FEVER	Fluctuating	High
COUGH	Yes	No
DYSPHAGIA	No	Yes
CAUSE	Viral	Bacterial
EMERGENCY	Not typically	Yes



HOME CARE

*Self-limiting
(Usually resolves on its own)*

- * Corticosteroids (↓ inflammation)
- * Racemic epinephrine
- * Humidified air (steamy bathroom or mist humidifier)
- * Encourage rest & fluid intake
- * Calm environment for the child



SEEK HELP

When the child is indicating respiratory distress


- * Child is confused/restless
- * Blue lips/nails
- * ↑ respiration rate (breathing faster, but less air is going in)
- * Retractions
- * Nasal flaring
- * Drooling/can't swallow

TREATMENT

FEVER MANAGEMENT

NORMAL TEMP

97.5°F to 98.6°F
36.4°C to 37.0°C




FEVER

> 100.4°F (38.0°C)

SIGNS & SYMPTOMS

- * Flushed skin
- * Diaphoresis (sweating)
- * Chills
- * Restlessness
- * Lethargy



TREATMENT

- * Administer antipyretics (ibuprofen) → Do not administer aspirin (risk for Reye's Syndrome)
- * Monitor for S&S of dehydration & electrolyte imbalances → Provide adequate fluids!
- * Sponge bath → Tepid water for 20-30 min. Squeeze over back & body
- * Remove excess clothing & coverings to ↓ the temp
- * Cool compress on the forehead

FEBRILE SEIZURE

WHAT IS IT?

Seizures associated with a **FEVER**

Not related to:

- ➔ intracranial infection
- ➔ metabolic imbalance
- ➔ viral illness

Usually **DOES NOT** have long term complications such as epilepsy or intellectual disability

SIGNS & SYMPTOMS

- * Rapid ↑ in core temperature
- * Child may be drowsy during postictal period

RISK FACTORS

- * 6 months - 5 years
- * Rapidly developed fever
- * HIGH fever
- * Family history of febrile seizures
- * Certain vaccines
 - ➔ DTP & MMR

TREATMENT


- * NOT anticonvulsants therapy
- * Rectal Diazepam
- * Educate the parents to seek help if...
 - ➔ Last > 5 min
 - ➔ Repeated seizures

CYSTIC FIBROSIS (CF)

PATHO

- * Multisystem disorder of the **EXOCRINE GLANDS** with increased production of thick mucus
- * Gene mutation (CFTR): prevents exocrine glands from properly functioning
- * **EXOCRINE GLANDS:** Produce & transfer secretions (mucus, tears, sweat, & enzymes) via ducts
- * ↑ viscosity of mucous = ↑ resistance to ciliary action = slowing the flow rate of mucous, leading to **mucous plugging**

CF IS AN AUTOSOMAL RECESSIVE GENETIC DISORDER



DIAGNOSIS

- * Ambyr test
- * Positive sweat sodium chloride test
- * Genetic screen

TREATMENT

- * **Treatment of the mucous**
 - ➔ Chest physiotherapy (PT)
 - ➔ Postural drainage
 - ➔ Huff coughing
 - ➔ Nebulizers
 - Bronchodilators, mucolytics, anti-inflammatories
- * **Treat & prevent infection**
 - ➔ Wear a mask, hand washing, up-to-date on vaccines, avoid those who are sick.
- * **Nutrition**
- * **Prevent GI blockage**
 - ➔ Fluids & stool softeners



CHEST PT

- * Drains airways of thick mucous to be coughed up
 - ➔ Stimulates cough
 - ➔ Helps loosen mucous
 - ➔ Results in deep breathing
 - ➔ Builds up strength and endurance of respiratory muscles
 - ➔ Improves cardiovascular fitness
- * Done multiple times a day between 1-2 hour increments
 - ➔ NOT done right before or after meals!
- * Causes vibrations & percussions to break apart the mucus (vests, manual vibration)



- * ↑ protein, ↑ fat, ↑ calorie
 - ➔ Fat soluble vitamin supplementation A, K, E, D
- * Possible supplemental oral feeding or enteral feeding
- * **Pancreatic enzymes:**
 - ➔ Pancrelipase or Pancreatin
 - ➔ Can swallow a capsules or sprinkle enzymes on foods that are acidic such as apple sauce!

All Kids Eat Donuts



MANIFESTATIONS OF CF

RESPIRATORY

- **INFECTION:** Thick mucous creates a great environment for bacterial growth
 - ➔ Pseudomonas
 - ➔ Staph. aureus
- Pneumonia
- Bronchitis
- **Thick mucus = blocked airways**
 - ➔ Obstructive pulmonary disease (Emphysema)
 - Clubbing
 - Barrel-shape chest
- Pneumothorax
- Strain on lungs = pulmonary hypertension

CARDIOVASCULAR

- Pulmonary hypertension puts strain on the heart
 - ➔ **Right-sided heart failure**

INTEGUMENTARY

- Sweat glands produce ↑ chloride = salty skin
- Salty sweat & salty tears which leads to
 - ➔ Dehydration
 - ➔ Electrolyte imbalance

REPRODUCTIVE

BOYS

- Thick mucus blocks the vas deferens = Infertility

GIRLS

- Thick cervical mucus blocks sperm from penetrating = Infertility

**BOTH
HAVE
DELAYED
PUBERTY**

NOSE & SINUSES

- Sinusitis
- Nasal polyps (snoring, stuffiness)

PANCREAS

PANCREAS SECRETES THICK MUCUS


- Deficient in pancreatic enzymes: (Protease, Amylase, Lipase)
 - ➔ Weight loss
 - ➔ Inadequate protein absorption
 - ➔ Deficiency of protein
 - ➔ Failure to thrive
- Insulin deficiency
 - ➔ Hyperglycemia
 - ➔ CF-related diabetes

LIVER

- Bile duct blocked from **THICK mucus**
 - ➔ Gallstones
 - ➔ Biliary cirrhosis

STOMACH & INTESTINES

- Fecal impaction
- Rectal prolapse
- Bowel obstruction
- Intussusception
- Back up of stool in intestine
 - ➔ Constipation
 - ➔ Vomiting
 - ➔ Abdominal distention
 - ➔ Cramping
 - ➔ Anorexia
 - ➔ RLQ pain
- Meconium ileus in infants
- Steatorrhea
 - ➔ Frothy (bulky), fatty, foul-smelling stools



FETAL CIRCULATION IN UTERO

FORAMEN OVALE

Blood is **SHUNTED** from the right atrium to the left atrium by the **FORAMEN OVALE**

Blood bypasses the lungs...why?

It's already oxygenated blood from the placenta (mom)

How can blood be shunted from the right atrium to the left atrium?

PRESSURE DIFFERENCE!


Blood flows from **high resistance** to **low resistance**.

Lungs: High resistance from all the fluid. So the blood does not want to go in the lungs!

RIGHT ATRIUM

Blood goes from the interior vena cava to the right atrium as well as some **deoxygenated blood** coming from the **SUPERIOR VENA CAVA**.

So the blood is now **MIXED** (oxygen-rich & oxygen-poor blood)



DUCTUS VENOSUS

Umbilical vein is carrying **oxygenated** blood from the placenta. It passes the **LIVER** (Some blood will go to the liver) but most will be **SHUNTED** to the **INFERIOR VENA CAVA** by the **DUCTUS VENOSUS**

Liver not fully functioning yet

DUCTUS ARTERIOSUS

Blood is **SHUNTED** from the pulmonary artery into the aorta by the **ductus arteriosus**

AORTA

Mixed blood is now in the aorta and being pushed out to oxygenate the fetus

BLOOD GOES BACK TO THE PLACENTA TO GET OXYGENATED AGAIN!

THE PLACENTA IS THE "LIFELINE" BETWEEN MOTHER & BABY



The Placenta is like **"TEMPORARY LUNGS"** for the fetus while in utero

2 UMBILICAL ARTERIES
1 UMBILICAL VEIN



A think AWAY

Takes deoxygenated blood + waste **AWAY** from the baby back to the placenta

Gives oxygen rich blood **TO** the baby

SHUNTS TO KNOW


- * **DUCTUS VENOSUS**
- * **FORAMEN OVALE**
- * **DUCTUS ARTERIOSUS**

START

DEVELOPMENT DYSPLASIA OF THE HIPS (DDH)

PATHO

- * Abnormal development of the hip joint
- * A baby's bones are not ossified yet so they have the ability to dislocate & relocate easily




- DISLOCATION** No contact between femoral head & acetabulum
- SUBLUXATION** Partial dislocation (acetabulum is not completely in contact with the hip joint)
- DYSPLASIA** Hip joint doesn't have the proper shape to fit together correctly

DIAGNOSIS

- * Ultrasound for in utero
- * X-ray for those older than 6 months
- * Barlow test & Ortolani

Listen for any noises during the exam. There should be no "clunks" heard or felt. If "clunks" are felt or heard = a positive sign for DDH



COMPLICATIONS

- * Avascular necrosis of the femoral head
- * ↓ ROM
- * Leg-length discrepancy
- * Early osteoarthritis
- * Femoral nerve palsy

RISK FACTORS

- * FEMALE → more lax ligaments from maternal hormones
- * Breech positioning
- * Oligohydramnios

TREATMENT

Early detection & treatment are crucial. The bones are not ossified in early infancy, so you want to manipulate them to grow properly. If DDH is not treated early the bones will ossify and develop incorrectly.

> 6 MONTHS

- * **Pavlik harness:** Stabilizes the hip by preventing hip extension

4 MONTHS - 2 YEARS

- * **Closed reduction:**
 - Requires general anesthesia where the hips will be placed back into the acetabulum by the surgeon
 - Spica cast is worn after surgery to maintain reduction
 - After spica cast the child will wear a brace until acetabulum is fully normal

> 2 YEARS OR NO IMPROVEMENTS WITH SURGERY OR HARNESS

- * Open surgical reduction followed by casting



INSTRUCTIONS FOR PAVLIK HARNESS

- * Must wear the harness at all times!
- * Do not adjust the straps or remove harness until instructed by the HCP
- * Change the diaper while the baby is in the harness
- * Check for redness, irritation or breakdown 2-3 times per day
- * Place baby on their back to sleep
- * Place long knee socks and undershirt to prevent rubbing of the harness

SCARLET FEVER

SCARLET-FEVER
THINK STREP!

PATHO

- * Complication of **group A streptococcal infection** AKA **Strep throat**
- * Not all children who have strep will develop scarlet fever
- * **TRANSMISSION:** Droplets & respiratory tract secretions.
Transmission happens in close contact such as schools & daycares.

SIGNS & SYMPTOMS

- * Onset: **ABRUPT!**
- * **RED RASH!**
 ➔ Sandpaper-like rash
- * Pharyngitis
- * Fever, body aches, chills
- * Strawberry tongue
- * Tender cervical nodes
- * Tonsils are red
- * Exudate may be present

Begins on the **NECK & CHEST** and spreads outwards to **THE EXTREMITIES!**
Rash is usually not seen on the palms & soles of the feet



S's of SCARLET FEVER

- * **S**trawberry tongue
- * **S**andpaper rash



COMPLICATIONS

- * Rheumatic fever
- * Glomerulonephritis
- * Abscesses of the throat
- * Pneumonia

Early diagnosis & treatment are very important to prevent complications!

TREATMENT

Most children can be cared for at home

- * Antibiotics (Penicillin V)
 ➔ Erythromycin for those allergic to Penicillin
- * Fluids & soft foods
- * Provide comfort
- * Cool mist humidifier

Take antibiotics as directed....
Finish the medication even if the child appears to be better!



SOUPS, TEAS, POPSICLES, SLUSHIES





MED-SURG

BROUGHT TO YOU BY




KIDNEY OVERVIEW

FUNCTIONS

- A** CID-BASE BALANCE
- W** ATER BALANCE
- E** LECTROLYTE BALANCE
- T** OXIN REMOVAL
- B** LOOD PRESSURE CONTROL
- E** RYTHROPOIETIN
- VITAMIN **D** METABOLISM

ANATOMY OF THE KIDNEY




TERMS TO KNOW

- DYSURIA**Pain while urinating
- NOCTURIA**Excessive urination at night
- HEMATURIA**Bloody urine
- FREQUENCY**Voiding more than every 3 hours
- URGENCY**Strong desire to void

- INCONTINENCE**Involuntary voiding
- ENURESIS**Involuntary voiding during sleep
- PROTEINURIA**Abnormal amounts of protein in the urine
- OLIGURIA**Urine output: <400 mL/day
- ANURIA**Urine output: <50 mL/day
- MICTURITION**Voiding

URINE FORMATION



LAB VALUES RELATED TO THE KIDNEYS


GFR	Glomerular Filtration Rate: rate of blood flow through the kidneys.	90 - 120 ML/MIN
BUN	Blood Urea Nitrogen: Normal waste product resulting from the breakdown of proteins. ↑ Levels can indicate a kidney problem & be toxic in the body.	7 - 20 MG/DL
CREATININE	End product of muscle metabolism solely filtered from the blood via glomerulus	0.6 - 1.2 MG/DL
URINE SPECIFIC GRAVITY	Measures the kidney's ability to excrete or conserve water	1.010 - 1.030
CREATININE CLEARANCE	The amount of blood the kidneys makes per minute that is FREE of creatinine	FEMALES: 85 - 125 ML/MIN MALES: 95 - 140 ML/MIN

You will see **INCREASED BUN & Creatinine** levels during kidney injury/failure

ACUTE GLOMERULONEPHRITIS (POSTSTREPTOCOCCAL)

PATHOLOGY


- 1 Untreated strep
- 2 Immune system response by creating **antigen-antibody complexes** (14 days after infection)
- 3 These antibodies get "lodged" in the glomeruli
- 4 Inflammation & scarring
- 5 ↓ GFR



*It's not the strep that causes the inflammation of the kidneys. It's the **antigen-antibody complexes** that form due to the strep that causes the inflammation & damage to the glomeruli*

SIGNS & SYMPTOMS

- Hematuria → Blood in the urine
- Azotemia → Excessive nitrogenous waste in the blood
Tea colored urine (cola color)
- Malaise
- Headache
- Proteinuria (mild)
- Hypoalbuminemia
- ↓ GFR = Oliguria
- Edema
 - Swelling in the face/eyes
- ↑ Blood pressure
- Retaining sodium
- ↑ Urine specific gravity
- ↑ BUN & creatinine
- (+) ASO (Antistreptolysin) Titer



MAIN CAUSE:
RECENT GROUP A BETA-HEMOLYTIC
STREPTOCOCCAL INFECTION

INTERVENTIONS

- Fix the cause! (strep)
- Diet modifications
 - Fluid restriction
 - Sodium restriction
 - ↓ Protein
 - Provide a lot of carbohydrates
- Monitor
 - Daily intake & output
 - Daily weight
- Bed rest
- Monitor blood pressure
 - Antihypertensives
 - Diuretics


Carbohydrates provide energy & stop the breakdown of protein

A weight gain of 1 kg is equal to 1,000 mL of retained fluid

ACUTE KIDNEY INJURY (AKI)

WHAT IS IT?

Sudden renal damage! Causes a build-up of waste, fluid, and electrolyte imbalance. It can be reversible. Formerly called *Acute Renal Failure*.



PHASES "OH OH DARN RENAL"

INTERVENTIONS

OH ONSET / INITIATION	➔ Triggering event (Prerenal, Intrarenal or Postrenal Failure)	➤ Correct & identify the underlying cause to prevent long term damage to nephrons!
OH OLIGURIC	➔ ↓ Urine output < 400 mL/24 hrs Glomerulus decreases the ability to filter blood (↓ GFR)	<ul style="list-style-type: none"> ➤ Low protein diet ➤ Limit fluid intake ➤ Strict I&O + daily weights ➤ Monitor EKG & labs <ul style="list-style-type: none"> • Watch for HYPERkalemia > 5.0 • ↑ BUN & Creatinine ➤ Dialysis may be needed until kidney function returns
DARN DIURETIC	➔ Cause of AKI is corrected Gradual ↑ in urinary output	<ul style="list-style-type: none"> ➤ Large amount of dilute urine with electrolytes ➤ Monitor the patient for dehydration & hypokalemia
RENAL RECOVERY	➔ ↑ in kidney function May take up to 6 - 12 months	➤ Some patients may never recover and may develop chronic kidney disease (CKD)

NEPHROTIC SYNDROME

PATHOLOGY

INFLAMMATORY RESPONSE IN THE GLOMERULUS

DAMAGE TO MEMBRANE

LOSS OF PROTEIN (ALBUMIN)

Albumin regulates oncotic pressure

HYPOALBUMINEMIA

LOW ALBUMIN LEVELS

CAUSES
SYNTHESIS OF
CHOLESTEROL
& TRIGLYCERIDES

HYPERLIPIDEMIA

FLUID
SHIFT

GENERALIZED EDEMA

ALBUMIN IS A
PROTEIN WHICH
PREVENTS CLOT
FORMATION

POSSIBLE BLOOD CLOTS
(THROMBOSIS)

CAN LOSE
PROTEIN THAT
HELPS FIGHT
INFECTIONS
(Immunoglobulins)

RISK FOR INFECTION

CAUSES


- Bacteria or viral infection
- Cancer
- Genetic predispositions
- Systemic disease (lupus or diabetes)
- NSAIDs

SIGNS & SYMPTOMS

- Hypoalbumenia
 - Edema
 - Fatigue & loss of appetite
 - Hyperlipidemia
- Proteinuria (> 3 g/day)
 - Large amounts of protein in the urine

INTERVENTIONS

- Monitor fluid status
 - Daily weights & I&O's
 - Swelling & abdominal girth
- Diet modifications
 - ↓ Cholesterol & saturated fats
 - ↓ Na⁺ intake
 - Moderate protein intake
- Medications
 - Diuretics
 - Statins (lipid-lowering drugs)
 - Prednisone to ↓ inflammation
 - Antineoplastic agent
 - Immunosuppressant
- Monitor signs of...
 - Infection
 - Blood clots



CHRONIC KIDNEY DISEASE (CKD)

PATHO

- Progressive & irreversible loss of kidney function.
- Occurs over a long period of time.

CAUSES

- Untreated acute kidney injury (AKI)
- Diabetes mellitus
- Hypertension
- Family history
- Recurrent infections
- Autoimmune disorders

STAGES

Stages are based on the GFR rate
AS CKD WORSENS... GFR DECREASES ↓

STAGE 1 GFR > 90

STAGE 2 60 - 89

STAGE 3 A: 45 - 59
B: 30 - 44

STAGE 4 15 - 29

STAGE 5 < 15
(END STAGE RENAL DISEASE) ⚠

TREATMENT

- Dialysis
- Kidney transplant

SIGNS & SYMPTOMS

In the end stages of CKD,
ALMOST EVERY BODY SYSTEM is negatively affected




- ↓ Urinary output (UOP)
 - Oliguria = <400 mL/day
 - Anuria = <100 mL/day
- Proteinuria & hematuria



- Lethargy
- Altered LOC/confusion
- Seizures



- Hypertension
- Fluid volume excess (Hypervolemia)
- Heart failure



- Anorexia
- Nausea/vomiting
- Uremic fetor (ammonia breath)
- Metallic taste




- Impaired immune & inflammatory response



- Anemia (↓ erythropoietin [EPO])
- ↑ Risk for bleeding
- Prolonged bleeding time



- Amenorrhea
- Erectile dysfunction
- ↓ Libido



- Uremic frost
- Pruritus




- LABS**
- ↑ BUN
 - ↑ Creatinine
 - ↑ K+
 - ↑ Magnesium
 - ↓ Calcium
 - ↑ Phosphate

TYPES OF DIALYSIS

HEMODIALYSIS

MOST COMMON METHOD

3X a week (3 - 5 hours per treatment)



THE DIALYZER (Artificial kidney)

Brings blood to the dialyzer

Filters out toxins/waste products

Brings clean blood back to the body

VASCULAR ACCESS

FISTULA

Joining an artery to a vein

GRAFT

Inserting synthetic graft material between an artery and vein

Needs time to heal and mature

BOTH REQUIRE SURGERY

EVALUATION OF PATENCY

- ✓ Feel the thrill...
- ✓ Hear the bruit...

COMPLICATIONS


- Hypotension
- Disequilibrium syndrome
- Hemorrhage
- Air embolus
- Electrolyte imbalances

AVOID...

- ✗ Compression
- ✗ Blood draws
- ✗ Blood pressure readings
- ✗ Tight clothing
- ✗ Carrying bags
- ✗ Sleeping on that arm

PERITONEAL DIALYSIS

INSIDE THE BODY



Warm the solution!

Dialysate is infused into the peritoneal cavity by gravity

Close the clamp on the infusion line

Dialysate dwells for a set amount of time (dwell time)

The drainage tube is unclamped

Fluid drains from the peritoneal cavity by gravity

A new container of dialysate is infused as soon as drainage is complete

REPEAT!

PERITONEAL CATHETER

Performed at the bedside or in the operating room

COMPLICATIONS

- Peritonitis (infection)
 - Cloudy or bloody drainage
 - Fever
 - Abdominal pain
 - Malaise

URINARY TRACT INFECTION

PATHO

Infection within the urinary system caused by either a **BACTERIA**, **VIRAL**, or **FUNGUS**.

BACTERIA
IS MOST COMMON
SPECIFICALLY
E. COLI

CAUSES


- Most common in women (shorter urethra & urethra is close to the rectum)
- Overuse of antibiotics
- Indwelling catheters
- Hormone changes (pregnancy changes)
- Diabetes
- Lifestyle
 - Baths, scented tampons, perfumes etc.

EDUCATION

- Take entire antibiotics course
- Wipe from front to back
- Void after intercourse
- Avoid caffeine & ETOH
- Void frequently
- Avoid bubble baths, perfumes, or sprays!
- Wear non-tight cotton underwear


NURSING CONSIDERATIONS

- Maintain fluid status
 - 2 - 3 L per day → "flushing" out the urinary tract
 - Remove the catheter ASAP (per HCP order)
- Medications
 - Antibiotics → Take urine culture BEFORE giving first dose of antibiotics
 - Analgesia (control pain)
 - Phenazopyridine (Pyridium) → Analgesic to ↓ pain
May turn urine orange



SIGNS & SYMPTOMS

- Smelly urine
- Chills & fever
- Costovertebral angle (CVA) tenderness**
- Nausea & vomiting
- Headache/malaise
- Painful urination (dysuria)
- Burning on urination
- Frequency & urgency
- Nocturia
- Incontinence
- Hematuria
- Fever
- WBC's in the urine



**ELDERLY CLIENTS
MAY SHOW
DIFFERENT SYMPTOMS**

- Confusion
- Lethargy
- New incontinence

RENAL CALCULI


PATHO

Stones (calculi) found in the urinary tract & kidney!

NEPHROLITHIASIS:
stones in the **kidneys**

URETEROLITHIASIS:
stones in the **ureter**

- Stones can be very large or very small
- They can be found inside the **KIDNEYS, URETERS,** or the **BLADDER**



TREATMENT

MOST COMMONLY, THE STONE WILL PASS ON IT'S OWN!

- MEDICATIONS** to control the ***PAIN***
 - NSAIDs
 - Opioids analgesics

↓ **Pain & inflammation**
(makes the stone easier to pass)
- STRAIN THE URINE**
 - keep any stones & send them to the lab to evaluate the type of stone
- GET THEM MOVING OR FREQUENTLY TURNING THEM!**
- ↑ FLUIDS!**

Push stone forward & out!
Decreases risk of infection
- DIET**
 - Limit protein, NA+ foods, & calcium
- PROCEDURES:**

NON INVASIVE Extracorporeal Shock Wave Lithotripsy (ESWL)
Sends shock waves to break up the stone!

INVASIVE! Percutaneous Nephrolithotomy
Stone removed by an incision made on the back where the kidneys are located.

SIGNS & SYMPTOMS

- PAIN!**
- Discomfort
- Hematuria → (RBCs)
- Pyuria → (WBCs)
- Nausea & vomiting

DIAGNOSIS

- KUB: X-ray of kidneys, ureters, bladder
- IVP: intravenous pyelogram
- Ultrasound or CT scan
- Urine test

What is URIC ACID?

Uric acid is a waste products of the breakdown of **purines**

	MOST COMMON! CALCIUM	URIC ACID	STRUVITE	RARE! CYSTINE
STONE TYPE	Forms due to ↑ amounts of calcium & oxalate in the urine	Too much uric acid in the urine (acidic urine)	Persistent alkaline environment that is ammonia-rich urine Due to a bacteria	Rare, genetic, inherited disorder that affects renal absorption of cystine
CAUSES	Hypercalcemia Hypercalciuria Hyperparathyroidism ↑ Intake of Na+ Dehydration GI disorders ↑ Intake of calcium supplements with vit D	Gout Foods high in purine or animal proteins Dehydration Metabolic issues (Diabetes)	Chronic urinary tract infections (UTI's) Foreign bodies Neurogenic bladder	

CARDIAC TERMS

CARDIAC OUTPUT (CO)

Total volume pumped per minute

Normal 4 - 8 L/min

Less volume = ↓ CO

More volume = ↑ CO

CO = HR x Stroke Volume

Cardiac Output = Heart Rate



↓ CO = ↓ perfusion to the body

- ↓ LOC
- Lungs sound wet due to back flow
- Shortness of breath
- Skin will be cold & clammy
- ↓ UOP
- Weak peripheral pulses

STROKE VOLUME

Amount of blood pumped out of the ventricle with each beat or contraction

CONTRACTILITY

Force / strength of contraction of the heart muscle

EJECTION FRACTION (EF)

% of blood expelled from the left ventricle with every contraction


Normal EF: 50 - 70%

EXAMPLE:

If the EF is 55%, the heart is pumping out 55% of what's inside of the left ventricle

PRELOAD

Amount of blood returned to the right side of the heart at the end of diastole



AFTERLOAD


Pressure that the left ventricle has to pump against (the resistance it must overcome to circulate blood)

Clinically measured by systolic blood pressure!

HEMODYNAMIC PARAMETERS

Cardiac output (CO)	Total volume pumped per minute	Normal 4 - 8 L/min
Cardiac Index (CI)	Cardiac output per body surface area $CI = \frac{CO}{\text{surface area}}$	2.5 - 4.0 L/min/m²
Central Venous Pressure (CVP)	Pressure in the superior vena cava. Shows how much pressure from the blood is returned to the right atrium from the superior vena cava.	2 - 8 mmHg
Mean Arterial Pressure (MAP)	Average pressure in the systemic circulation (your body) through the cardiac cycle	70 - 100 mmHg <i>At least 60 mm Hg is required to adequately perfuse the vital organs</i>
Systemic Vascular resistance (SVR)	The resistance it takes to push blood through the circulatory system to create blood flow	800 - 1200 dynes/sec/cm

FLOW OF BLOOD THROUGH THE HEART



RIGHT

Deoxygenated Blood

- 1 Superior Vena Cava / Inferior Vena Cava
- 2 Right Atrium (RA)
- 3 Tricuspid Valve (TV)
- 4 Right Ventricle (RV)
- 5 Pulmonary Valve (PV)
- 6 Pulmonary Artery*

carries
DEOXYGENATED
blood to the **LUNGS**

LEFT

Oxygenated Blood

- 7 Pulmonary Vein*
- 8 Left Atrium
- 9 Bicuspid/Mitral Valve
- 10 Left Ventricle
- 11 Aortic Valve
- 12 Aorta

carries
OXYGENATED
blood to the
TISSUES/BODY

OVERVIEW OF BLOOD VESSELS

ARTERIES

Carry oxygenated blood to tissues

VEINS

Carry deoxygenated blood back to the heart



MEMORY TRICK


Arteries think **A**way from the heart

* EXCEPTIONS

The only exception to this is the **PULMONARY ARTERY** and **PULMONARY VEIN**

brings deoxygenated blood from the heart to the lungs

carries oxygenated blood from the lungs to the heart



ELECTRICAL CONDUCTION OF THE HEART

CARDIAC CONDUCTION SYSTEM:

Generates & transmits **ELECTRICAL IMPULSES** which stimulates contractions of the atria and then the ventricles.



MNEMONIC

STEPS IN THE HEART'S CONDUCTION SYSTEM

SEND
A
BIG
BOUNDING
PULSE

SA node (SinoAtrial node)
AV node (AtrioVentricular)
Bundle of His
Bundle branches (right & left)
Purkinje fibers

Primary pacemaker of the heart. Creates electrical impulses of **60 - 100 bpm**

Secondary pacemaker of the heart "backup pacemaker." If the SA node malfunctions, the AV node takes over at a rate of **40 - 60 bpm**

If the SA & the AV nodes fail, the Purkinje fibers can fire at a rate of **30 - 40 bpm**


NOTE:
This is a normal heart rate

AUSCULTATING HEART SOUNDS

5 AREAS FOR LISTENING TO THE HEART



All People Enjoy Time Magazine



Aortic Right 2nd intercostal space

Pulmonic Left 2nd intercostal Space

ERB's Point (S1, S2) Left 3rd intercostal space

Tricuspid Lower left sternal border 4th intercostal

Mitral Left 5th intercostal, medial to midclavicular line



Think **M** for **M**idclavicular & **MITRAL** has **5** letters for "**5**th intercostal space"

NORMAL

S1

LUB

Tricuspid & mitral valve closure



S2

DUB

Aortic & pulmonic valve closure

CLOSING OF THE VALVES

Valve opening does not normally produce a sound

ABNORMAL

S3

EARLY DIASTOLE in rapid ventricle filling



S4

LATE DIASTOLE & high atrial pressure (forcing blood into a stiff ventricle)

ABNORMAL VENTRICULAR FILLING

Extra ♥ sounds

SYSTOLE:

Ventricle
pump / ejection
=
LUB (S1)

DIASTOLE:

Ventricle
relax / filling
=
DUB (S2)




"COZY RED"

CO (contract) **ZY** (systole)
RE (relax) **D** (diastole)



EKG WAVEFORMS



- P WAVE** Atrial contraction (**DE**polarization)
- PR SEGMENT** Movement of electrical activity from atria to ventricles
- QRS COMPLEX**... Ventricle contraction (**DE**polarization)
- ST SEGMENT** Time between ventricular depolarization & repolarization
- T WAVE** Ventricle relaxing (**RE**polarization)
- TP INTERVAL** Ventricle are relaxing & filling

HEART RHYTHM MEASUREMENTS

- PR INTERVAL**
0.12 - 0.2
- QRS COMPLEX**
0.06 - 0.12
- QT INTERVAL**
< 0.40 seconds

BASIC RHYTHMS

- NORMAL SINUS** 60 - 100 bpm
- SINUS TACHYCARDIA** > 100 bpm
- SINUS BRADYCARDIA** < 60 bpm



DEpolarization think...
DEcompressing



REpolarization think...
RElaxing
REpolarizing
REfilling with blood

PR INTERVAL

Movement of electrical activity from atria to ventricles


ST SEGMENT

Time between ventricular depolarization and repolarization (ventricular contraction)

QT INTERVAL

Time take from ventricles to depolarize, contract, and repolarize

5-LEAD PLACEMENT



6 STEPS TO INTERPRETING EKGs

#1 P-WAVE

Identify & examine the P-waves

- Should be present & upright
- Comes before QRS complex
- One P-wave for every QRS complex

#2 PR INTERVAL

Measure PR interval

Normal PR interval:
0.12 - 0.2 seconds

#3 QRS COMPLEX

Is every P-wave followed by a QRS complex?

- Should not be widened or shortened – this may indicate problems!

Normal QRS complex:
0.06 - 0.12

Widen is often seen in PVCs, Electrolyte imbalances & drug toxicity!

#4 R-R

Are the R-R intervals consistent?


- Regular or irregular?

BASIC RHYTHMS

NORMAL SINUS 60 - 100 bpm

SINUS TACHYCARDIA > 100 bpm

SINUS BRADYCARDIA < 60 bpm



1 large box = 0.20 seconds

5 large boxes = 1 second

1 small box = 0.04 seconds

#5 DETERMINE THE HEART RATE

6 SECOND METHOD

Count the number of R's in between the 6 second strips & multiply by 10



6 R's X 10 = 60 beats per minutes

Be sure and check that the strip is 6 seconds!
Count the boxes

BIG BOX METHOD

300 divided by the number of big boxes between 2 R's



300 / 5 = 60 BPM

#6 IDENTIFY THE EKG FINDING!

EKGs

NORMAL SINUS RHYTHM



RATE	60 - 100 bpm
RHYTHM	Regular
P-WAVE	Upright & uniform before each QRS
PR INTERVAL	Normal
QRS COMPLEX	Normal



SINUS BRADY



KEY The sinus node creates an impulse at a **slower**-than-normal rate

RATE	< 60 bpm
RHYTHM	Regular
P-WAVE	Upright & uniform before each QRS
PR INTERVAL	Normal
QRS COMPLEX	Normal

CAUSES

- ♥ Lower metabolic needs
 - Sleep, athletic training, hypothyroidism
- ♥ Vagal stimulation
- ♥ Medications
 - Calcium channel blockers, beta blockers, Amiodarone

TREATMENT

- ♥ Correct the underlying cause!
- ♥ ↑ the heart rate to normal

SINUS TACHY



KEY The sinus node creates an impulse at a **faster**-than-normal rate

RATE	> 100 bpm
RHYTHM	Regular
P WAVE	Upright & uniform before each QRS
PR INTERVAL	Normal
QRS COMPLEX	Normal


CAUSES

- ♥ Physiologic or psychological stress
 - Blood loss, fever, exercise, dehydration
- ♥ Certain medications
 - Stimulants - caffeine, nicotine
 - Illicit drugs - cocaine, amphetamines
 - Stimulate sympathetic response - epinephrine
- ♥ Heart failure
- ♥ Cardiac tamponade
- ♥ Hyperthyroidism

TREATMENT

- ♥ Identify the underlying cause!
- ♥ ↓ the heart rate to normal

VENTRICULAR TACHYCARDIA (VT)



looks like tombstones

Irregular, coarse waveforms of different shapes. The ventricles are quivering and there is **no contractions or cardiac output** which may be **fatal!**

RATE	100 - 250 bpm
RHYTHM	Regular
P-WAVE	Not visible
PR INTERVAL	None
QRS COMPLEX	Wide (like tombstones) > 0.12 seconds

CAUSES

- ♥ Myocardial ischemia / infarction
- ♥ Electrolyte imbalances
- ♥ Digoxin toxicity
- ♥ Stimulants: caffeine & methamphetamine

MANIFESTATIONS

- ♥ Patient is usually awake (unlike V-fib)
- ♥ Chest pain
- ♥ Lethargy
- ♥ Anxiety
- ♥ Syncope
- ♥ Palpitations



TREATMENT

STABLE CLIENT WITH A PULSE

- ♥ Oxygen
- ♥ Antidysrhythmics (ex. Amiodarone...stabilizes the rhythm)
- ♥ Synchronized Cardioversion

- Synchronized administration of shock (delivery in sync with the QRS wave).
- Cardioversion is NOT defibrillation! (defibrillation is only given with deadly rhythms!)

UNSTABLE CLIENTS WITHOUT A PULSE

Also called **PULSELESS V-TACH**


- ♥ CPR
- ♥ Follow ACLS protocol for defibrillation
- ♥ Possible intubation
- ♥ Drug therapy
 - Epinephrine, vasopressin, amiodarone

SHOCK!

UNTREATED VT can lead to → **VENTRICULAR FIBRILLATION** → **DEATH** ☠

EKGs

VENTRICULAR FIBRILLATION (V-FIB)



Rapid, disorganized pattern of electrical activity in the ventricle in which electrical impulses arise from many different foci!

RATE	Unknown
RHYTHM	Chaotic & irregular
P-WAVE	Not visible
PR INTERVAL	Not visible
QRS COMPLEX	Not visible

CAUSES

- ♥ Cardiac injury
- ♥ Medication toxicity
- ♥ Electrolyte imbalances
- ♥ Untreated ventricular tachycardia

MANIFESTATIONS

- ♥ Loss of consciousness
- ♥ May not have a pulse or blood pressure
- ♥ Respirations have stopped
- ♥ **Cardiac arrest & death!**

NO CARDIAC OUTPUT
=
NO OXYGEN TO THE BODY

TREATMENT

- ♥ CPR
- ♥ Oxygen
- ♥ Defib (follow ACLS protocol for defibrillation) ⚡
- ♥ Possible intubation

- ♥ Drug Therapy
 - Vasoconstriction: Epinephrine
 - Antiarrhythmic: Amiodarone, lidocaine
 - Possibly magnesium



"Defib the Vfib"

CARDIOVERSION VS. DEFIBRILLATION



CARDIOVERSION


- ♥ Synchronized shock
- ♥ Lower amount of energy
- ♥ Not done with CPR
- ♥ Stable clients
 - ♥ Ex. A-fib

DEFIBRILLATION

- ♥ Asynchronous
- ♥ Higher amount of energy
- ♥ Resume CPR after shock
- ♥ Unstable clients
 - ♥ Example: pulseless VT or VF

ATRIAL FIBRILLATION (A-FIB)

IRREGULAR R-R INTERVALS



Uncoordinated electrical activity in the atria that causes rapid & disorganized "fibbing" of the muscles in the atrium.

RATE	Usually over 100 BPM
RHYTHM	Irregular
P-WAVE	None. They are irregular (fibrillary waves)
PR INTERVAL	Visible
QRS COMPLEX	Normal

**THE ATRIA IS
QUIVERING!**

CAUSES

- ♥ Open heart surgery
- ♥ Heart failure
- ♥ COPD
- ♥ Hypertension
- ♥ Ischemic heart disease

MANIFESTATIONS

- ♥ Most commonly asymptomatic
- ♥ Fatigue
- ♥ Malaise
- ♥ Dizziness
- ♥ Shortness of breath
- ♥ Tachycardia
- ♥ Anxiety
- ♥ Palpitations

**ALL DUE
TO
LOW O₂**

TREATMENT

STABLE PT.

- ♥ Oxygen
- ♥ Drug therapy!
 - Beta blockers
 - Calcium channel blockers
 - Digoxin
 - Amiodarone
 - Anticoagulant therapy to prevent clots

RISK FOR CLOTS!

The atria quiver causes pooling of blood in the heart which increases the risk for clots = increased risk for MI, PE, CVAs, & DVTs!

UNSTABLE PT.

- ♥ Oxygen
- ♥ Cardioversion
 - Synchronized administration of shock (delivery in sync with the QRS wave).
 - Cardioversion is NOT defibrillation!

DEFIBRILLATION

Defibrillation is only given with deadly rhythms!

EKGs

PREMATURE VENTRICULAR CONTRACTIONS (PVCs)



Early "premature" conduction of a QRS complex

RATE	Depends on the underlying rhythm
RHYTHM	Regular but interrupted due to early P-waves
P-WAVE	Visible but depends on timing of PVC (may be hidden)
PR INTERVAL	Slower than normal but still 0.12 - 0.20 seconds
QRS COMPLEX	Sharp, bizarre, and abnormal during the PVC

CAUSES

- ♥ Heart failure
- ♥ Myocardial ischemia / infarction
- ♥ Drug toxicity
- ♥ Caffeine, tobacco, alcohol
- ♥ Stress or Pain
- ♥ Increased workload on the heart

EXERCISE
FEVER
HYPERVOLEMIA
HEART FAILURE
TACHYCARDIA

BIGEMINY: every **other** beat

TRIGEMINY: every **3rd** beat

QUADRAGEMINY: every **4th** beat

R-ON-T PHENOMENON: PVC arises spontaneously from the repolarization gradient (T-wave) may precipitate V-fib

TREATMENT

TX based on underlying cause

- ♥ May not be harmful if the client has a healthy heart
- ♥ Oxygen
- ♥ Decrease caffeine intake
- ♥ Correct the electrolyte imbalances
- ♥ D/C or adjust the drug causing toxicity
- ♥ Decrease stress or pain



MANIFESTATIONS

- ♥ May be asymptomatic
- ♥ Feels like your heart...
 - "Skipped a beat"
 - "Heart is pounding"
- ♥ Chest pain

CHEST PAIN

Notify the healthcare provider if the client complains of chest pain, if the PVCs increase in frequency or if the PVCs occur on the T-wave (R-on-T phenomenon).

ASYSTOLE



RATE
RHYTHM
P-WAVE
PR INTERVAL
QRS COMPLEX

FLATLINE

CAUSES

- ♥ Myocardial ischemia/infarction
- ♥ Heart failure
- ♥ Electrolyte imbalances (common: hypo/hyperkalemia)
- ♥ Severe acidosis
- ♥ Cardiac tamponade
- ♥ Cocaine overdose

TREATMENT

- ♥ **High quality CPR**
 - Heel of hand on the center of the chest
 - Arms straight
 - Shoulders aligned over hands
 - Compress at 2 - 2.4 inches at a rate of 100 - 120 min
 - 30 compressions to 2 rescue breaths
 - Minimal interruptions

ATRIAL FLUTTER

SAWTOOTH



Similar to A-fib, but the heart's electrical signals spread through the atria. The heart's upper chambers (atria) beat too quickly but at a regular rhythm.

RATE	75-150 BPM
RHYTHM	Usually regular
P-WAVE	"Sawtooth" P-wave configuration shaped flutter waves
PR INTERVAL	Unable to measure
QRS COMPLEX	Usually normal & upright

CAUSES

- ♥ Coronary artery disease (CAD)
- ♥ Hypertension
- ♥ Heart failure
- ♥ Valvular disease
- ♥ Hyperthyroidism
- ♥ Chronic lung disease
- ♥ Pulmonary embolism
- ♥ Cardiomyopathy

MANIFESTATIONS

- ♥ May be asymptomatic
- ♥ Fatigue / syncope
- ♥ Chest pain
- ♥ Shortness of breath
- ♥ Low blood pressure

TREATMENT

STABLE PT.

- ♥ Drug therapy!
 - Calcium channel blockers
 - Antiarrhythmics
 - Anticoagulants

RISK FOR CLOTS!

Atrial flutter causes pooling of blood in the atria = risk for clots

UNSTABLE PT.

- ♥ Cardioversion
 - Synchronized administration of shock (delivery in sync with the QRS wave).
 - Cardioversion is NOT defibrillation!


DEFIBRILLATION

Defibrillation is only given with deadly rhythms!

HEART FAILURE

SIGNS & SYMPTOMS

LEFT SIDED HF



MEMORY TRICK Left Side think **L**ungs
Fluid is backing up into the **LUNGS** = pulmonary symptoms

- D**yspnea
- R**ales (crackles)
- O**orthopnea
- W**eakness / fatigue
- N**octurnal paroxysmal dyspnea
- I**ncreased HR
- N**agging cough (*frothy, blood tinged sputum*)
- G**aining weight (2 -3 lbs a day)

OTHER S&S
↑ UOP
Hypotension
S3 Gallop

RIGHT SIDED HF

Fluid is backing up into the **VENOUS SYSTEM**

- S**welling of the legs & hands
- W**eight gain
- E**dema (pitting)
- L**arge neck veins (JVD)
- L**ethargy / fatigue
- I**rrregular heart rate
- N**octuria
- G**irth (Ascites)

OTHER S&S
Hepatomegaly
Splenomegaly
Anorexia

SYSTOLIC HF VS. DIASTOLIC HF

SYSTOLIC HF

Weakened heart muscle

The ventricle does not **EJECT** properly

DIASTOLIC HF

Stiff & non-compliant heart muscle

The ventricle does not **FILL** properly

EJECTION FRACTION (EF)

Amount of blood **PUMPED OUT**

Amount of blood **IN THE CHAMBER**

% EF

EF REDUCED



NORMAL EF



**NORMAL EJECTION FRACTION
50% - 70%**

HEART FAILURE: DIAGNOSIS & INTERVENTIONS

DIAGNOSIS

BNP

B-TYPE NATRIURETIC PEPTIDE


Secreted when there is
↑ pressure in the ventricle

BNP 100 - 300 pg/mL.....HF is suspected

BNP > 300 pg/mL.....Mild HF

BNP > 600 pg/mL.....Moderate HF

BNP > 900 pg/mL.....Severe HF



CHEST X-RAY

Enlarged heart &
pulmonary infiltrates

ECHOCARDIOGRAM

LOOKS AT:
ejection fraction,
back flow,
& valve problems

Ejection Fraction (EF)

EF is ↓ in
most types of HF

NORMAL RANGE:
55% - 75%

INTERVENTIONS

MONITOR:

- Strict I&Os
- Daily weights
- Edema




Report
weight gain
(2-3 lbs)

REPORT S&S OF FLUID RETENTION

- Edema
- Weight gain

DIET MODIFICATIONS:

- Fluid restrictions
- ↓ Sodium
- ↓ Fat
- ↓ Cholesterol



ELEVATE HOB

(Semi-Fowler's position)

BALANCE PERIODS OF ACTIVITY & REST

CORONARY ARTERY DISORDERS (CAD)


RISK FACTORS

NON-MODIFIABLE

Age
Gender
Race
Family history

MODIFIABLE

Diabetes Obesity
Hypertension Physical inactivity
Smoking High cholesterol
Metabolic Syndrome



PATHO

Fatty plaques develop



Called **ATHEROSCLEROSIS**



Restriction of blood flow to the heart

SIGNS & SYMPTOMS

ISCHEMIA

Inadequate blood supply to the heart = ↓ O₂ to the heart.



ANGINA PECTORIS

Chest pain that is caused by **myocardial ischemia**

ISCHEMIA: ↓ O₂
INFARCTION: Death

- Chest pain w/ activity
- Shortness of breath
- Fatigued

PREVENTION

- Management of hypertension
- Management of diabetes
- Smoking cessation
- Diet
- Exercise

DIAGNOSIS

BLOOD TEST → Lipoprotein profile

- LDL
- HDL
- Total cholesterol
- Triglycerides

ECG

- Assess for changes in ST segments or T-waves!

TREATMENT

- Lipid-lowering medications "Statins"
- Heart-healthy diet
- Physical activity
- Smoking cessation
- Stress management
- Hypertension management
- Diabetes management
- Coronary stent / angioplasty
- Coronary Artery Bypass Graft (CABG)

WEEKLY EXERCISE GOALS
Moderate: 75 min
Vigorous: 150 min

CHOLESTEROL

LDL
Low Density Lipoprotein



Want **LOW** Levels (<100 mg/dL)
BAD Cholesterol

HDL
High Density Lipoprotein




Want **HIGH** Levels (>60 mg/dL)
HAPPY Cholesterol

PERIPHERAL VASCULAR DISEASE

is an umbrella term for...

PERIPHERAL VENOUS DISEASE (PVD)

Deoxygenated blood can't get back to the heart.
Pooling of oxygenated blood in the extremities.



- PAIN ?** ✓ Dull, constant, achy pain!
- PULSE ?** ✓ May not be palpable due to edema
- EDEMA ?** ✓ Blood is POOLING in the leg
- TEMP ?** Warm legs (Blood is warm)
- COLOR ?** Stasis dermatitis (Brown/yellow)
- WOUNDS ?** Venous STASIS ulcers, Irregular shaped wounds, shallow
- GANGRENE ?** ✗ We have too much blood! Gangrene is caused by insufficient amounts of blood.
- POSITIONING ?** Elevate Veins Positions that make it worse: dangling, sitting/standing for long periods of time

PERIPHERAL ARTERIAL DISEASE (PAD)

Think "BAD"

Narrow artery (atherosclerosis) where oxygenated blood can't get to the distal extremities (hands & feet).

Ischemia & necrosis of the extremities

- PAIN ?** ✓ Sharp pain: Gets worse at night "rest pain"
Intermittent claudication
- PULSE ?** ✓ Very poor or even absent
- EDEMA ?** ✗ No blood in the extremities
- TEMP ?** Cool No blood = cool leg (blood is warm)
- COLOR ?** Pale, hairless, dry, scaly, thin skin due to lack of nutrients (↓ O₂)
- WOUNDS ?** Regular in shape, red sores round appearance "punched out"
- GANGRENE ?** ✓ Tissue death caused by a lack of blood supply
- POSITIONING ?** Dangle arteries

CAUSES OF BOTH

Smoking • Diabetes • High cholesterol • Hypertension

DX: Doppler Ultrasound or Ankle Brachial Index (ABI)

TREATMENT — KEEP VEIN OPEN!

- Elevate Veins
- Medications
 - Aspirin or Clopidogrel
 - Cholesterol lowering drugs "statin"
- Surgery
 - Angioplasty
 - Bypass (CABG)
 - Endarterectomy

TREATMENT — GET BLOOD MOVING!

- **D**Angle Arteries (Dependent position)
- Perform daily skin care with moisturizer
- Stop smoking
- Avoid tight clothing (vasoconstriction)
- No heating pads!
- Medications
 - Vasodilators
 - Antiplatelets

PERIPHERAL VASCULAR DISEASE WORKSHEET


TEST YOUR KNOWLEDGE!
NO CHEATING! :)

is an umbrella term for...

PERIPHERAL VENOUS DISEASE (PVD)

Deoxygenated blood can't get back to the heart.

Pooling of oxygenated blood in the extremities.



PAIN ?

PULSE ?

EDEMA ?

TEMP ?

COLOR ?

WOUNDS ?

GANGRENE ?

POSITIONING ?

PERIPHERAL ARTERIAL DISEASE (PAD)

Think "BAD"

Narrow artery (atherosclerosis) where oxygenated blood can't get to the distal extremities (hands & feet).

Ischemia & necrosis of the extremities

PAIN ?

PULSE ?

EDEMA ?

TEMP ?

COLOR ?

WOUNDS ?

GANGRENE ?

POSITIONING ?

CAUSES OF BOTH

DX: _____

TREATMENT

• Position

• Medications

• Surgery

TREATMENT

• Position

• Perform _____

• Stop _____

• Avoid _____


• No _____

• Medications

WANT MORE WORKSHEETS?
Check out The Complete Laminated Study Templates!



ANGINA PECTORIS



Angina is chest pain associated with ischemia. It's due to narrowing of at least one major coronary artery.

TYPES OF ANGINA

STABLE

“Predictable”

Occurs with **EXERTION**

EXAMPLE Exercise or strenuous activity

UNSTABLE

“Preinfarction”

Occurs at **REST & MORE FREQUENTLY**

PRINZMETAL'S / VARIANT

“Coronary artery vasospasm”

Pain at **REST** with reversible **ST-ELEVATION**

SIGNS & SYMPTOMS

- Chest pain (heavy sensation) may radiate to neck, jaw, or shoulders
- Unusual fatigue
- Weakness
- Shortness of breath
- Pallor
- Diaphoresis



INTERVENTIONS

GOAL: ↓ oxygen demand

REPERFUSION PROCEDURES

PCI

Percutaneous
Coronary Interventions

CABG

Coronary Artery
Bypass Graft

DRUG THERAPY

NITRATES

Vasodilators
↓ ischemia = ↓ pain
Usually administered sublingual

CALCIUM CHANNEL BLOCKERS

Relaxes blood vessels
↑ oxygen supply to the heart
↓ workload of heart

BETA BLOCKERS

↓ myocardial oxygen consumption

ANTIPLATELET / ANTICOAGULANT

Prevents platelet aggregation & thrombosis

PATIENT TEACHING

SUBLINGUAL NTG OR SPRAY

- 1 tab/spray sublingual every 5 minutes, up to 3 doses.
- If angina is not relieved or is worse 5 min after the first dose, call 911!


Keep in original container (dark, glass bottle) in a dry, cool place. Do not swallow or chew these tablets



MYOCARDIAL INFARCTION (MI)


PATHO

Complete blockage in one or more arteries of the heart




ATHEROSCLEROSIS

Coronary arteries become narrow due to plaque build-up




ANGINA

Due to ischemia (low O₂)



MYOCARDIAL INFARCTION (MI)

Plaque rupture become a blood clot that blocks arteries of the heart



SIGNS & SYMPTOMS

Sudden, crushing, radiating chest pain that continues despite rest & medications

- Shortness of breath
- Nausea & vomiting
- Sweating
- Pale & dusty skin

WOMEN PRESENT WITH DIFFERENT SYMPTOMS

- Fatigue
- Shoulder blade discomfort
- Shortness of breath

PAIN FELT IN THE... Left arm • Mid back/shoulder • Heartburn

DIAGNOSIS

* ECG

- ST-Elevation (no O₂)
- ST-Depression (low O₂)
- T-wave inversion

* TROPONIN

* STRESS TESTS

- Chemical & exercise

TREATMENT

IMMEDIATE

M MORPHINE

↓ workload of the heart & ↓ pain

O OXYGEN

↑ O₂ to the heart

N NITROGLYCERIN

opens up the vessels

A ASPIRIN

Prevents platelets from sticking together

CATH LAB OR CLOT BUSTER

MEDICATIONS

- Thrombolytics (clot busters)
- Example: Streptokinase

SURGERY

- PCI "Percutaneous Coronary Intervention"
- CABG
- Endarterectomy - Cut out the blockage

Suffixes:
-teplase
-ase

PREVENTION & REST

PREVENT / STABILIZE CLOT

- Heparin IV

REST THE HEART WITH...

- Nitro
- Beta-Blockers
- Calcium channel blockers

Any time you give a thrombolytic, watch for signs of bleeding!



CARDIAC BIOMARKERS

TROPONIN

Protein released in the blood stream when the heart muscle is damaged.

There are 3 isomers of troponin

- ➔ **Troponin C:** binds calcium to activate muscle contraction
- ➔ **Troponin I & T:** specific for cardiac muscle

BEST indicator of an acute MI

TROPONIN I
RANGE

0 - 0.03 ng/mL

PEAK

CAN REMAIN ELEVATED FOR AS LONG AS 2 WEEKS

CK-MB

CREATINE KINASE - MB

An enzyme released in the bloodstream when the heart, muscles or brains are damaged!

Cardiac-specific isoenzyme
BUT less reliable than Troponin

RANGE

0 - 5 ng/mL

PEAK

24 HOURS

MYOGLOBIN

Myoglobin is found in cardiac & skeletal muscle

NOT a specific indicator of an acute MI, but a (-) sign is good for ruling out an acute MI



Myoglobin think Muscle

RANGE

5 - 70 ng/mL

PEAK

12 HOURS

BNP

BRAIN NATRIURETIC PEPTIDE

A peptide released when the ventricle is filled with too much fluid and **STRETCHES!**

Indicates heart failure (HF)

RANGE

Normal: **<100 pg/mL**

Mild HF: **100 - 300**

Moderate HF: **300 - 700**

Severe HF: **>700**

HYPERTENSION (HTN)

HYPERtension = **HIGH** BP

MOST ACCURATE DIAGNOSIS FOR HTN

CATEGORIES	SYSTOLIC (SQUEEZE)	DIASTOLIC (DECOMPRESS)
NORMAL	< 120	< 80
PRE-HTN	120 - 139	80 - 89
STAGE 1 HTN	140 - 159	90 - 99
STAGE 2 HTN	> 160	> 100
HTN CRISIS	> 180	> 120

AFFECTED ORGANS




CONGESTIVE HEART FAILURE (CHF)

Overworking of the heart muscle (ventricle enlarges)



STROKE

Weak & narrow vessels could lead to rupture of vessels



RENAL FAILURE

Too much blood flowing to the kidneys at a fast rate & high pressure



VISUAL CHANGES

Damages blood vessels in the retina (blurred vision, can't focus on objects)

RISK FACTORS

MOST COMMON

PRIMARY HTN

Also called **ESSENTIAL** or **IDIOPATHIC HTN**

- Cause is unknown
- Not curable, only controllable

- F** Family HX
- A** Advanced age
- C** ↑ Cholesterol
- R** Race (African Americans)
- T** Too much caffeine
- I** Intake of Na/ETOH
- O** Obesity
- S** Smoking
- R** Restricted activity
- K** Low K+ & vitamin D levels
- S** Sleep apnea

SECONDARY HTN

- Has a direct cause / preexisting condition
 - Chronic kidney disease
 - Diabetes
 - Hypo/Hyperthyroidism
 - Cushing syndrome
 - Pregnancy
 - Certain drugs (oral contraceptives)

SIGNS & SYMPTOMS

Usually asymptomatic!

Commonly called the **"SILENT KILLER"**

- Symptoms: (if seen)
- Blurred vision
 - Headache
 - Chest pain
 - Nose bleeds

EDUCATION

- Limit sodium intake
- Limit alcohol intake
- Smoking cessation
- Teach how to measure BP & keep a record
- Exercise programs for weight loss if needed

CHECKING BLOOD PRESSURE



- Place stethoscope over brachial artery
- Patients should not smoke, exercise, etc. within 30 minutes of having their BP checked (could lead to inflated BP)
- Instruct the client to:
 - Sit in a chair with legs uncrossed
 - Arm at ♥ level
 - Correct size cuff
- No BPs should be auscultated in arms with:
 - Mastectomy
 - HX of AV shunt
 - Blood clots
 - PICC lines/central lines

Too small = false high BP


Too large = false low BP

ANTIHYPERTENSIVE MEDICATION OVERVIEW

" ABCDD "

- A** ACE inhibitors
 - B** BETA Blockers
 - C** Calcium Channel Blockers
 - D** Digoxin
 - D** Diuretics
- SUFFIXES**
- PRIL
 - OLOL
 - PINE -AMIL


ENDOCRINE SYSTEM OVERVIEW



DIABETES TYPE 1 & 2

How the pancreas works (WITH FOOD)

Consume food
↓
Blood sugar increases
↓
This causes the pancreas to release insulin
↓
Insulin puts sugar & potassium into the cells!



How the pancreas works (WITH NO FOOD)

NO food
↓
Pancreas "back up plan"
↓
Glucagon hormone is released
↓
Breaks down stored glucose (glucagon) in the liver
↓
Releases glucose into the blood stream

TYPE 1 DIABETES

NO INSULIN PRODUCTION



Type **ONE** we have **NONE**

PATHOLOGY

- Caused by an autoimmune response
- The cells are starved of glucose since there is no insulin to bring it into the cells
- The cells break down protein and fat into energy causing ketones to build up = **ACIDOSIS!**
- Usually diagnosed in **childhood**



Easy to remember because **childhood** comes 1ST in life and **adulthood** comes 2ND

TYPE 2 DIABETES

DOES NOT PRODUCE ENOUGH INSULIN OR PRODUCES "BAD" INSULIN THAT DOES NOT WORK PROPERLY



Terrible **T**wo are **B**AD

PATHOLOGY

- Insulin resistance
- Insulin receptors are worn out & not working properly!
- Usually diagnosed in **adulthood** (due to a poor diet, sedentary lifestyle, and obesity)

ONSET: **ABRUPT**

TREATMENT

only has **1** treatment:
INSULIN

Oral hypoglycemic agents will not work for this pt.

Insulin dependent for life!

SIGNS & SYMPTOMS

Hyperglycemia

Glucose >115 & HbA1C 6.5 +



- Polyuria:** Excessive peeing
- Polydipsia:** Excessive thirst
- Polyphagia:** Excessive hunger

ONSET: **GRADUAL**

TREATMENT

has **2+** treatments:

DIET & EXERCISE
ORAL HYPOGLYCEMIC AGENTS

Example: Metformin

POSSIBLY INSULIN

Insulin is not administered routinely in a type 2 diabetes patient. Only in times of stress, surgery, or sickness will insulin need to be administered.

COMPLICATIONS

DIABETIC KETOACIDOSIS (DKA)

ONSET: **ABRUPT**

PATHOLOGY

Not enough insulin
↓
Blood sugar becomes **VERY** high
↓
Cells break down protein & fat into energy
↓
Ketones build up = **Acidosis!**

SIGNS & SYMPTOMS

- Ketosis & acidosis
- Hyperglycemia
- Dehydration
- Kussmaul respirations (trying to blow off CO₂)
- Acid breath "fruity breath"

TREATMENT

IV INSULIN • Fluid replacement
Correction of electrolyte imbalances

COMPLICATIONS

HYPERGLYCEMIC HYPEROSMOLAR NONKETOTIC SYNDROME (HHNS)

ONSET: **GRADUAL**

PATHOLOGY

NO acidosis present!
Simply high amounts of glucose in the blood

SIGNS & SYMPTOMS

Hyperglycemia
>600 +

TREATMENT

Fluid replacement
Correction of electrolyte imbalances
Possible Insulin administration

LONG TERM COMPLICATIONS



KIDNEY
NEPHROPATHY
Renal Failure



NERVES
RENAL NEUROPATHY
Loss of sensation



EYE
RETINOPATHY



HEART
HTN & ATHEROSCLEROSIS

HYPERGLYCEMIA VS. HYPOGLYCEMIA

HYPERGLYCEMIA

↑ BLOOD SUGAR


>200 mg/dL
Gradual (hours to days)

BLOOD GLUCOSE GOAL:
70 - 110 mg/dL

HYPOGLYCEMIA

↓ BLOOD SUGAR

<70 mg/dL
Happens suddenly


THE BRAIN NEEDS GLUCOSE... NO GLUCOSE CAUSES BRAIN DEATH!

3 P's
MOST COMMON SYMPTOMS

SIGNS & SYMPTOMS



- Polyuria
- Polydipsia
- Polyphagia
- Hot & dry skin
- Dry mouth (dehydration)
- Fruity breath
- Deep, rapid breaths (air hunger)
- Numbness & tingling
- Slow wound healing
- Vision changes

CAUSES

4 S's

- Sepsis (infection)
- Stress
- Steroids
- Skipping insulin or oral diabetic medication
- Not eating a diabetic diet

DIABETIC DIET

✓	✗
Complex carbohydrates Fiber-rich foods Heart-healthy fish "Good fats" Sugar-free fluids	Saturated fats Trans fats Cholesterol Sodium
	

TREATMENT

- Administer insulin as needed
- Test urine for ketones

	RAPID	SHORT	INTERMEDIATE	LONG
GENERIC NAMES	Lispro Aspart Glulisine	regular	nph	Glargine Detemir
BRAND NAMES	Humalog Novolog Apidra	Humulin R Novolin R	Humulin N Novolin N	Lantus Levemir

SIGNS & SYMPTOMS

- Cool & clammy skin
- Sweating (Diaphoresis)
- Palpitations
- Fatigue & weakness
- Confusion
- Headache
- Shakiness
- Inability to arouse from sleep
→ Can lead to coma ⚠️

CAUSES

- Exercise
 - Swimming, cycling, college athlete etc.
- Alcohol
- Peak times of insulin

RAPID INSULIN has the highest risk for Hypoglycemia

TREATMENT

CONSCIOUS PATIENTS

15 X 15 X 15

- Oral intake of **15 GRAMS** of carbohydrates
Juices, soda, low fat milk. NOT peanut butter or high fat milk
- Recheck blood glucose in **15 MIN**
- Give another **15 GRAMS** of carbohydrates if needed

UNCONSCIOUS PATIENTS


Do not put anything in an unconscious client's mouth, they can **ASPIRATE!**
Administer IV 50% dextrose (D50)
OR Glucagon (IM, IV, SubQ)

EMERGENCY
call a rapid response

THYROID DISORDERS

FUNCTION

- ☞ The thyroid gland produces 3 hormone (T3, T4, & Calcitonin)
 - You need **IODINE** to make these hormones
- ☞ Thyroid gives you **ENERGY!**



HYPERTHYROIDISM

PATHOLOGY

Excessive production of thyroid hormone

TOO MUCH ENERGY!


- Graves disease
- Too much Iodine (helps makes T3 + T4)
- Toxic Nodular Goiter
- Thyroid replacement medication (Toxicity)

LAB VALUES

↑ T3 & T4 ↓ TSH

SIGNS & SYMPTOMS

- Hyper-excitable
- Nervous/tremors
- Irritable
- ↓ Attention span
- Increased appetite
- Weight loss
- Hair loss
- Goiter (enlarged thyroid)
- Hot
- **EXOPHTHALMOS**
- Increased:
 - Blood pressure
 - Pulse
 - GI function



⚠️ LIFE-THREATENING COMPLICATIONS

⚡ THYROID STORM!
🚨 ACUTE / LIFE THREATENING EMERGENCY!

TREATMENT

- Anti-Thyroid Medications
 - Methimazole or PTU
- Beta Blockers (↓ HR & BP)
- Iodine Compounds
- Radioactive Iodine Therapy
- Thyroidectomy



HYPOTHYROIDISM

PATHOLOGY

Low production of thyroid hormone

NOT ENOUGH ENERGY!

MOST COMMON

- Hashimoto's disease
- Not enough Iodine
- Thyroidectomy
- Anti-thyroid medications
- Pituitary hormone
- Affects women more often than men

LAB VALUES

↓ T3 & T4 ↑ TSH

SIGNS & SYMPTOMS

- No energy
- Fatigue
- No expressions
- Weight gain
- Cold
- Amenorrhea
- Slurred speech
- Dry skin
- Coarse hair
- Decreased
 - HR
 - GI function (constipation)
 - Blood sugar (Hypoglycemia)

⚠️ LIFE-THREATENING COMPLICATIONS

MYXEDEMA COMA!


TREATMENT

- Hormone replacement (replacing levothyroxine)
 - Synthetic levothyroxine
 - Synthroid or Levothroid
 - Will be on this medication forever

*For more information about thyroid medications, see the Pharmacology Bundle

PARATHYROID GLAND DISORDERS

The parathyroid gland produces and secretes PTH (parathyroid hormone) which controls the levels of **CALCIUM** in the blood



↑ PTH

HYPERPARATHYROIDISM

↑ CALCIUM ↓ PHOSPHORUS

CAUSES

PRIMARY CAUSE:

Tumor or hyperplasia of the parathyroid

SECONDARY CAUSE:

Chronic kidney failure

SIGNS & SYMPTOMS

- **STONES:** Kidney stones (↑ calcium)
- **BONES:**
 - Skeletal pain
 - Pathological fractures from bone deformities
- Abdominal **MOANS**
 - Nausea, vomiting, and abdominal pain
 - Weight loss / anorexia
 - Constipation
- Psychic **GROANS**
 - Mental irritability
 - Confusion

**STONES,
BONES,
MOANS, &
GROANS**

TREATMENT

- Parathyroidectomy
- Removal of more than one gland
- Administer
 - Phosphates, calcitonin, & IV or oral bisphosphonates
- DIET: ↑ fiber & moderate calcium

↓ PTH

HYPOPARATHYROIDISM

↓ CALCIUM ↑ PHOSPHORUS

CAUSES

- Can occur due to accidental removal of the parathyroid
 - Thyroidectomy, parathyroidectomy, or radical neck dissection
- Genetic predisposition
- Exposure to radiation
- Magnesium depletion

SIGNS & SYMPTOMS

- Numbness & tingling
- Muscle cramps
- Tetany
- Hypotension
- Anxiety, irritability, & depression

Same S&S of hypocalcemia!

POSITIVE TROUSSEAU'S:

Carpal spasm caused by inflating a blood pressure cuff

CHVOSTEK'S SIGNS:

Contraction of facial muscles with light tap over the facial nerve

TREATMENT


- IV Calcium
- Phosphorus binding drugs
- DIET: ↑ Calcium ↓ Phosphorus

ADRENAL CORTEX DISORDERS

ADRENAL CORTEX HORMONES:

RETAINS: Na^+ & H_2O
LOSES: K^+

Glucocorticoids • Mineralocorticoids • Sex hormones



CUSHING'S

Disorder of the adrenal cortex
TOO MANY STEROIDS



They "have a **CUSHION**"

CAUSES

- Females
- Overuse of cortisol medications
- Tumor in the adrenal gland that secretes cortisol

SIGNS & SYMPTOMS

- Muscle wasting
- Moon face
- Buffalo hump
- Truncal obesity w/ thin extremities
- Supraclavicular fat pads
- Weight gain
- Hirsutism (masculine characteristics)
- ↑ Glucose ↑ Na^+
- ↓ K^+ ↓ Ca^{++}
- Hypertension

TREATMENT

- Adrenalectomy
 - Requires lifelong glucocorticoid replacement
- Avoid infection
- Adm. chemotherapeutic agents if adrenal tumor is present



ADDISON'S

Disorder of the adrenal cortex
NOT ENOUGH STEROIDS



We need to "**ADD**" some

CAUSES

- Surgical removal of both adrenal glands
- Infection of the adrenal glands
- TB, cytomegalovirus, & bacterial infections

SIGNS & SYMPTOMS

- Fatigue
- Nausea / vomiting / diarrhea
- Anorexia
- Hypotension & Hypovolemia!
- Confusion
- ↓ Blood sugar
- ↓ Na & H_2O ↑ K^+
- Hyperpigmentation of the skin
- Vitiligo: white areas of depigmentation



ADDISONIAN CRISIS

SIGNS & SYMPTOMS

- Profound fatigue
- Dehydration.....shock!
- Renal failure
- Vascular collapse
- Hyponatremia
- Hyperkalemia

TREATMENT

Fluid resuscitation & high-dose hydrocortisone

TREATMENT

- Adm. glucocorticoid and/or mineralocorticoid
- Diet: high in protein & carbs

PITUITARY GLAND DISORDERS

ANTIDIURETIC HORMONE (ADH):

ADH regulates & balances the amount of water in your blood



ADH is found in the **PITUITARY GLAND!**



SYNDROME OF INAPPROPRIATE ANTIDIURETIC HORMONE (SIADH)

SIADH is often of non-endocrine origin

TOO MUCH ADH

RETAINS WATER

INCREASED ICP

can lead to an ADH problem

CAUSES

- Pulmonary disease
 - ➔ TB
 - ➔ Severe pneumonia
- Disorders of the CNS
 - ➔ Head injury
 - ➔ Brain surgery
 - ➔ Tumor
- HIV
- Medications
 - ➔ Vincristine
 - ➔ Phenothiazines
 - ➔ Antidepressants
 - ➔ Thiazide diuretics
 - ➔ Anticonvulsants
 - ➔ Antidiabetic drugs
 - ➔ Nicotine

SIGNS & SYMPTOMS

- Low urinary output of concentrated urine
- Fluid volume overload
- Weight gain without edema
- Hypertension
- Tachycardia
- Nausea & vomiting
- Hyponatremia

TREATMENT

- Implement seizure precautions
- Elevate HOB to promote venous return
- Restrict fluid intake
- Adm. loop diuretics
- Adm. vasopressin antagonists



DIABETES INSIPIDUS (DI)



DI think Dry Inside

NOT ENOUGH ADH

LOSES WATER

CAUSES

- Head trauma, brain tumor
- Manipulation of the pituitary
 - ➔ Surgical ablation, craniotomy, sinus surgery, hypophysectomy
- Infections of the central nervous system (CNS)
 - ➔ Meningitis, encephalitis, or TB
- Failure of the renal tubules to respond to ADH

SIGNS & SYMPTOMS

- Excretes large amounts of diluted urine
- Polydipsia (increased thirst)
- Polyuria (increased urine output)
- Dehydration
- Decreased skin turgor
- Dry mucous membranes
- Muscle pain & weakness
- Headache
- Postural hypotension
- Tachycardia
- Low urinary specific gravity

Normal specific gravity: **1.005 - 1.030**

TREATMENT

- Adequate fluids
- IV hypotonic saline
- ADH replacement (replace the missing hormone!)
 - ➔ Vasopressin or desmopressin
- Monitor
 - ➔ Intake & output
 - ➔ Weight



ADRENAL MEDULLA DISORDER

ADRENAL MEDULLA HORMONES:

Epinephrine ▪ Norepinephrine




PHEOCHROMOCYTOMA

RARE tumor on the adrenal gland that secretes excessive amounts of epinephrine & norepinephrine

CAUSES

- Family history that makes them prone to developing the tumor



SIGNS & SYMPTOMS



- Hypertension (severe)
- Headache
- Heat (excessive sweating)
- Hypermetabolism
- Hyperglycemia

TREATMENT

- Adrenalectomy (if a tumor is present)
- Tell the client not to smoke, drink caffeine or change position suddenly
- Adm. anti-hypertensives
- Promote rest & calm environment
- Diet: high in calories, vitamins, & minerals



Avoid Stimuli!

It may cause a hypertensive crisis!


AUSCULTATING LUNG SOUNDS

TIPS FOR LISTENING


- 1 Listen directly on the skin with the diaphragm
- 2 Listening inside the **IN**tercostal spaces (*IN* between the ribs)
- 3 Listen to the anterior & posterior chest
- 4 Have the client sit upright (high fowler's), arms resting across the lap.
- 5 Instruct client to take deep breaths
- 6 Listen from top to bottom (comparing sides)

Listen for a
FULL INHALATION TO EXPIRATION
on each spot

ANTERIOR
Will hear
UPPER lobes well



POSTERIOR
Will hear
LOWER lobes well



NORMAL SOUNDS

BRONCHIAL (TRACHEAL)

DESCRIPTION

High, loud & hollow tubular

LOCATION HEARD

Anteriorly only
(heard over trachea & larynx)

DURATION

Inspiration < expiration



VESICULAR

DESCRIPTION

Soft, low pitched,
breezy / rushing sound

LOCATION HEARD

Heard anterior & posteriorly

DURATION

Inspiration > expiration



BRONCHOVESICULAR

DESCRIPTION


Medium pitched, hollow

LOCATION HEARD

Heard anterior & posteriorly

DURATION

Inspiration = expiration



ABNORMAL (ADVENTITIOUS) SOUNDS

DISCONTINUOUS SOUNDS

DISCRETE CRACKLING SOUNDS

FINE CRACKLES (RALES)

DESCRIPTION: High pitched, crackling sounds
(Sound like fire crackling, or velcro coming part)

DUE TO: Previously deflated airways that are popping back open

EXAMPLE: Pulmonary edema, asthma, obstructive diseases

COARSE CRACKLES (RALES)

DESCRIPTION: Low pitched, wet bubbling sound

DUE TO: Inhaled air collides with secretion in the trachea or large bronchi

EXAMPLE: Pulmonary edema, pneumonia, depressed cough reflex

PLEURAL FRICTION RUB

DESCRIPTION: Low pitched, harsh / grating sounds

DUE TO: Pleura is inflamed and loses its lubricant fluid.
It's literally the surfaces rubbing together during respirations

EXAMPLE: Pleuritis

CONTINUOUS SOUNDS

CONNECTED MUSICAL SOUNDS

WHEEZES

DESCRIPTION: High-pitched musical instrument with more than one type of sound quality (polyphonic)

DUE TO: Air moving through a narrow airway

EXAMPLE: Asthma, bronchitis, chronic emphysema

STRIDOR

DESCRIPTION: High pitched whistling or gasping with harsh sound quality

DUE TO: Disturbed airflow in larynx or trachea

EXAMPLE: Croup, epiglottitis, any airway obstruction

REQUIRES MEDICAL ATTENTION

CHRONIC OBSTRUCTIVE PULMONARY DISEASE (COPD)

PATHOLOGY

Pulmonary disease that causes chronic airflow obstruction



EMPHYSEMA or **CHRONIC BRONCHITIS**

OTHER FACTS

- COPD is a progressive disorder which means the disease gets worse over time; it's irreversible!
- Alveoli sac lose their elasticity (inability to fully exhale)

DIAGNOSTIC

- Arterial blood gases (ABG's)
- Chest x-ray
- Pulmonary function test: Spirometry

Obstructive lung disease
FEV1 / **FVC** ratio of less than 70%

FEV1
 =
FORCED EXPIRATORY VOLUME

FVC
 =
FORCED VITAL CAPACITY

RISK FACTORS

- Smoking **MOST COMMON**
 ↳ Breathing in harmful irritants
- Occupation exposure
- Infection
- Air pollution
- Genetic abnormalities
- Asthma
- Severe respiratory infection in childhood

Deficiency of Alpha1- antitrypsin
 (Protects the lining of the lungs)

EMPHYSEMA VS CHRONIC BRONCHITIS

EMPHYSEMA

Abnormal distention of airspaces

Enlargement & destruction of airspace distal to the terminal bronchiole

Hyperventilation (breathing fast)
 Trying to blow off CO₂

LIMITED AIRFLOW

↓ O₂
 &
 ↑ CO₂

Mucus secretion


Airway obstruction (inflammation)

Chronic productive cough & sputum production for >3 months (within 2 consecutive years)

SIGNS & SYMPTOMS

"PINK PUFFERS"

- Hyperinflation of the lungs (barrel chest)
- Thin - weight loss
 ↳ Burning a lot of calories from breathing a lot!
- Shortness of breath
- Severe dyspnea



SIGNS & SYMPTOMS

"BLUE BLOATERS"

- Overweight
- Cyanotic (blue) - Hypoxemia
 ↳ ↓ O₂ & ↑ CO₂
- Peripheral edema
- Rhonchi & wheezing
- Chronic cough

CHRONIC OBSTRUCTIVE PULMONARY DISEASE (COPD)

NURSING MANAGEMENT & EDUCATION

MONITOR RESPIRATORY SYSTEM

- * Lung sounds
- * Sputum production
- * Oxygen status

LIFESTYLE MODIFICATIONS

- * Smoking cessation
 - ➔ Determine readiness
 - ➔ Develop a plan
 - ➔ Discuss nicotine replacement

DIET MODIFICATIONS

- * Promote nutrition
- * Increase calories
- * Small frequent meals
- * Stay hydrated
 - ➔ Thins mucous secretions

TEACH PROPER BREATHING TECHNIQUES

- * Pursed lips
- * Diaphragmatic breathing

SURGERY

- * Bullectomy
- * LVRS: lung volume reduction surgery
- * Lung transplant

STAY UP TO DATE ON VACCINES

- * Influenza & pneumococcal vaccine
 - ↓ the incidence of pneumonia

OXYGEN THERAPY

- COPD clients are stimulated to breathe due to ↓ O₂ (if you give too much O₂...they lose their "drive to breathe")
- Healthy clients are stimulated to breathe due to ↑ CO₂

Adm. O₂ during exacerbations or showing signs of respiratory distress


Adm. oxygen with caution to clients with **CHRONIC HYPERCAPNIA** (elevated PaCO₂ levels)
1 - 2 liters max

Clients with COPD (especially emphysema) are using a lot of their energy to breathe, therefore burning a lot of calories



PROMOTES CARBON DIOXIDE ELIMINATION

Allows better expiration by ↑ airway pressure that keeps air passages open during exhalation!



We want to use the **DIAPHRAGM** rather than the accessory muscles to breathe!

- ➔ This strengthens the diaphragm and slows down breathing rate



MEDICATION

BRONCHODILATORS

- * Relaxes smooth muscle of lung airways = better airflow
- * Symbicort (steroid + long-acting bronchodilator)

CORTICOSTEROIDS

- * ↓ inflammation (oral, IV, inhaled)
- * Example: Prednisone, Solumedrol, Budesonide

BUPROPION (ANTI-DEPRESSANT)

*For more information about respiratory medications, see the Pharmacology Bundle

SUFFIX:
"-asone"
"-inide"
"-olone"

ORDER OF EVENTS

- 1 Bronchodilator**
Dilated airways
- 2 Corticosteroids**
Airways are open now in order for the steroids to do its job!



PNEUMONIA

PATHOLOGY

Lower respiratory tract infection that causes inflammation of **ALVEOLI SACS!**

REMEMBER

gas exchange takes place in the alveoli... so pneumonia causes **impaired gas exchange.**



SYMPTOMS

- * ↑ Temperature: **mild - high** fever
- * ↑ HR
- * ↑ RR
 - * Attempting to blow off CO₂
- * ↓ O₂ saturation
- * Chills
- * Chest pain
- * Difficulty breathing
- * Productive cough
- * Unusual breath sounds: coarse crackles & wheezes
- * Respiratory acidosis
- * ↑ CO₂ ↓ O₂

RISK FACTORS

Can be **COMMUNITY-ACQUIRED** or **HOSPITAL-ACQUIRED!**

- * Prior infection
- * Immunocompromised
 - * HIV, young/old, auto immune infections
- * Postoperative
- * Lung diseases
 - * COPD
- * Immobility
- * Aspiration risk

INTERVENTIONS

- * Monitor...
 - * Respiratory status
 - * Vital signs: HR, temp, & pulse oximetry
 - * Color, consistency & amount of sputum
- * Diet
 - * ↑ Calorie
 - * ↑ Protein
 - * ↑ Fluids (oral or IV)
 - * Small frequent meals
- * Medications
 - * Antipyretics
 - * Antibiotics (*only for bacteria*)
 - * Antivirals
 - * Bronchodilators
 - * Cough suppressants
 - * Mucolytic agents
- * Semi Fowler's position
 - Helps lung expansion

Thins secretions & compensates dehydration from fever

DIAGNOSTIC

Chest X-ray * ↑ White blood cells * Sputum culture

↓
shows pulmonary infiltrates or pleural effusions

↓
can be BACTERIAL, VIRAL, or FUNGAL

EDUCATE

- * Use of Incentive Spirometer
 - * Helps to pop open the alveoli sacs & get the air moving
- * Up to date vaccines
 - * Annual flu shot
 - * Pneumococcal vaccine
- * Smoking cessation
- * Hand washing & avoiding sick people!



ASTHMA


PATHOLOGY

Chronic lung disease that causes an inflamed, narrow, & swollen airway (bronchi & bronchioles)

CAUSES

NOT COMPLETELY KNOWN!

- Genetic
- Environmental
 - Smoke, pollen, perfumes, dust mites, pet dander, cold or dry air, etc.
- GERD
- Exercise-induced asthma
- Certain drugs
 - NSAIDs, aspirin



CLASSIFICATIONS BASED ON SYMPTOMS

MILD INTERMITTENT

< 2 a week

MILD PERSISTENT

> 2 a week
Not daily

MODERATE PERSISTENT

Daily symptoms & exacerbations that happen 2x a week

SEVERE PERSISTENT

Continually showing symptoms with frequent exacerbations

SIGNS & SYMPTOMS

CHARACTERIZED BY FLARE-UPS
(meaning: it comes & goes)

- Dyspnea (shortness of breath)
- Tachypnea (fast respiratory rate)
- Chest tightness
- Anxiety
- Wheezing
- Coughing
- Mucus production
- Use of accessory muscles
- AIR TRAPPING**

Air trapping causes the client to retain CO₂ which is **ACIDIC** = **RESPIRATORY ACIDOSIS**

NURSING CARE

- Assess client's airway
- High Fowler's position
- Provide frequent rest periods
- Adm. oxygen therapy
 - Goal: keep the O₂ at 95 - 100%
- Maintain a calm environment to ↓ stress
- Asses **PEAK FLOW METER** reading
- Asses for cyanosis & retractions





STATUS ASTHMATICUS

Life-threatening asthma episode
Medical emergency!

- OXYGEN
- ↓
- HYDRATION
- ↓
- NEBULIZATION
- ↓
- SYSTEMIC CORTICOSTEROID

MEDICATIONS

- BronchoDILATORS**
 - Short-acting (Albuterol) → **RAPID RELIEF**
 - Long-acting (Salmeterol) → **PREVENTS ASTHMA ATTACKS**
 - Methylxanthines (Theophylline)
- Corticosteroids**
 - Suffix **-ASONE & -IDE** → **ANTI-INFLAMMATORY AGENTS**
 - Ex: Beclomethasone
- Leukotriene Modifiers
- Anticholinergics



PEAK FLOW METER

- Shows how controlled the asthma is & if it's getting worse
- Establish a baseline by performing a "personal best" reading
 - Client will exhale as hard as they can & get a reading

GREEN = GOOD
YELLOW = NOT TOO GOOD
RED = BAD

*For more information about respiratory medications, see the Pharmacology Bundle

IRON DEFICIENCY ANEMIA

MOST COMMON TYPE OF ANEMIA

PATHOLOGY

TYPE OF ANEMIA CAUSED BY ↓ IRON LEVELS

- Iron is **ESSENTIAL** to hemoglobin in red blood cells.
- The body uses **IRON** to make hemoglobin. Hemoglobin carries oxygen to the cells!

RED BLOOD CELLS ROLE

Transports O₂ & removes CO₂ from the body with the help of hemoglobin (Hgb)



HEMOGLOBIN (HGB)

Found in the RBC's
It's a protein that contains IRON

CAUSES

- Blood loss / hemorrhage
- Malabsorption
- Inadequate dietary intake of iron

SYMPTOMS

- Pallor
- Weakness & fatigue
- Microcytic (small) red blood cells
- ↓ hemoglobin & ↓ hematocrit



NORMAL VALUES

Hemoglobin (Hgb)

Female: 12 - 16 g/dL Male: 13 - 18 g/dL

Hematocrit (HCT)

Female: 36% - 48% Male: 39% - 54%

IRON-RICH FOODS




EAT
Egg Yolks
Apricots
Tofu

LOTS
Legumes
Oysters
Tuna
Seeds

POF
Potatoes
Fish



IRON
Iron-fortified cereals
Red meats
Poultry
Nuts



INTERVENTIONS

- Diet changes
- ↑ Iron
- ↑ Protein
- ↑ Vitamins
- Administer iron
 - Oral, IM, or IV
- D/C any damaging drugs
- If active bleed is suspected, identify cause & control bleeding!

Administering Iron Supplements

↓ ABSORPTION

Calcium:

Milk & antacids

↑ ABSORPTION

Vitamin C:

Fruit juice & multivitamin

Liquid iron stains the teeth!

1. Take with a straw

2. Brush teeth after



Side Effects of Iron Supplements

- Black stool
- Constipation
- Foul aftertaste

THROMBOCYTOPENIA

PATHOLOGY

↓ PLATELETS


👉 Platelets help clot the blood

👉 Platelet aggregation

➡ The clumping together of platelets that form a plug at the site of the injury

👉 ↓ platelets = think **BLEEDING**

Normal Platelet Count
150,000 - 450,000 per microliter



Thrombocytopenia
< 150,000

CAUSES

- P** latelet disorders
- L** eukemia
- A** nemia
- T** rauma
- E** nlarged spleen
- L** iver disease
- E** thanol (alcohol-induced)
- T** oxins (drug-induced)
- S** epsis

SYMPTOMS

- ◆ Weakness, dizziness, tachycardia, hypotension
- ◆ Prolonged bleeding time
- ◆ Petechiae (pinpoint bleeding)
- ◆ Purpura (bruising)
- ◆ Bleeding from the gums & nose
- ◆ Heavy menstrual cycles
- ◆ Blood in stool or urine
- ◆ ↑ INR & ↑ PT/PTT

DIAGNOSIS

- ◆ Bleeding time
- ◆ aPTT - Activated partial thromboplastin time
- ◆ PT - Prothrombin time
- ◆ INR - International normalized ratio
- ◆ ↓ Hgb & Hct

NURSING MANAGEMENT

- ◆ Platelet transfusion
- ◆ Bone marrow transplant
 - Platelets are made in the bone marrow
- ◆ Splenectomy
 - For those unresponsive to medical therapy

BLEEDING PRECAUTIONS



- ◆ Use electric razors
- ◆ NO aspirin
- ◆ Use small needle gauges
- ◆ Decrease needle sticks
- ◆ Protect from injury



IMMUNE THROMBOCYTOPENIC PURPURA (ITP)

Formerly called "idiopathic thrombocytopenia purpura"

PATHOLOGY

Autoimmune disease where the body produces antibodies against its own thrombocytes (Platelets)

"Purpura" is in the name because it causes easy bruising & petechiae in the trunk & extremities!

ITP
< 20,000

CAUSES

- ◆ Children after viral illness
- ◆ Females (ages 20 - 40)
- ◆ Pregnancy

HEMATOLOGY LAB VALUES

EXPECTED RANGE

DESCRIPTION



	EXPECTED RANGE	DESCRIPTION	↓	↑
RED BLOOD CELLS (RBCs)	<p>F 4.2 – 5.2 X 10⁶ / uL</p> <p>M 4.7 – 6.1 X 10⁶ / uL</p>	<p>Red blood cells transport oxygen to the body's cells.</p>	<ul style="list-style-type: none"> Fluid volume overload Hemorrhage Anemia Renal disease (lack of erythropoietin production) <p>more volume dilutes the RBCs</p>	<p>Less volume concentrates the RBCs</p> <ul style="list-style-type: none"> Dehydration /fluid volume deficit Hyperactivity of the bone marrow
WHITE BLOOD CELLS (WBCs)	<p>4,500 – 11,000 / uL</p>	<p>The white blood cells are a part of the immune system and help to fight infections and diseases.</p>	<p>LEUKOPENIA</p> <p>WBCs < 4,500 /uL</p> <ul style="list-style-type: none"> Immunosuppression 	<p>LEUKOCYTOSIS</p> <p>WBCs > 11,000 /uL</p> <ul style="list-style-type: none"> Current or recent INFECTION & inflammation
PLATELETS (PLT)	<p>150,000 – 450,000 /uL</p>	<p>Platelets help clot the blood. Platelet aggregation is the clumping together of platelets that form a plug at the site of the injury.</p>	<p>THROMBOCYTOPENIA</p> <p>PLTs < 150,000 /uL</p> <p>↓ Platelets think BLEEDING</p>	<p>THROMBOCYTOSIS</p> <p>PLTs > 450,000 /uL</p> <ul style="list-style-type: none"> Certain cancers Infection
HEMOGLOBIN (HGB)	<p>F 12 – 16 g/dL</p> <p>M 13 – 18 g/dL</p>	<p>Hemoglobin is an iron containing protein found in red blood cells. It transports oxygen from the lungs to the tissues. It also returns CO₂ from the tissues back to the lungs.</p>	<ul style="list-style-type: none"> Fluid retention (hemodilution) Anemia Hemorrhage 	<ul style="list-style-type: none"> Dehydration (hemoconcentration)
HEMATOCRIT (HCT)	<p>F 36% – 48%</p> <p>M 39% – 54%</p>	<p>The percent of blood that is made up of red blood cells (expressed as a %).</p>	<ul style="list-style-type: none"> Fluid retention (hemodilution) Anemia Hemorrhage 	<ul style="list-style-type: none"> Dehydration (hemoconcentration) Low oxygen availability (smoking, pulmonary diseases (COPD), high altitudes)
ACTIVATED PARTIAL THROMBOPLASTIN TIME (aPTT)	<p>NORMAL (not on anticoagulants) 30 – 40 seconds</p> <p>ON HEPARIN THERAPY 1.5 – 2.0 x the normal value</p>	<p>aPTT measures how long it takes for a blood clot to form. It's also used to monitor the effectiveness of the anticoagulant: HEPARIN.</p>	<ul style="list-style-type: none"> Hypercoagulable <p>MEMORY TRICK Numbers are LOW = (lots will GROW)</p>	<ul style="list-style-type: none"> Heparin therapy <p>MEMORY TRICK Numbers are too HIGH = Patient will DIE (from increased bleeding)</p>
PROTHROMBIN TIME (PT)	<p>NORMAL (not on anticoagulants) 10 – 12 seconds</p> <p>ON HEPARIN THERAPY 1.5 – 2.0 x the normal value</p>	<p>Prothrombin time measures the amount of time needed to form a clot. It's also used to monitor the effectiveness of the anticoagulant: WARFARIN.</p>		<ul style="list-style-type: none"> Deficiency in vitamin K Deficiency in clotting factor Liver disease
INTERNATIONAL NORMALIZED RATIO (INR)	<p>NORMAL (not on anticoagulants) < 1</p> <p>ON HEPARIN THERAPY INR 2.0 – 3.0 INR 2.5 – 3.5 (heart valve replacement)</p>	<p>INR is calculated from the prothrombin time and is used to monitor oral anticoagulants such as WARFARIN.</p>	<p>MEMORY TRICK Numbers are LOW = (lots will GROW)</p>	<ul style="list-style-type: none"> Warfarin therapy <p>MEMORY TRICK Numbers are too HIGH = Patient will DIE (from increased bleeding)</p>
D-DIMER	<p>< 0.5 mcg/mL</p>	<p>D-dimers are fragments of fibrin that are in the blood when a clot dissolves or is broken down. D-dimer helps to determine if a clot is present somewhere in the body.</p>	<ul style="list-style-type: none"> Blood clot is ruled out 	<p>Additional tests are needed to confirm and determine a specific diagnosis</p> <ul style="list-style-type: none"> Blood clot may be present in the body

ACUTE & CHRONIC PANCREATITIS

PATHO

The islets of Langerhans secrete **INSULIN & GLUCAGON INTO THE BLOOD STREAM**

Pancreatic tissue: secrete digestive enzymes that break down **CARBOHYDRATES, PROTEINS & FATS**

PANCREATITIS is an **AUTO-DIGESTION** of the pancreas by its own digestive enzymes released too early in the pancreas

LABS


- ↑ Amylase
- ↑ Lipase
- ↑ WBCs
- ↑ Bilirubin
- ↑ Glucose
- ↓ Platelets
- ↓ Ca & Mg

CAUSES

- Gallstones
 - Blocks the bile duct
- Alcohol (ETOH)
 - Damages the cells of the pancreas
- Infection
- Medications
- Tumor
- Trauma

SIGNS & SYMPTOMS

- In **ACUTE**, there will still be working functions of the pancreas.
- Sudden sever **PAIN!**
 - Mid-epigastric pain LUQ
 - Nausea & vomiting
 - Fever
 - ↑ HR & ↓ BP
 - ↑ Glucose
 - Mental confusion & agitation
 - Abdominal guarding
 - Rigid/board-like abdomen
 - Grey-Turner's Sign
 - Bluish discoloration at the flanks!
 - Cullen's Sign
 - Bluish discoloration of the umbilicus
 - Cullens = Circle belly button



VS

ACUTE

Sudden inflammation that is **REVERSIBLE** if prompt recognition and treatment is done

CHRONIC

Chronic inflammation that is **IRREVERSIBLE**

- Repeated episodes of acute pancreatitis
- Excessive & prolonged consumption of alcohol (ETOH)
 - Recurrent damage to the cells of the pancreas
- Cystic Fibrosis

In **CHRONIC**, you will see different S&S due to the prolonged damage & loss of function

- Chronic epigastric pain or no pain
- Pain ↑ after drinking ETOH or after a fatty meal
- Steatorrhea "fatty stools"
 - Oily/greasy frothy stool
- Weight loss
 - Can't digest food properly
- Jaundice
 - Yellowish color of the skin from build up of bile
- Diabetes Mellitus
 - Damage to the islet of Langerhans
- Dark urine
 - From excess bile in the body

MEDICATIONS

- Opioid analgesics
- Antibiotics
- Proton Pump Inhibitors (PPI's), H2 antagonists, antacids
- Pancreatic enzymes
- Insulin

DIGESTIVE ENZYMES (EXOCRINE)

AMYLASE: Breaks down carbs to **glucose**

PROTEASE: Breaks down **proteins**

LIPASE: Breaks down **fats**

DIET

- NO ETOH!
- ↑ protein
- Limit sugars
- ↓ fat (no greasy, fatty foods)
- Complex carbohydrate (fruits, vegetables, grains)



INTERVENTIONS

- Rest the pancreas!
 - NPO (we don't want stimulation of the enzymes)
- IV fluids
- Pain management
- Positioning
 - Side lying → fetal position, NOT supine!
- Insert NG tube
 - Remove stomach contents



INFLAMMATORY BOWEL DISEASE (IBD)

CROHN'S DISEASE


PATHO

Inflammation that occurs anywhere in the GI tract (mouth - anus)

NO CURE!

SIGNS & SYMPTOMS

- Cobble-stone appearance
- Fever
- Cramping after meals
- Mucus like diarrhea (semisolid)
- Abdominal distention
- Nausea & vomiting



ULCERATIVE COLITIS

PATHO

Inflammation & **ulceration** of only the large intestine & rectum

CURE: Colectomy with ileostomy

SIGNS & SYMPTOMS

- Ulcers cause
 - Rectal bleeding
 - Bloody diarrhea
 - Abdominal cramping
- ↑ HR & ↓ BP
 - Hypovolemic shock
- Malnutrition
- Malaise
- Dehydration
- Vitamin K deficiency

INTERVENTIONS

for the Acute Phase

Adm. fluids, electrolytes or parenteral nutrition

NPO

DIET

- Clear liquids to ↓ fiber
- ↑ Protein
- Vitamins & iron supplements
- Avoid gas-forming foods

AVOID SMOKING

MEDICATIONS

Dairy
Fruits & vegetables

Whole-wheat grains

Alcohol

Nuts


Caffeine

Corticosteroids
Immunosuppressants
Antidiarrheals
Salicylate compounds

MONITOR

- Bowel sounds
- Bowel perforation
- Peritonitis
- Hemorrhage
 - Stool
 - Color
 - Consistency
 - Presence of blood

TYPES OF HEPATITIS



HEPATITIS
↓
LIVER INFLAMMATION
"INFLAMMATION OF THE LIVER"

CAUSED BY:

- **VIRAL (A, B, C, D, E)** **MOST COMMON**
- EXCESSIVE USE OF ALCOHOL
- HEPATOTOXIC MEDICATIONS

	TRANSMISSION	SIGNS & SYMPTOMS	DIAGNOSTIC	TREATMENT	VACCINE
<p>HAV ACUTE ONLY</p>	<p>Fecal & oral</p> <ul style="list-style-type: none"> • Food & water 	<p>GI symptoms (N&V, stomach pain, anorexia)</p> <p>Dark-colored urine</p> <p>Clay-colored stool</p> <p>Vomiting</p> <p>Flu-like symptoms</p> <p>Jaundice</p> <p>↓</p> <p>YELLOW DISCOLORATION of the skin from the buildup of bilirubin</p>	<p>Anti-HAV</p> <p>IgM = Active infection</p> <p>IgG = Recovered (It's Gone)</p>	<p>Supportive therapy... REST!</p>	<p>✓</p>
<p>HBV B IS BOTH ACUTE & CHRONIC</p>	<p>B think Body fluids (Semen, saliva)</p> <ul style="list-style-type: none"> • Birth & blood • Childbirth, sex, & IV drugs 		<p>HBsAG = Active infection</p> <p>Anti-HBs = Immune / recovery</p>	<p>ACUTE Supportive therapy & rest</p> <p>CHRONIC Antivirals</p>	<p>✓</p>
<p>HCV ACUTE & CHRONIC</p>	<p>Body fluids</p> <ul style="list-style-type: none"> • Most common: IV drug users 		<p>Anti-HCV</p> <p>No post exposure immunoglobulin</p>	<p>Antivirals</p> <p>Interferon</p>	<p>✗</p>
<p>HDV ACUTE & CHRONIC</p>	<p>Depends on B</p> <p>B & D = BuDs</p> <p>Hep D occurs with Hep B</p>		<p>HDAg</p> <p>Anti-HDV</p>	<p>Antivirals</p> <p>Interferon</p>	<p>✗</p>
<p>HEV ACUTE ONLY</p>	<p>Fecal & oral</p> <ul style="list-style-type: none"> • Food & water uncooked meats, 3rd world countries 		<p>Anti-HEV</p>	<p>Supportive therapy... REST!</p>	<p>✗</p>

EDUCATION FOR ALL TYPES OF HEPATITIS!

- Rest
- Diet
- Small frequent meals
- ↑ Carbohydrates
- ↑ Calories
- ↓ Protein & fat
- Proper hand hygiene
- Do not share personal hygiene products
- Avoid sex until hepatitis antibodies are negative
- Educate on toxic substances to avoid
- Alcohol, acetaminophen, aspirin, sedatives

LABS:

Liver enzymes
ALT: 7 - 56 U/L
AST: 5 - 40 U/L

Bilirubin: <1 mg/dL

Ammonia: 15 - 45 mcg/dL


ALL WILL BE ELEVATED IN HEPATITIS

CIRRHOSIS

FUNCTIONS OF A HEALTHY LIVER

- 1 **DETOX THE BODY**
- 2 **HELPS TO CLOT THE BLOOD**
- 3 **HELPS TO METABOLIZE (BREAKDOWN) DRUGS**
- 4 **SYNTHESIS (MAKES) ALBUMIN**

STAGES OF LIVER DAMAGE



PATHOLOGY

Liver cells are **DESTROYED** and replaced with fibrotic (**scar**) tissue.

Loss of normal function of the liver.

CAUSES

- ETOH consumption
- Nonalcoholic fatty liver disease (NAFLD)
 - Viral hepatitis B & C
 - Autoimmune
 - Hepatotoxic drugs
 - Toxins & parasites
 - Fat collection in the liver (obesity, diabetes, ↑ cholesterol)

SIGNS & SYMPTOMS

- Asterixis
 - Liver flap
- Jaundice
 - Yellow discoloration in the eyes & skin
- Ascites
- Edema
- Abdominal pain
- Chronic dyspepsia (GI upset)
- Itchy skin
- ↑ Bilirubin & ammonia
- ↓ Platelets
 - Risk for bleeding
- ↓ WBC's
 - Risk for infection

COMPLICATIONS

- Portal HTN
 - Portal veins become narrow due to scar tissue
- GI bleeding (esophageal varices)
- Splenomegaly
- Anemia
- Hepatic encephalopathy/coma
 - Due to ↑ ammonia levels (ammonia is a sedative)
- Gynecomastia
 - Breast development in men
- Hepatorenal Syndrome
 - Acute kidney injury *in clients with liver failure*

TREATMENT

- No more alcohol
- Rest
- Prevent bleeding
 - Bleeding precautions
- Measure abdominal girth
- Daily weights & I&O's
- Electric razor
- Soft-bristled tooth brush
- Pressure on all venipuncture
- Paracentesis
 - Removal of fluid from the peritoneal cavity (ascites)
- Liver transplant

MEDICATIONS

- Antacids
- Vitamins
- Diuretics
- Lactulose
 - ↓ serum ammonia through the stool
- Avoid narcotics

THE LIVER CAN'T METABOLIZE DRUGS WELL WHEN IT'S SICK



DO NOT GIVE ACETAMINOPHEN TO PEOPLE WITH LIVER ISSUES!

NEUROLOGICAL ASSESSMENTS

LEVEL OF CONSCIOUSNESS (LOC)

Level of CONSCIOUSNESS (LOC) is always **#1** with neurological assessment

A change in LOC may be the only sign that there is a **PROBLEM!**



PUPILLARY CHANGES

PERRLA

Pupils, **E**qual, **R**ound, **R**eactive to **L**ight & **A**ccommodation



NORMAL PUPIL SIZE : 2 - 6 mm

GLASGOW COMA SCALE

TOOL FOR ASSESSING A CLIENT'S RESPONSE TO STIMULI

EYE OPENING RESPONSE	Spontaneous	4
	To speech	3
	To pain	2
	No response	1
VERBAL RESPONSE	Oriented	5
	Confused	4
	Inappropriate words	3
	Unclear sounds	2
MOTOR RESPONSE	None	1
	Obeys command	6
	Localizes pain	5
	Withdraws	4
	Flexion	3
	Extension	2
TOTAL	None	1
		3 - 15

INTERPRETATION

WORST 3 Severe impairment of neurological function, coma, or brain death

<8 Unconscious patient

BEST 15 Fully alert & oriented

MENTAL STATUS

- ARE THEY AWARE OF THEIR SURROUNDINGS?**
- ARE THEY ORIENTED TO PERSON, PLACE, TIME, & SITUATION?**
- DO THEY HAVE THEIR SHORT TERM & LONG TERM MEMORY?**

Ask these types of questions to assess mental status:

- What is your name?
- Do you know where you are?
- Do you know what month it is?
- Who is the current U.S. president?
- What are you doing here?



DEEP TENDON REFLEX (DTR) RESPONSES

- 0** = No response **ABSENT**
- 1+** = Present, but sluggish or diminished
- 2+** = Active or expected response **NORMAL**
- 3+** = More brisk than excited; Hyperactive
- 4+** = Brisk, Hyperactive, with intermittent, or transient clonus



BABINSKI REFLEX (PLANTAR REFLEX)

ELICITED BY STROKING THE LATERAL SIDE OF THE FOOT



INTACT CNS

The lateral sole of the foot is stroked and the toes contract & draw together.

BRAIN DYSFUNCTION

Toes fan out when stroked. Remember this is only normal in newborns & infants up to 2 years of age, but abnormal in adults!

Babinski think:

Normal in **B**abies & the **B**ig toe fans out

SEIZURES


WHAT IS A SEIZURE? Abnormal & sudden electrical activity of the brain

WHAT IS EPILEPSY? Chronic seizure activity due to a chronic condition

CAUSES

- ↑ fever (Febrile seizure in child)
- CNS infection
- Drug or alcohol withdrawal
- ABG imbalance
- Hypoxia
- Brain tumor
- Hypoglycemia
- Head injury
- Hypertension

STAGES OF A SEIZURE



GENERALIZED SEIZURES



THE ENTIRE BRAIN IS AFFECTED

TONIC-CLONIC

"Used to be called grand-mal" May begin with an aura. Stiffening (tonic) and/or rigidity (clonic) of the muscles.

MYOCLONIC

Sudden jerking or stiffening of the extremities (arms or legs).

ABSENCE

Usually looks like a blank stare that lasts seconds. Often goes unnoticed

ATONIC

Sudden loss of muscle tone. May lead to sudden falls or dropping things.

PARTIAL (FOCAL) SEIZURES



ONE AREA OF THE BRAIN IS AFFECTED

SIMPLE PARTIAL


Sensory symptoms with motor symptoms and stays aware. They may report an aura.

COMPLEX PARTIAL

Altered behavior/awareness and loses consciousness for a few seconds.

CARE DURING THE SEIZURE

SEIZURE PRECAUTIONS



DON'T

- Restrain the client
- Force the jaw open
- Place anything in their mouths
- Leave the client

CEREBROVASCULAR ACCIDENT (CVA) "STROKE"

ISCHEMIC STROKE

BLOCKAGE

"Thrombotic or embolic"

THROMBOSIS: blood clot that formed on the artery wall

EMBOLISM: A clot has left part of the body

Blood flow is cut off which leads to **ISCHEMIA**.

TRANSIENT ISCHEMIC ATTACKS: "TIA'S"

"Mini strokes"

The same pathology as a stroke but no cerebral infarction occurs

HEMORRHAGIC STROKE

BLEEDING

RUPTURED ARTERY

ANEURYSM (weakening of the vessel)

UNCONTROLLED HYPERTENSION

The collection of blood in the brain leads to ischemia & increased ICP

TREATMENT

FIBRINOLYTIC THERAPY

(TPA) TISSUE PLASMINOGEN ACTIVATOR

DISSOLVES DOWN THE BLOOD CLOT!

- Avoid IM injections
- Avoid unnecessary IV punctures
- Prevent injury (bed rest)
- Check for bleeding

STOP THE BLEEDING PREVENT INCREASED ICP

- Poor prognosis
- Needs careful monitoring in an intensive care unit
- Blood may need to be removed to ↓ pressure on the brain

SIGNS & SYMPTOMS

Face drooping

- Uneven smile

Arm weakness

- Arm numbness; can't lift arm

Speech difficulty

- Slurred speech

Time to call 911

REMEMBER!

If the stroke occurs on the left side of the brain, the right side of the body will be affected

TYPES OF APHASIA

RECEPTIVE

Unable to comprehend speech
(**WERNICKE'S AREA**)

EXPRESSIVE

Can comprehend speech (but can't respond back with speech)
(**BROCA'S AREA**)

RISK FACTORS

MODIFIABLE

- Hypertension
- Atherosclerosis
- Anticoagulation therapy
- Diabetes Mellitus
- Obesity
- Stress
- Oral contraceptives

NON-MODIFIABLE

- Family history of strokes
- Older age
- Male gender
- Black
- Hispanic

NURSING MANAGEMENT

• Assist with safe feeding

- Do not feed until gag reflex has come back
- ↓ chances of aspiration
- Keep suction at the bedside
- Crush medications

• Positioning of the client

- Elevate head of the bed to ↓ ICP
- Place a pillow under the affected arm in a neutral position

• Assist with communication skills

- Encourage passive range of motion every 2 hours
- Preventative DVT measures
- Assist with Activities of Daily Living (ADL's)
- Communication
 - Be patient
 - Make clear statements
 - Ask simple questions
 - Don't rush!

PREVENTATIVE DVT MEASURES

- Compression stockings
- Frequent position change
- Mobilization

- Frequent rest periods
- Dress the affected side first
- Support affected side

LIQUID

- Thin
- Nectar-like
- Honey-like
- Spoon-thick

FOOD

- Pureed
- Mechanically altered
- Mechanically softened
- Regular

CRANIAL NERVES

WHAT ARE CRANIAL NERVES?

Nerves that originate from the brain stem. They send information to & from various parts of the body.

SE SENSORY
M MOTOR
B BOTH



XII: HYPOGLOSSAL **M**

FUNCTION: **GLOSSO MEANS TONGUE!**

Tongue movement (swallowing & speech)

TEST:

Inspect tongue & ask to stick tongue out

XI: SPINAL ACCESSORY **M**

FUNCTION:

Controls strength of neck & shoulder muscles

TEST:

Ask the client to rotate their head & shrug their shoulders

X: VAGUS **B**

FUNCTION:

MOTOR - Swallowing, speaking, & cough
SENSORY - Facial sensation

TEST:

Sensation coming from skin around the ear

IX: GLOSSOPHARYNGEAL **B**

FUNCTION: **GLOSSO MEANS TONGUE!**

MOTOR - Tongue movement & swallowing
SENSORY - Taste (sour & bitter)

TEST:

Test tongue by giving client sour, bitter, & salty substance.


VIII: VESTIBULOCOCHLEAR / ACOUSTIC **SE**

FUNCTION:

Balance & hearing

TEST:

- Stand with eyes closed
- Otoscopic exam
- Rinne & Weber Tests



VII: FACIAL **B**

FUNCTION:

MOTOR - Facial expression
SENSORY - Taste (sweet & salty)

TEST:

- Ask client to do different facial expression (Frown, smile, raise eyebrows, close eyes, blow etc)
- Test tongue by giving client sour, sweet, bitter, and salty substances.

VI: ABDUCENS **M**

FUNCTION:

Controls parallel eye movement
Abduction - moving laterally
AKA away from midline

TEST:

- Look up, down, & inward
- Ask the client to follow your finger as you move it towards their face

V: TRIGEMINAL **B**

FUNCTION:


MOTOR - Mastication (biting & chewing)
SENSORY - Facial sensation

TEST:

- Pressure on the forehead cheek & jaw with a cotton swab to check sensation
- Ask client to open mouth & then bite down

MNEMONICS

Ooh, Olfactory	Some Sensory
Ooh, Optic	Say Sensory
Ooh, Oculomotor	Marry Motor
To Trochlear	Money Motor
Touch Trigeminal	But Both
And Abducens	My Motor
Feel Facial	Brother Both
Very Vestibulocochlear / Acoustic	Says Sensory
Good Glossopharyngeal	Big Both
Velvet. Vagus	Brains Both
Such Spinal Accessory	Matter Motor
Heaven! Hypoglossal	More Motor



I: OLFACTORY **SE**

FUNCTION:

Sense of smell

TEST:

Smell substance with eyes closed (test each nostril separately)

II: OPTIC **SE**

FUNCTION:

Vision

TEST:

- Snellen chart
- Ophthalmoscopic exam
- Confrontation to check peripheral vision



III: OCULOMOTOR **M**

FUNCTION:

Ocular (eye) motor (movement)
Controls most eye movements, pupil constriction, & upper-eyelid rise

TEST:

- Look up, down, & inward
- Ask the client to follow your finger as you move it towards their face

IV: TROCHLEAR **M**

FUNCTION:

Controls downward & inward eye movement

TEST:

- Look up, down, & inward
- Ask the client to follow your finger as you move it towards their face

CRANIAL NERVES

WHAT ARE CRANIAL NERVES?

Nerves that originate from the brain stem. They send information to & from various parts of the body.

LABEL THE FLAGS:

- SE **SENSORY**
- M **MOTOR**
- B **BOTH**



XII:

FUNCTION:

TEST:

XI:

FUNCTION:

TEST:

X:

FUNCTION:

TEST:

IX:

FUNCTION:

TEST:

VIII:

FUNCTION:

TEST:

VII:

FUNCTION:

TEST:

MNEMONICS

Ooh, O _____	Some S _____
Ooh, O _____	Say S _____
Ooh O _____	Marry M _____
To T _____	Money M _____
Touch T _____	But B _____
And A _____	My M _____
Feel F _____	Brother B _____
Very V _____ / A _____	Says S _____
Good G _____	Big B _____
Velvet. V _____	Brains B _____
Such S _____	Matter M _____
Heaven! H _____	More M _____

I:

FUNCTION:

TEST:


II:

FUNCTION:

TEST:

• _____ chart

E
F P
T O Z
L P E D
P E C F D



III:


FUNCTION:

TEST:

IV:

FUNCTION:

TEST:



VI:

FUNCTION:

TEST:

V:

FUNCTION:

TEST:



WANT MORE WORKSHEETS?

Check out The Complete Laminated Study Templates!

BURNS



WHAT IS A BURN?

Damage to skin integrity

TYPES OF BURNS

THERMAL

MOST COMMON

Superficial heat
Examples: liquid, steam, fire

CHEMICAL

Burn caused by a toxic substance
Can be Alkali or Acidic
Examples: bleach, gasoline, paint thinner

RADIATION

Sunburns (UV radiation)
& cancer treatment (radiation therapy)

INHALATION

Caused by inhaling smoke which can cause flame injury or carbon monoxide poisoning

FRICTION

Burn caused when an object rubs off the skin
Examples: road rash, scrapes, carpet burn

COLD

Skin has been overexposed to cold
Example: frostbite

ELECTRIC

Electrical current that passes through the body causing damage within

POTENTIAL COMPLICATIONS

Dysrhythmias, Fracture of bones.
Release of **myoglobin & hemoglobin** into the blood which can clog the kidneys.

BURNS INJURY DEPTH

1ST DEGREE

SUPERFICIAL

- Epidermis
- Pink & painful (still has nerves)
- No scarring

BLANCHING:
present

HEALS:
a few days

2ND DEGREE

SUPERFICIAL PARTIAL THICKNESS

- Epidermis & dermis
- Blisters, shiny, & moist
- Painful

BLANCHING:
present

HEALS:
2 - 6 weeks

3RD DEGREE

FULL THICKNESS

- Epidermis, dermis, & hypodermis
- May look black, yellow, red & wet
- No pain/limited pain (nerve fibers are destroyed)
- Skin will not heal (need skin grafting)
- **Eschar:** dead tissue, leathery; must be removed!

LAYERS of the skin


↓

EPIDERMIS

DERMIS

HYPODERMIS

subcut/fatty tissue



BURN LOCATION

RESPIRATORY

- Face
- Neck
- Chest
- Torso

DISABILITY

- Hands
- Feet
- Joints
- Eyes

TROUBLE HEALING

- Poor blood supply
- Diabetes
- Infection

INFECTION

Any open area where bacteria can easily enter

- Perineum
- Ears
- Eyes

COMPARTMENT SYNDROME

- In the extremities
Tight skin such as eschar acting like a band around the skin cutting off blood circulation

INHALATION INJURY

Damage to the respiratory system!
Happens mostly in a **closed area**

SIGNS OF INHALATION INJURY

- Hair singed around the face, neck or torso
- Trouble talking
- Soot in the nose or mouth
- Confusion or anxiety

CARBON MONOXIDE (CO) POISONING

Carbon monoxide travels faster than oxygen, making it bind to hgb first.

Now oxygen cannot bind to hgb = **HYPOXIC**

Classic symptom: cherry red skin
Treatment: 100% O₂

NOTE:

Oxygen saturation may appear normal



PHASES OF BURN MANAGEMENT



EMERGENT PHASE

Onset of Injury to the restoration of capillary permeability

24 - 48 HOURS
after burn

PATHO

↑ Capillary permeability (leaky vessels) causing:

- Plasma leaves the intravascular space
 - Albumin & sodium follows
- Fluids shift to the interstitial tissue

Leads to
Edema

Leads to fluids volume deficit (FVD)
in the intravascular space

VITAL SIGNS

- ↑ Pulse
- ↓ Blood pressure
- ↓ Cardiac output
- ↓ Urine output
(from ↓ perfusion to the kidneys)

THINK:
HYPOVOLEMIC
SHOCK!

LABS

- ↑ Potassium (K+)
- ↑ Hematocrit (HCT)
- ↓ White Blood Cells (WBCs)
- ↑ BUN/Creatine

NURSING CONSIDERATIONS

- Establish IV access (preferably 2)
- Fluids (Lactated Ringer's, crystalloids)
- Parkland formula
- Foley catheter to monitor urinary output (UOP)
GOAL: > 30 mL/hr of UOP
- Decrease edema
 - Elevate extremities above heart level

Think
ABCs



ACUTE PHASE

Capillary permeability stabilized - to wound closure

48 - 72 HOURS
after burn & until wounds have healed

PATHO

Capillary permeability is restored which leads to the body diuresing (increased urine production). All the excess fluid that shifted from the interstitial tissue shifts back into the intravascular space.

GOALS

- **PREVENT INFECTION**
 - Systemic antibiotic therapy
- **ENSURE PROPER NUTRITION**
 - Needs ↑ calories
 - Protein & Vit C to promote healing
- **ALLEVIATE PAIN**
- **WOUND CARE**
 - Always premedicate before wound care!
 - Debridement or grafting

NURSING CONSIDERATIONS

- **RENAL**
 - Diuresis is happening
 - Foley catheter to monitor UOP
- **RESPIRATORY**
 - Possible intubation if respiratory complications occurred
- **GASTROINTESTINAL**
 - Since the client is in FVD, there is ↓ perfusion to the stomach
 - Paralytic ileus
 - Curlings ulcer
 - Medication to decrease chance of ulcers
 - H2 histamine blocking agent (↓HCl)
 - Monitor bowel sounds
 - May need NG tube for suctioning



REHABILITATIVE PHASE

Burn healed and the patient is functioning mentally & physically

GOALS

- Psychosocial
- Occupational therapy (OT)
- Activities of daily living (ADLs)
- Cosmetic corrections
- Physical therapy (PT)



FLUID RESUSCITATION FOR BURNS

THE PARKLAND FORMULA

Used to calculate the total volume of fluids (mL) that a patient needs **24 hours** after experiencing a burn
 Apply only in 2nd & 3rd degree burns.


$$4 \text{ ML} \times \text{TBSA (\%)} \times \text{BODY WEIGHT (KG)} = \text{TOTAL ML OF FLUID NEEDED}$$

Give the first 1/2 of the solution in the **FIRST 8 HOURS**

Over the **NEXT 16 HOURS**, give the second 1/2 of the solution

RULE OF NINES

Quick estimate of the % of the **total body surface area (TBSA)** has been effected by a partial & full-thickness burn in an adult client.



PRACTICE QUESTION

PART 1

A 25 year old male patient who weighs **79 kg** has sustained burns to the back of the right arm, posterior trunk, front of the left leg, and their anterior head and neck. Using the **Rule of Nines**, calculate the total body surface area percentage that is burned.

- Back of right arm - 4.5%
- Posterior trunk - 18%
- Front of left leg - 9%
- Anterior head & neck- 4.5%

ANSWER:
36%

PART 2

Use the Parkland formula to calculate the total amount of Lactated Ringer's solution that will be given over the next 24 hours.

ANSWER:
11,376 mL

NOTE:

The formula uses TBSA (%). However, you must calculate using 36. Not 0.36 (also written as 36%).

$$4 \text{ mL} \times 36\% \times 79 \text{ kg} = 11,376 \text{ mL}$$

$$11,376 / 2 = 5,688 \text{ mL}$$

FIRST 8 HOURS

$$11,376 / 2 = 5,688 \text{ mL}$$

NEXT 16 HOURS

Keep in mind: the question could ask you for mL given in the first 24 hours, the first 8 hours, etc., so read the question carefully.

SHOCK

WHAT IS SHOCK?

A life-threatening condition resulting from **INADEQUATE TISSUE PERFUSION**. This leads to possible cell dysfunction, cell death, and even organ failure.

HYPVOLEMIC-SHOCK

MOST COMMON TYPE OF SHOCK

ETIOLOGY

HYPVOLEMIC

"LOW" "VOLUME" "IN THE BLOOD"

Decreased intravascular volume

CAUSES

NON-HEMORRHAGIC (not from bleeding)

- FLUID SHIFT (edema or ascites)
- SEVERE DEHYDRATION (vomiting, diarrhea, burns)

HEMORRHAGIC (from bleeding)

- TRAUMA
- GI BLEED
- POSTPARTUM

SIGNS & SYMPTOMS

Pulse	CO	HR	BP
Weak, thready pulse	↓ Not a lot of blood being pumped by the heart	↑ Tachycardia Compensating to increase blood flow	↓ Hypotension
Skin	CVP	SVR	O ₂ Sat
Cyanosis (Bluish tint of the lips, tongue, and fingertips) Cool, pale skin ↓ capillary refill (>3 seconds)	↓	↑ Vasoconstriction	↓ ↓ blood being perfused to the body = low O ₂

TREATMENT

- Large gauge IVs (at least 2)
- Fluids & blood replacement
 - Crystalloids (example: normal saline or Lactated Ringers)
 - Colloids (albumin)
 - Blood products (plasma, PRBCs, & PLTs)

Other Signs & Symptoms

LABS CAN BE:

- ↑ HCT hemoconcentration
- ↓ HCT actually hemorrhaging the RBCs
- Oliguria (urine output of <30 mL/hr)
- Confused, agitated


due to decreased blood flow to the brain



ETIOLOGY

The heart can't pump enough blood to meet the perfusion needs of the body

NOTE: There is enough blood, the heart just can't pump it to the body which causes **fluid accumulation** in the lungs!



CAUSES


- Damage from an acute MI
- Severe hypoxemia
- Acidosis
- Hypoglycemia
- Cardiomyopathy
- Cardiac tamponade
- Dysrhythmias

SIGNS & SYMPTOMS

Pulse	CO	HR	BP
Weak peripheral pulses	↓ Not a lot of blood being pumped by the heart	↑ Tachycardia Compensating to increase blood flow	↓ Hypotension
Skin	CVP	SVR	O ₂ Sat
Cool, clammy skin ↓ capillary refill (>3 seconds)	↑	↑ Vasoconstriction	↓ ↓ blood being perfused to the body = low O ₂

TREATMENT

- For an MI: Angioplasty
Thrombolytics
- Oxygen
- Vasopressors (example: epinephrine, dobutamine, dopamine)
 - Vasopressors cause vasoconstriction which ↑ blood flow and increases perfusion to the organs
- Diuretics
 - ↓ the workload of the heart
 - ↓ extra blood volume
- Intra-aortic balloon pump (helps to improve coronary artery blood flow & ↑ CO)



Other Signs & Symptoms

- Jugular vein distention (JVD)
 - Chest pain
 - Oliguria (urine output of <30 mL/hr)
 - Confused, agitation
- due to decreased blood flow to the brain
- From fluid accumulation in the lungs:
- Dyspnea
 - Pulmonary edema



BP = Blood pressure HR = Heart rate CO = Cardiac output SVR = Systemic vascular resistance CVP = Central venous pressure

DISTRIBUTIVE SHOCK

(Septic, Neurogenic, Anaphylactic)

DISTRIBUTIVE:

Leaky blood vessels
Excessive **vasodilation**
(widening of vessels)



Intravascular volume
pools in the peripheral
blood vessels



Since the blood is in the
peripherals, it is NOT perfusing
the vital organs which causes
relative hypovolemia

MOST COMMON TYPE OF DISTRIBUTIVE SHOCK

ETIOLOGY

Caused by widespread infection or sepsis 

CAUSES

- Pneumonia
- Wound infection
- Urosepsis
- Invasive procedures
- Bacteria
- Indwelling medical devices (catheters)
- Intra-abdominal infections

SIGNS & SYMPTOMS

Pulse	CO	HR	BP	Other Signs & Symptoms
Bounding pulses	↑	↑ Tachycardia	↓ Hypotension	→ Hyperthermia & fever → Increased respiratory rate → GI upset: Nausea, vomiting, diarrhea, decrease gastric motility → ↑ Inflammatory markers ↑ WBCs ↑ C-reactive protein (CRP)
Skin Initially warm & flushed, but as the BP drops, the skin becomes cool, pale & mottled	↓ CVP	↓ SVR Vasodilation	↑ O ₂ Sat	

TREATMENT

CORRECT THE UNDERLYING CAUSE

- Fluid replacement
- **Broad-spectrum antibiotics**
- Vasopressors (norepinephrine & dopamine)
- Neuromuscular blockade agents & sedation
 - ↓ metabolic demands & provides comfort
- Medications to prevent stress ulcers
 - H2 blocking agents
 - Proton pump inhibitors (PPIs)




Broad spectrum antibiotics are used when the organism is not yet known/determined. Once the organism is known, the client can be put on more specific antibiotics.

SEPTIC SHOCK (SEPSIS)

ETIOLOGY

Vasodilation due to a loss of balance between

In neurogenic shock, the client mainly experiences parasympathetic stimulation which causes VASODILATION for an extended period



PARASYMPATHETIC STIMULATION
(Rest & digest)

→ Causes dilation (relaxing) of the smooth muscles

P think Peaceful

SYMPATHETIC STIMULATION
(Fight or flight)

→ Causes constriction (tightening) of the smooth muscles

CAUSES

- Spinal cord injury (above T6, cervical)
- Spinal anesthesia
- Nervous system damage
- Insulin reaction



NEURogenic = Issue with **NER**vous system

SIGNS & SYMPTOMS


EVERYTHING IS DECREASED

Remember parasympathetic means **RELAXED EVERYTHING**

CO	HR	BP	Other Signs & Symptoms
↓	↓	↓ Hypotension	RELATIVE HYPOVOLEMIA: There is enough blood volume. However, the vascular space is dilated , so blood volume is displaced causing hypovolemia.
Skin Dry, warm extremities (venous blood pooling) Hypothermia: warm/dry extremities, cold body	↓ CVP	↓ SVR Vasodilation	

the sympathetic NS is not working to compensate & ↑ the HR

TREATMENT DEPENDS ON THE CAUSE OF THE SHOCK

- Spinal cord injury → **PROTECT THE SPINE:** Keep spine immobilized (cervical collar, backboards, log-rolling)
- Assess & manage airway
May need intubation or mechanical ventilation
- Elevate the head of the bed
- IV fluids  Watch for fluid volume overload
- Increased risk for clots due to pooling of blood
 - Watch for signs of a clot
 - Compression devices
 - Antithrombotic agents (heparin)
- Vasopressors (example: epinephrine, dobutamine, dopamine)

S&S OF BLOOD CLOTS:

- Pain in the extremities
- Redness
- Tenderness
- Warmth




BP = Blood pressure HR = Heart rate CO = Cardiac output SVR = Systemic vascular resistance CVP = Central venous pressure

DISTRIBUTIVE SHOCK





(Septic, Neurogenic, Anaphylactic)



ETIOLOGY




CAUSES/TRIGGERS

- Often unknown (idiopathic)
- Foods (example: peanuts) 
 - Medications 
 - Insects (example: bee sting) 
 - Latex 
 - Exercise-induced anaphylaxis (EIA)

Signs & symptoms usually occur within 2 - 30 minutes of exposure to antigen

TREATMENT

- High-flow oxygen
- First-line drug: **EPINEPHRINE** ★ 
- Causes vasoconstriction & bronchodilation
- Other possible medications
 - Antihistamines
 - Diphenhydramine (Benadryl)
 - Albuterol (Proventil)
 - Corticosteroids
- Fluids
- Stay with the client & monitor

BIPHASIC ANAPHYLAXIS:
A recurrence of anaphylaxis after appropriate treatment


SIGNS & SYMPTOMS

Pulse	CO	HR	BP
Rapid, weak pulse	↓	↑ Tachycardia	↓ Hypotension

CAPILLARY PERMEABILITY:
Fluid is leaving the intravascular space

Skin	CVP	SVR	O ₂ Sat
Generalized flushing	↓	↓ Vasodilation	↓

Other Signs & Symptoms

- * **CARDIAC**
 - Cardiac dysrhythmias or cardiac arrest
- * **GI**
 - Nausea/vomiting
 - Acute abdominal pain
- * **FEELING OF IMPENDING DOOM**
- * **RESPIRATORY**
 - Bronchoconstriction
 - Difficulty breathing
 - Wheezing
 - Coughing
 - Unable to speak
- * **SKIN**
 - Itching, generalized flushing, redness, hives, or a rash may be present 

HOW TO USE AN EPINEPHRINE AUTO-INJECTOR (EAI)


EDUCATION POINTS:

- Store in dark room
- Administer EAI immediately after the first sign of an allergic reaction

EXPECTED SYMPTOMS AFTER ADMINISTRATION:

- Tachycardia
- Palpitations
- Dizziness

INJECT IN THE OUTER THIGH AT A 90° ANGLE




BP = Blood pressure HR = Heart rate CO = Cardiac output SVR = Systemic vascular resistance CVP = Central venous pressure

ANAPHYLACTIC SHOCK

ABGs

4 MUST-KNOW COMPONENTS



ABGs MEASURE HOW ACIDIC OR ALKALOTIC THE BLOOD IS IN THE ARTERIAL CIRCULATION.

*also a measure of gases such as O₂ & CO₂

PH	Measurement of how acidic or alkalotic your blood is	regulated by both lungs & kidneys	7.35 - 7.45
PACO₂	Measurement of carbon dioxide in the blood CO₂ think aCid	Regulated by the lungs	35 - 45
HCO₃	Measurement of bicarbonate in the blood Bicarbonate think Base	Regulated by the kidneys	22 - 26
PAO₂	Measurement of oxygen in the blood	Regulated by the lungs	80 - 100

Value not needed to interpret alkalosis or acidosis. It just tells you if the patient is hypoxic or not.

ABG INTERPRETATION

1 KNOW YOUR LAB VALUES!

	ACIDOSIS	NORMAL	ALKALOSIS
PH	< 7.35 ↓	7.35 - 7.45	> 7.45 ↑
CO₂	> 45 ↑	35 - 45	< 35 ↓
HCO₃	< 22 ↓	22 - 26	> 26 ↑

2 RESPIRATORY OR A METABOLIC PROBLEM?

there are 2 ways to analyze the information

ROME METHOD

Respiratory	PH ↑	CO ₂ ↓	Alkalosis
Opposite	PH ↓	CO ₂ ↑	Acidosis
Metabolic	PH ↑	HCO ₃ ↑	Alkalosis
Equal	PH ↓	HCO ₃ ↓	Acidosis

TIC-TAC-TOE METHOD

ACID	NORMAL	BASE

3 UNCOMPENSATED, PARTIALLY COMPENSATED, OR FULLY COMPENSATED?

If the **PH** is out of range & **CO₂** or **HCO₃** is in range =


UNCOMPENSATED

If **CO₂**, **HCO₃** & **PH** are ALL out of range =

PARTIALLY COMPENSATED

If **PH** is in range (7.35 - 7.45) =

FULLY COMPENSATED



PH IN RANGE? Just because the PH is "normal", it can still fall on a acidotic side or alkalotic side

How do the organs compensate?

KIDNEYS



B think BASE

BICARB HYDROGEN



hours - days to compensate

Excreting excess **ACID & BICARB (HCO₃)** OR Retaining **HYDROGEN & BICARB (HCO₃)**

LUNGS



CO₂ think ACID

CO₂



compensates FAST!

HYPERventilation = **ALKALOSIS**
↓ CO₂ = **ALKALOSIS**
HYPOventilation = **ACIDOSIS**
↑ CO₂ = **ACIDOSIS**

ABG PRACTICE QUESTION EXAMPLE

QUESTION

A client with a bowel obstruction has been treated with gastric suctioning for 4 days. The nurse notices an increase in nasogastric drainage. Which Acid-base imbalance does that nurse correctly identify?

The patient labs are the following →

Ph **7.50**
PaCO₂ **50** mm Hg
PaO₂ **90** mm Hg
HCO₃ **32** mEq/L

Value not needed to interpret alkalosis or acidosis. It just tells you if the patient is hypoxic or not.

TIC-TAC-TOE METHOD

1

What does the problem give you?

PH	7.50	<input type="radio"/> ACIDIC	<input checked="" type="radio"/> ALKALOTIC	<input type="radio"/> NORMAL
CO ₂	50	<input checked="" type="radio"/> ACIDIC	<input type="radio"/> ALKALOTIC	<input type="radio"/> NORMAL
HCO ₃	32	<input type="radio"/> ACIDIC	<input checked="" type="radio"/> ALKALOTIC	<input type="radio"/> NORMAL

2

ACID	NORMAL	BASE
CO ₂		PH
		HCO ₃

- RESPIRATORY ACIDOSIS
- RESPIRATORY ALKALOSIS
- METABOLIC ACIDOSIS
- METABOLIC ALKALOSIS

3

UNCOMPENSATED, PARTIALLY COMPENSATED, or FULLY COMPENSATED?

- Is the pH in range? YES NO
- Is the CO₂ in range? YES NO
- Is the HCO₃ in range? YES NO
- UNCOMPENSATED
- PARTIALLY COMPENSATED
- FULLY COMPENSATED

If CO₂, HCO₃ & PH are ALL out of range

FINAL ANSWER:

METABOLIC ALKALOSIS, PARTIALLY COMPENSATED

ROME METHOD

1

What does the problem give you?

PH	7.50	<input type="radio"/> ACIDIC	<input checked="" type="radio"/> ALKALOTIC	<input type="radio"/> NORMAL
CO ₂	50	<input checked="" type="radio"/> ACIDIC	<input type="radio"/> ALKALOTIC	<input type="radio"/> NORMAL
HCO ₃	32	<input type="radio"/> ACIDIC	<input checked="" type="radio"/> ALKALOTIC	<input type="radio"/> NORMAL

2

Which of the four scenarios from the ROME method matches the information given in your problem?

Respiratory	PH ↑	CO ₂ ↓	Alkalosis
Opposite	PH ↓	CO ₂ ↑	Acidosis
Metabolic	PH ↑	HCO₃ ↑	Alkalosis
Equal	PH ↓	HCO ₃ ↓	Acidosis

- RESPIRATORY ACIDOSIS
- RESPIRATORY ALKALOSIS
- METABOLIC ACIDOSIS
- METABOLIC ALKALOSIS

3

UNCOMPENSATED, PARTIALLY COMPENSATED, or FULLY COMPENSATED?

- Is the pH in range? YES NO
- Is the CO₂ in range? YES NO
- Is the HCO₃ in range? YES NO
- UNCOMPENSATED
- PARTIALLY COMPENSATED
- FULLY COMPENSATED

If CO₂, HCO₃ & PH are ALL out of range

FINAL ANSWER:

METABOLIC ALKALOSIS, PARTIALLY COMPENSATED

RESPIRATORY ACIDOSIS VS. RESPIRATORY ALKALOSIS

RESPIRATORY ACIDOSIS

PATHOPHYSIOLOGY



LUNG PROBLEM

The lungs are **RETAINING** too much **CO₂**



KIDNEYS COMPENSATE

The kidneys excrete excess **HYDROGEN** & retain **BICARB (HCO₃)**

PH < 7.35
CO₂ > 45

CAUSES

RETAINING CO₂: "Depress" breathing

- D**rugs (opioids & sedatives)
- E**dema (fluid in the lungs)
- P**neumonia (excess mucus in the lungs)
- R**espiratory center of the brain is damaged
- E**mboli (pulmonary emboli)
- S**pasms of the bronchial (asthma)
- S**ac elasticity damage (COPD & emphysema)

All these things cause impaired gas exchange

SIGNS & SYMPTOMS

- ↓ Blood pressure
- ↓ Respiration rate
- ↑ Heart rate
- Restlessness
- Confusion
- Headache
- Sleepy / coma

INTERVENTIONS

- Administer O₂
- Semi-Fowler's position
- Turn, cough, & deep-breathe (TCDB)
- Pneumonia: ↑ fluids to thin secretions & administer antibiotics
- If CO₂ > 50, they may need an endotracheal tube
- Monitor potassium levels

NORMAL K+
3.5 - 5.0 mmol/L

RESPIRATORY ALKALOSIS

PATHOPHYSIOLOGY



LUNG PROBLEM

The lungs are **LOSING** too much **CO₂**



KIDNEYS COMPENSATE

The kidneys excrete excess **BICARB (HCO₃)** & retain **HYDROGEN**

PH > 7.45
CO₂ < 35

CAUSES

LOSING CO₂: "Tachypnea"

- ↑ Temperature
- Aspirin toxicity
- Hyperventilation

SIGNS & SYMPTOMS

- ↑ Respiratory rate > 20 breaths/min
- ↑ Heart rate
- Confused & tired
- Tetany
- EKG changes
- (+) Chvostek's sign

Twitching of the facial muscles when tapping the facial nerve in response to **HYPOCALCEMIA**

INTERVENTIONS

- Provide emotional support
- Fix the breathing problem!
- Encourage good breathing patterns
- Rebreathing into a paper bag
- Give anti-anxiety medications or sedatives to ↓ breathing rate
- Monitor K+ & Ca- levels

NORMAL CA-
9 - 11 mg/dL

METABOLIC ACIDOSIS VS. METABOLIC ALKALOSIS

METABOLIC ACIDOSIS


METABOLIC ALKALOSIS

PATHOPHYSIOLOGY



KIDNEY PROBLEM

Too much **HYDROGEN**
Too little **BICARB** (HCO_3)



LUNGS COMPENSATE

The lungs will blow off **CO_2**


PH < 7.35 **HCO_3** < 22

PATHOPHYSIOLOGY



KIDNEY PROBLEM

Too much **BICARB** (HCO_3)
Too little **HYDROGEN**



LUNGS COMPENSATE

The lungs will retain **CO_2**

PH > 7.45 **HCO_3** > 26

CAUSES

- Diabetic ketoacidosis → Not enough insulin = ↑ fat metabolism = excess **ketones (acid)**
- Acute/chronic kidney injury
- Malnutrition → Breaking down of fats = excess **ketones (acid)**
- Severe diarrhea → Remember **Bicarb** comes out of your **Base**

CAUSES

- Too many antacids → Too much **sodium bicarbonate (BASE)**
- Diuretics
- Excess vomiting → Excess loss of **hydrochloric acid (HCL)** from the stomach
- Hyperaldosteronism

SIGNS & SYMPTOMS

- ↑ Respiratory rate → **KUSSMAUL'S BREATHING**
Deep rapid breathing > 20 breaths per minute
- Hyperkalemia
 - Muscle twitching
 - Weakness
 - Arrhythmias
- ↓ Blood pressure
- Confusion

SIGNS & SYMPTOMS

- ↓ Respiratory rate → **HYPOVENTILATION**
 < 12 breaths per minute
- ↓ Potassium (K^+)
 - Dysrhythmias
 - Muscle cramps/weakness
 - Vomiting
- Tetany
- Tremors
- EKG changes

Metabolic **ACIDOSIS** = ↑ serum potassium
Metabolic **ALKALOSIS** = ↓ serum potassium

INTERVENTIONS

- Monitor intake & output
- Administer IV solution of sodium bicarb to ↑ bases & ↓ acids
- Initiate seizure precaution
- Monitor K^+ levels

NORMAL K^+
3.5 - 5.0 mmol/L

DIABETIC KETOACIDOSIS (DKA)

- Give insulin (this stops the breakdown of fats which stops **ketones** from being produced)
- Monitor for hypovolemia due to polyuria

KIDNEY DISEASE

- Dialysis to remove toxins
- Diet
 - ↑ Calories
 - ↓ Protein

INTERVENTIONS

- Monitor K^+ and Ca^{2+} levels
- Administer IV fluids to help the kidneys get rid of bicarbonate
- Replace K^+
- Give antiemetics for vomiting (Zofran or Phenergan)
- Watch for signs of respiratory distress

NORMAL Ca^{2+}
9 - 11 mg/dL

FRACTURES


WHAT IS A FRACTURE? A fracture is a complete or incomplete disturbance in the progression of bone structure

TYPES OF FRACTURES



COMMINUTED


The bone is crushed causing lots of little fragments



TRANSVERSE

The bone is fractured straight across

commonly seen in children




OBLIQUE

The fracture runs at an angle across the bone




GREENSTICK

One side of the bone is bent, the other is broken



IMPACTED

The fractured bone is driven into another bone



SPIRAL

The break partially encircles the bone



OPEN/COMPOUND

A fracture where the bone breaks through the skin

increased risk for INFECTION

STAGES OF BONE HEALING



STAGE I

HEMATOMA FORMATION

- First 1-2 days of fracture
- Bleeding into the injured site occurs

MNEMONIC
"HE FELL BECAUSE HE WAS RUNNING"



STAGE II

FIBROCARTILAGINOUS CALLUS FORMATION

- Formation of granulation tissue
- Reconstruction of bone begins
- Still not strong enough to bear weight



STAGE III

BONY CALLUS FORMATION (OSSIFICATION)

- 3rd - 4th week of fracture healing
- Mature bone is replacing the callus



STAGE IV

REMODELING

- This may take months to years!
- Compact bone replaces spongy bone
- X-rays are used to monitor the progress of bone healing

NURSING ASSESSMENT

POST-FRACTURE *Neuromuscular assessments*

- 5 P'S**
- Pain**
- Pallor**
- Pulselessness**
- Paresthesia**
- Paralysis**

burning or tingling sensation

COMPARTMENT SYNDROME

Increased pressure and build-up, causes tissue impairment leading to cell death!

Pressure ↑



Blood flow cut off



Tissue damage due to **HYPOXIA** (lack of oxygen)

SIGNS & SYMPTOMS

- Deep, throbbing, unrelenting pain
- Pain unrelieved by medications
- Disproportional to the injury
- Intensifies with passive ROM

TREATMENT

IMMEDIATE

- Place extremity at the heart level (not above heart level)
- Open the cast or splint

FASCIOTOMY

Fascia is cut to relieve tension & pressure

GOUT

PATHOLOGY

Gout is a form of arthritis characterized by increased uric acid levels.

HYPERURICEMIA

"high" "uric acid" "in the blood"

This causes deposits of uric acid crystals in the joints.


TOPHI

Accumulation of sodium urate crystals in joints such as the big toe and hands, or other areas such as the ears.



MEMORY TRICK

Tophi think Toe



WHAT IS URIC ACID?

Uric acid is created from purine breakdown during digestion. It's produced by the liver and is mostly excreted by the kidneys.

EXPECTED RANGE:

F: 2.5 - 8 mg/dL

M: 1.9 - 7.5 mg/dL

CAUSES

- Diet high in purines
- Certain medications
 - Diuretics (causes dehydration)
 - Aspirin
 - Cyclosporine
- Disorder of purine metabolism
- Kidney problems
 - Inadequate excretion of uric acid by the kidneys

SIGNS & SYMPTOMS

Can be **ACUTE** or **CHRONIC**

- Acute gouty arthritis
- Pain (severe)
- Swelling
- Warmth at the site
- Bone deformity
- Joint damage
- Tophi
- Renal calculi

EDUCATION

Educate on avoiding:

- Foods high in purines
- Medications (aspirin)
- Alcohol
- Dehydration


foods high in purines:



Stay hydrated: 2-3 liters per day



- Uric acid deposits can cause kidney stones, fluids help prevent this!
- Weight loss program if overweight



MEDICATIONS

GENERIC	TRADE NAME
allopurinol	Aloprim, Zyloprim, Lopurin



MEMORY TRICK

AlloPurinol → Prevents gout

GENERIC	TRADE NAME
colchicine	Mitigare, Colcrys



MEMORY TRICK

Colchicine → for aCute gout attacks

*For more information about gout medications, see the musculoskeletal section in the Pharmacology Bundle


OSTEOPOROSIS

PATHOLOGY

OSTEOPOROSIS

"relating to bone" "porous"

Osteoporosis essentially means: **HAVING POROUS BONES**



Normal bone marrow has small holes in it, but osteoporosis causes much larger holes

The rate of **BONE RESORPTION** (osteoclasts) is greater than the rate of **BONE FORMATION** (osteoblasts) = **↓ DECREASED TOTAL BONE MASS**

DIAGNOSTIC

Bone density test: Dual-energy x-ray absorptiometry (**DEXA**)

This process takes X-ray images measuring calcium and other minerals in the bones


RISK FACTORS

- C** Calcium & vitamin intake is **LOW**
- A** Age: **women after menopause** (the decrease in estrogen at menopause causes increase bone resorption)
- L** Lifestyle (smoking, excessive alcohol intake, sedentary lifestyle, immobility)
- C** Caucasian or Asian women
- I** Inherited (family history)
- U** Underweight/malabsorption disorder (Celiac disease, bariatric surgery, eating disorders)
- M** Medications: **long-term use of corticosteroids**, anticonvulsants, levothyroxine, long-term use of proton pump inhibitors, etc.



SIGNS & SYMPTOMS

- May be asymptomatic until a fracture occurs
- FRACTURES** (hips, spine, wrist)
- Low back, neck, or hip pain
- The back will be rounded (hunch back) causing height loss



FRACTURES

Clients often think they fell and broke something, **BUT** bones may break first causing them to fall.

NURSING INTERVENTIONS

ASSESSING FOR RISK FACTORS
Educate on stopping smoking & limiting alcohol

EDUCATE ON WAYS TO PREVENT OSTEOPOROSIS

TEACHING ABOUT PREVENTING INJURY

AT HOSPITAL

- Use call light
- Non-slip socks
- Communicate falls risk
- Clutter-free environment

PREVENTION

- Weight-bearing exercises (weights, hiking, etc).
- Consume foods rich in calcium & vitamin D



AT HOME

- No area rugs (risk for falling)
- Watch out for pets
- Keep glasses near by



MEDICATIONS

- Calcium supplements with Vitamin D
- Bisphosphonates (ends in "**DRONATE**")




*For more information about bisphosphonates, see the Pharmacology Bundle

OSTEOARTHRITIS (OA) & RHEUMATOID ARTHRITIS (RA)

OSTEOARTHRITIS (OA)

PATHOLOGY

- OA is a noninflammatory degenerative disorder of the joints. It's caused by the breakdown of cartilage between the joints.
- The articular cartilage breaks down, which leads to damage to the bone.




RISK FACTORS

- OBESITY
- Older age
- Female gender
- Certain occupations (heavy labor)
- Genetics


TREATMENT

- Orthotic devices (splints, braces, knee braces)
- Walking aids (canes)
- Exercise
- Weight loss
- Occupational therapy (OT) & physical therapy (PT)
- Analgesics



DISTAL
Distal interphalangeal (DIP) called **HEBERDEN'S NODES**

PROXIMAL
Proximal interphalangeal (PIP) called **BOUCHARD'S NODES**



SIGNS & SYMPTOMS

- Pain
 - Stiffness (morning stiffness)
 - Functional impairment
 - Bony enlargements
- Occurring mostly at the weight-bearing joints (hips, knees)


MOVEMENT / EXERCISE → Aggravated / symptoms worsen
REST → Symptoms are relieved

RHEUMATOID ARTHRITIS (RA)

PATHOLOGY

Exact mechanism is unknown

- RA is a chronic, inflammatory type of arthritis. It's classified as an autoimmune disease.



SIGNS & SYMPTOMS

- Symmetric joint pain
- Symptoms are typically **BILATERAL** & symmetric
- Stiffness in the morning (lasting >1 hour)
- Swelling, warmth, and redness
- Deformity of the fingers
- Can effect all joints (fingers, wrists, neck, shoulders, etc).
- Systemic effects: heart, lungs, skin, etc.



STAGES OF RHEUMATOID ARTHRITIS

1 SYNOVITIS

- Inflammation of the synovium
- Synovial membrane thickens

2 PANNUS FORMATION

- Pannus is a layer of vascular fibrous tissue

3 FIBROUS ANKYLOSIS

- Joint invaded by fibrous connective tissue

4 BONY ANKYLOSIS

- When the bones are fused together

This causes loss of...

- Articular surfaces
- Joint motion
- Ligament elasticity

DIAGNOSIS

- Hard to diagnose because symptoms are very similar to other diseases
- (+) Rheumatoid factor
- Increase erythrocyte sedimentation
- C-reactive protein (indicates inflammation in the body)
- X-ray shows joint deterioration


RISK FACTORS

- May cause an inflammatory response & destructive synovial fluid
- Environmental factors (smoking, pollution)
 - Bacterial or viral illness
 - Cigarette smoking
 - Family history


TREATMENT

GOAL: Decrease joint pain & swelling. Decrease changes of joint deformity & minimize disability.

NO CURE



- Medications
- Surgery
 - SYNOVECTOMY:** removal of synovium
 - JOINT REPLACEMENT**
 - ARTHRODESIS:** "joint fusion"
- Joint support
 - Splints & assistive devices
- Range of motion (ROM) exercise
- Low impact exercise (walking, water aerobics, etc.)
- Occupational therapy (OT) & physical therapy (PT)
- Heat or cold? **HEAT** → For stiffness
COLD → For pain/inflammation



PHARMACOLOGY

BROUGHT TO YOU BY



ANTIBIOTICS / ANTIBACTERIALS

Broad spectrum antibiotics.....	-OXACIN
Tetracyclines	-CYCLINE
Sulfonamides	SULF-
Cephalosporins	-CEF CEPH-
Penicillins	-CILLIN
Aminoglycosides & macrolides.....	-MYCIN
Fluoroquinolones	-FLOXACIN

ANTIVIRALS

Antiviral (disrupts viral maturation)	-VIRIMAT
Antiviral (undefined group)	VIR- -VIR- -VIR
Antiviral (neuraminidase inhibitors)	-AMIVIR
Antiviral (acyclovir)	-CYCLOVIR
HIV protease inhibitors.....	-NAVIR
HIV / AIDS	-VUDINE

ANTIFUNGAL

Antifungal.....	-AZOLE
-----------------	---------------

ANESTHETICS / ANTIANXIETY

Local anesthetics	-CAINE
Barbiturates (CNS depressant)	-BARBITAL
Benzodiazepines (for anxiety/sedation)	-ZOLAM
Benzodiazepines (for anxiety/sedation)	-ZEPAM

ANTIDIABETIC

Oral hypoglycemics.....	-IDE -TIDE -LINIDE
Inhibitor of the DPP-4 enzyme.....	-GLIPTIN
Thiazolidinedione	-GLITAZONE

CARDIAC

ANTIHYPERTENSIVES

ACE inhibitors	-PRIL
Beta-blockers	-OLOL
Angiotensin II receptor antagonists	-SARTAN
Calcium channel blockers	-PINE -AMIL
Vasopressin receptor antagonists	-VAPTAN
Alpha-1 blockers	-OSIN
Loop diuretics	-IDE -SEMIDE
Thiazide diuretics	-THIAZIDE
Potassium sparing diuretics	-ACTONE

ANTHYPERLIPIDEMICS

HMG-CoA reductase inhibitor	-STATIN
-----------------------------------	---------

OTHER

Anticoagulant (Factor Xa inhibit)	-XABAN
Anticoagulants (Dicumarol type)	-AROL
Anticoagulants (Hirudin type)	-IRUDIN
Low-molecular-weight heparin (LMWH)	-PARIN
Thrombolytics (clot-buster)	-TEPLASE -ASE
Antiarrhythmics	-ARONE

ANALGESICS / OPIOIDS

Opioids	-DONE
Opioids	-ONE
NSAID's (anti-inflammatory)	-OLAC -PROFEN
Salicylates	ASPRIN (ASA)
Nonsalicylates	ACETAMINOPHEN

GASTROINTESTINAL

Histamine H2 antagonists (H2-blockers)	-TIDINE -DINE
Proton pump inhibitor (PPIs)	-PRAZOLE
Laxatives	-LAX

ANTIDEPRESSANTS

Selective serotonin reuptake inhibitors (SSRIs)	-OXETINE -TALOPRAM -ZODONE
Serotonin-norepinephrine reuptake inhibitors (SNRI/DNRI)	-FAXINE -ZODONE -NACIPRAM
Tricyclic antidepressants (TCAs)	-TRIPTYLINE -PRAMINE

MISCELLANEOUS

Corticosteroids	-ASONE -OLONE -INIDE
Triptans (anti-migraine)	-TRIPTAN
Ergotamines (anti-migraine)	-ERGOT-
Antiseptics	-CHLORO
Antituberculars (TB)	RIFA-
Bisphosphonates	-DRONATE
Neuromuscular blockers	-NUIM
Retinoids (anti-acne)	TRETIN-
Phosphodiesterase 5 inhibitors	-AFIL
Carbonic anhydrase inhibitors	-LAMIDE
Progestin (female hormone)	-TREL
Atypical antipsychotics	-RIDONE

RESPIRATORY











UPPER RESPIRATORY

Second-gen antihistamines (H1 antagonist)	-ADINE
Second-gen antihistamines (H1 antagonist)	-TIRIZINE
Second-gen antihistamines (H1 antagonist)	-TICINE
Nasal decongestants	-EPHRINE -ZOLINE


LOWER RESPIRATORY

Beta2-agonists (Bronchodilator)	-TEROL
Xanthine derivatives	-PHYLLINE
Cholinergic blockers	-TROPIMUM
Cholinergic blockers	-CLINDIDIUM
Immunomodulators & leukotriene modifiers	-ZUMAB -LUKAST

ANTIDOTES

Opioids / Narcotics	NALOXONE (NARCAN)  NO more Opioids NARCAN → OPIOIDS
Warfarin (Coumadin)	VITAMIN K  During WAR , Vitamin K Kills WAR farin
Heparin	PROTAMINE SULFATE  You will need HELP from a PRO to stop bleeding out
Digoxin	DIGIBIND OR DIGIFAB
Anticholinergic toxicity	PHYSOSTIGMINE
Benzodiazepines	FLUMAZENIL (ROMAZICON)  I FLU fast in my mercedes BENZ
Cholinergic toxicity	ATROPINE (ATROPEN)  We don't have time to CHAT , we have a toxic situation CHOLINERGIC → ATROPINE
Acetaminophen (Tylenol)	ACETYL CYSTEINE (MUCOMYST)  ACETAMINOPHEN → ACETYL CYSTEINE
Magnesium sulfate	CALCIUM GLUCONATE  MAG gie CAL Ls for help! MAGNESIUM → CALCIUM
Iron toxicity	DEFEROXAMINE  DEFEROXAMINE → FER rous means "containing iron"
Lead toxicity	SUCCIMER OR CALCIUM DISODIUM EDETATE 
Alcohol withdrawal	CHLORDIAZEPOXIDE (LIBRIUM)
Beta blockers	GLUCAGON  Beta blockers be GON e with Gluca GON
Calcium channel blockers	GLUCAGON, INSULIN, OR CALCIUM
Aspirin	SODIUM BICARBONATE  You take ASPIRIN when you have a headache. You may also want a SALTY snack when you have a headache.
Insulin reaction	GLUCAGON  If you want your insulin GON e, you give Gluca GON
Pyridoxine	DEFEROXAMINE
Tricyclic antidepressants	SODIUM BICARBONATE
Cyanide poisoning	HYDROXOCOBALAMIN

These are chelation agents

 Look at these memory tricks to help with remembering these antidotes!

COMMON THERAPEUTICS LEVELS

Digoxin	0.5 - 2.0 ng/mL (> 2 = TOXIC)
Lithium	0.6 - 1.2 mEq/L
Theophylline	10 - 20 mcg/dL
Dilantin (Phenytoin)	10 - 20 mg/L
Magnesium sulfate	4 - 7 mg/dL
Acetaminophen (Tylenol)	10 - 20 mcg/mL
Gentamicin	5 - 10 mcg/mL
Salicylate	100 - 300 mcg/mL
Vancomycin	Peak: 20 - 40 mcg/mL Trough: 5 - 15 mcg/mL
Valporic acid	50 - 100 mcg

SALICYLATES & NONSALICYLATES

SALICYLATES

GENERIC TRADE NAME

aspirin -

ACTION

- 👉 Analgesic & antipyretic
- 👉 Anti-inflammatory
- 👉 Anticoagulant

- Analgesic
 - Inhibits prostaglandins. Prostaglandins make pain receptors more sensitive to feel pain.
- Antipyretic
 - ↓ body temp by dilating the blood vessel & spreading the blood throughout the body.
- Aspirin
 - Prolongs bleeding times.
 - Inhibits the clumping of platelets.

USES

- Mild to moderate pain
- ↓ body temp
- Inflammatory conditions (RA, OA, & rheumatic fever)
- Aspirin is used to ↓ the risk of an MI & CVA

SIDE EFFECTS

- GI upset
 - Heartburn
 - Anorexia
 - Nausea / vomiting
 - GI bleeding



CONTRAINDICATIONS

- Known sensitivity to Salicylates or NSAIDS
- Any **BLEEDING TENDENCIES**
 - GI bleeding (peptic ulcers)
 - Blood dyscrasia
 - Bleeding disorders
 - On anticoagulants
 - Vit K deficiency
- Children with recent viral infection
 - Risk for **REYE'S SYNDROME!**



NURSING CONSIDERATIONS

- Stop taking salicylates 1-week prior to major surgery (remember ↑ risk for bleeding)
- Monitor for GI bleeding

ANTIDOTE: activated charcoal

TOXICITY?

- 1) Gastric lavage
- 2) Activated charcoal (within 2 hours of ingestion)

NONSALICYLATES

GENERIC TRADE NAME

acetaminophen Tylenol

ACTION

- 👉 Analgesic & antipyretic

Action is not completely known. Does NOT have any anti-inflammatory or antiplatelet effects.

USES

- Mild to moderate pain
- Aspirin substitute for those with:
 - Allergy to aspirin
 - bleeding tendencies
- Children with fever / flu-like symptoms

SIDE EFFECTS

- Hives
- Hemolytic anemia
- Pancytopenia
- Hypoglycemia
- Liver damage
 - Hepatotoxicity
 - Hepatic failure
 - Jaundice

These side effects rarely occur when the medication is taken as directed. They occur due to **CHRONIC USE** or **HIGHER DOSAGE THAN RECOMMENDED**



CONTRAINDICATIONS

- Known sensitivity to acetaminophen
- Those with liver dysfunction
 - Chronic alcohol use



NURSING CONSIDERATIONS

- Before adm. of acetaminophen, assess overall health & alcohol use
- Malnourished clients & those with chronic alcohol use (>3 drinks /day) are at increased risk for liver damage
- Limit dosage to 1000-2000 mg/day!



ANTIDOTE: Acetylcysteine (mucomyst)

↪ This protects the liver cells & destroys acetaminophen metabolism

TOXICITY?

- 1) Gastric lavage (within 4 hours of ingestion)
- 2) Give antidote via nebulizer within 24 hours of ingestion

NSAIDS

NSAIDS Non-steroidal Anti-Inflammatory Drugs


Gets their name because they produce an anti-inflammatory effect but they are not steroids!

ACTION



- Anti-inflammatory
- Analgesic
- Antipyretic

Inhibit prostaglandin synthesis by blocking cyclooxygenase (COX)



This means they inhibit pain, but also inhibit the enzyme that maintains the lining of the stomach!

GENERIC	TRADE NAME
ibu profen	Advil
fenop profen	Nalfon
flurbi profen	-
diclofenac	-
celecoxib	Celebrex
ketor olac	Sprix (nasal spray)
naproxen	Aleve
indomethacin	Indocin

SUFFIXES: -PROFEN, -OLAC

Inhibits COX 2 without inhibiting COX 1

USES

- Mild to moderate pain
- Menstrual cramps
- ↓ fever
- Musculoskeletal disorders
 - OA & RA

SIDE EFFECTS

- GI upset **MOST COMMON**
 - Nausea / diarrhea / vomiting
 - Anorexia
 - Abdominal pain / discomfort
- Heart
 - HTN & heart failure
- Kidney clogging
 - NSAIDs are nephrotoxic!
- Blood clots
 - Stroke

CONTRAINDICATIONS

- Known hypersensitivity to NSAIDs or aspirin
- Clients with clot history
 - MI, CVA, PE, DVT
- Clients with liver, kidney, or bleeding disorders




Certain medications are known to cause bronchospasms in clients with asthma.

We want to "BAN" these medications from asthma patients.

NURSING CONSIDERATIONS

- NSAIDs cause GI upset such as acid reflex
 - Administer proton pump inhibitors (PPIs)
 - ➔ Omeprazole
 - ➔ Pantoprazole
- Educate: take with food to decrease stomach upset
 - Don't take on an empty stomach



OPIOID ANALGESICS

GENERIC
hydromorphone
codeine
oxycodone
fentanyl
morphine sulfate
SUFFIXES: -DONE, -ONE

ACTION

CNS Depressant

Binds to opioid receptors in the brain which causes an analgesic sedative, & euphoric effect.



- Most commonly used opioid for chronic pain.
- Can be given in many forms: (PO, nasally, subcut, IM, IV, & suppository)

USES

- ↓ anxiety & sedate post-op
- ↓ anxiety in those with dyspnea
- Relieve pain (myocardial infarction)
- Manage opioid dependence
- Treat diarrhea & intestinal cramping

Opioids do **NOT** produce an anti-inflammatory effect or an antipyretic effect. So they are not used to reduce fevers or for gout / rheumatoid arthritis.

SIDE EFFECTS



- ↓ GI Function
- Constipation

LONG TERM SIDE EFFECTS

Client will **NOT** build tolerance

- ↓ Vital signs
 - ↓ HR
 - ↓ BP (hypotension)
 - ↓ RR
- ↓ CNS function
 - Sedation, insomnia, weakness, dizziness
- Pruritus (itching)
- Nausea
- IV admin
- Burning sensation

SHORT TERM SIDE EFFECTS


Client **WILL** build tolerance

TOLERANCE VS. DEPENDENCE

The body adapts to the drug (gets used to it)

Higher doses of medication are needed to achieve the same effect!

The body goes through "withdrawals" & experiences negative effects when the medication is STOPPED!



ANTIDOTE: Naloxone (Narcan)

- Reversal agent for opioid overdose
- Opioids last longer than the effect of naloxone (Narcan)
- Repeat doses may be needed



This reverses the opioid's effects and the client's pain will come back!

NURSING CONSIDERATIONS

- Preventative measures for **CONSTIPATION**
 - Adm. stool softeners or laxatives
 - Daily exercise
 - **F**luids, **F**iber, & **F**ruits → **F**ill up the toilet!
 - Encourage client to defecate when they feel the urge (do not wait)
- Client respirations begin to drop
 - Coaching the client to breath may increase the respiratory rate
 - Administer naloxone (Narcan)
- Preventative measures for falls
 - Opioids causes orthostatic hypotension
 - Educate to rise slowly, assist the client with ambulatory activities
 - Keep the room well lit
- Take PO opioids with food to decrease GI upset
- Do not drink ETOH!

TRANSDERMAL PATCH

- ✓ Remove old patch before placing a new one
- ✓ Dispose old patch in the sharps container
- ✓ Date & initial the patch
- ✓ Do not apply over hair
- ✓ Rotate sites
- ✓ Avoid the sun or heat (it increases absorption)

WHEN TO STOP THE MEDICATION:

- Respiratory depression
 - RR < 12
- If the client is unarousable

SULFONAMIDES & FLUOROQUINOLONES

OVERVIEW OF ANTIBIOTICS

- * Antibiotics and antibacterials are used interchangeably
- * Antibiotics are only used for bacterial infection (not viral)
- * Finish the entire prescription of antibiotics (even if you are feeling better)
- * NO alcohol (antibiotics are hard on the liver)
- * A culture & sensitivity test
 - **CULTURE** is a test to determine the type of bacteria
 - **SENSITIVITY TEST** is to determine what kind of medication will work best
 - Always obtain cultures *before* administering an antibiotic

SUPERINFECTION

Antibiotics disrupt the "normal flora" which can cause a super infection (secondary infection)

GENERIC

TRADE NAME

sulfadiazine	-
sulfasalazine	Azulfidine
sulfamethoxazole	Bactrim

PREFIX: SULFA-

SULFONAMIDES "SULFA DRUGS"

ACTION

Bacteriostatic (slow-growing)
Inhibit folic acid metabolism.
It slows the growth of the bacteria enough for the body to take over with its own defense mechanic (WBCs)



NURSING CONSIDERATIONS

- Increase **FLUIDS** intake because sulfas dry out the body
- Since sulfas cause photosensitivity, we want to use sunblock and avoid the sun!
- Take folic acid daily
- Patient may bruise easily
 - Monitor skin & handle with care



SIDE EFFECTS

- **GI UPSET!**
 - Nausea, vomiting, anorexia, diarrhea, abdominal pain, stomatitis
- Chills / fever
- Crystalluria
 - Crystals in the urine
- Photosensitivity
 - Increased risk for sunburn!
- **HEMATOLOGIC CHANGES**
 - Leukopenia (↓ WBCs)
 - Thrombocytopenia (↓ platelets)
 - Aplastic anemia (↓ RBCs)



USES


- UTIs (commonly caused by E.coli)
- Acute otitis media
- Ulcerative colitis
- Topical: used for burn wounds

CONTRAINDICATIONS

- Hypersensitive to sulfa drugs
- Allergy to sulfonyleureas like Glyburide (oral antibiotics)



Sulfas think SUN BURN!



GENERIC

TRADE NAME

ciprofloxacin	Cipro
gemifloxacin	Factive
ofloxacin	Floxin
moxifloxacin	Avalox

SUFFIX: -FLOXACIN

FLUOROQUINOLONES

ACTION

Interferes with the synthesis of bacterial DNA
(Causes death of the bacterial cell)



CONTRAINDICATIONS

- Clients with a history of hypersensitivity to the fluoroquinolones
- Children <18 years old
- Give with caution to:
 - Diabetics, those with renal impairment, history of seizures, & the elderly.



SIDE EFFECTS

- **GI UPSET!**
 - Nausea, diarrhea, abdominal pain
- Dizziness
- Photosensitivity



Your Tendon is near the **FLOOR** & can rupture due to **FLOOR**oquinolones

↑ **RISK FOR TENDONITIS & TENDON RUPTURE** (Especially the elderly taking corticosteroids)

USES

- Lower respiratory infections
- Bone & joint infections
- UTIs
- STIs
- Infections of the skin
- Ophthalmic solutions for eye infections

NURSING CONSIDERATIONS

- Fluoroquinolones cause photosensitivity. We want to use sunblock and avoid the sun!
- Take on an **EMPTY** stomach w/ full glass of water



PENICILLIN & CEPHALOSPORINS

GENERIC

- penicillin G
- penicillin V
- amoxicillin
- ampicillin
- piperacillin
- oxacillin

SUFFIX: -CILLIN

PENICILLIN

4 TYPES

- Natural
- Penicillinase-resistant
- Aminopenicillins
- Extended-spectrum

SIDE EFFECTS

- GI UPSET!**
 - Stomatitis & dry mouth
 - Gastritis, nausea, vomiting, diarrhea, & abdominal pain
- ORALLY** - Inflammation of the tongue (Glossitis)
- IM INJECTION** - Pain at the site
- IV INJECTION** - Irritation & inflammation (Phlebitis)

ACTION

Broad Spectrum Antibiotic

Inhibits the integrity of the bacterial cell wall



USES

- (UTIs)
- Septicemia
- Meningitis
- Intra-abdominal infections
- STIs (syphilis)
- Respiratory infections (pneumonia)

Penicillins are commonly used as Prophylaxis (prevention) against secondary infections.

CONTRAINDICATIONS

- History of allergies to **CEPHALOSPORINS** or **PENICILLIN**
- Renal disease, asthma, bleeding disorders, GI disease



NURSING CONSIDERATIONS

- Pregnancy & breast-feeding safe
- Penicillin makes oral contraceptive ineffective
 - use additional contraceptive
- Educate: take with **FOOD** to ↓ GI upset
- Penicillin allergy is very common!

Penicillin bumps the Pill

CROSS SENSITIVITY

Ask about allergy to **PENICILLIN** or **CEPHALOSPORINS** before administering the first dose!
A client who is allergic to penicillin also may be allergic to cephalosporins.

1ST GENERATION MEDICATIONS

GENERIC	TRADE NAME
cefadroxil	-
cefazolin	Ancef
cephalexin	Keflex

2ND GENERATION MEDICATIONS

GENERIC	TRADE NAME
cefaclor	-
cefoxitin	Mefoxin
cefotetan	-

3RD GENERATION MEDICATIONS

GENERIC	TRADE NAME
cefdinir	-
ceftriaxone	Rocephin
cefotaxime	Claforan

PREFIXES & SUFFIXES:
-CEF- & -CEPH-

CEPHALOSPORINS

ACTION

Bactericidal - kills bacteria

(Causes death of the bacterial cell)



USES

- Otitis media
- Respiratory infections
- Bone infections
- UTIs
- Use prophylactically pre-opt, intra-opt, and post-opt to prevent infection during surgery.

CONTRAINDICATIONS

- History of allergies to **CEPHALOSPORINS** or **PENICILLIN**
- Administer with caution: clients with renal disease, hepatic impairment, bleeding disorder




NURSING CONSIDERATIONS

- Cephalosporins make oral contraceptive ineffective
 - use additional contraceptive
- Do NOT drink alcohol while on this medication.

SIDE EFFECTS

- GI UPSET!**
 - Nausea, vomiting, diarrhea
- Dizziness
- Malaise
- Heartburn
- Fever
- Nephrotoxicity
- Aplastic anemia (↓ RBCs)
- Stevens-Johnson syndrome (SJS)
- Toxic epidermal necrolysis
- IV INJECTION** - Irritation & inflammation (Phlebitis)
- IM INJECTION** - Pain at the site



TETRACYCLINES & AMINOGLYCOSIDES

GENERIC

TRADE NAME

tetracycline	-
doxycycline	Atridox
minocycline	Arestin
demeclocycline	-

SUFFIX: -CYCLINE

TETRACYCLINES

USES

- **SKIN**
 - Skin & soft tissue infection
 - Severe acne
- Rocky mountain spotted fever
- Helicobacter Pylori (H. pylori)

NURSING CONSIDERATIONS

- Fluoroquinolones cause photosensitivity. We want to use sunblock and avoid the sun!
- Tetracyclines make oral contraceptives ineffective
 - Use additional contraception
- Take on an EMPTY stomach with a full glass of water
- Causes tooth discoloration
 - Do not give to children younger than 9
- Sit up for 30 min after taking medication
 - Do not lay down
 - Pill induced esophagus (HEARTBURN & scaring of the esophagus!)
- Avoid calcium/dairy products
 - These prevent the absorption of the drug

ACTION

Bacteriostatic (slow-growing)
Inhibits bacterial protein synthesis



Tetra think Teeth

SIDE EFFECTS

- **GI DISTRESS!**
 - Nausea / vomiting / diarrhea
 - Stomatitis
- Skin rashes
- Photosensitivity reaction



CONTRAINDICATIONS

- Known allergy to tetracyclines
- Contraindicated in lactation



Tetracyclines think Toxic to the developing fetus

GENERIC

TRADE NAME

gentamicin	-
kanamycin	-
neomycin	-
streptomycin	-

SUFFIXES: -MYCIN, -MICIN

AMINOGLYCOSIDES

ACTION

Bactericidal - kills bacteria
Blocks the ribosome from reading the mRNA. Then the bacterial can't multiply.



NURSING CONSIDERATIONS

- Monitor:
 - Renal status
 - Neuro status
 - Respiratory status
- Evaluate clients comments related to any hearing issues

SIDE EFFECTS

- **GI DISTRESS!**
 - Nausea / vomiting / anorexia
- Rash & hives

Aminoglycosides are A mean antibiotic because they have very harmful side effects.

CONTRAINDICATIONS

- Known allergy to aminoglycosides
- Hearing loss
- Musculoskeletal disorders (Myasthenia gravis & Parkinson's disease)
- Contraindicated for location



USES

- **BOWEL PREPARATION:** Decrease normal flora in the GI for those having abdominal surgery
- Management of hepatic coma
 - Decreasing the ammonia in the intestines



NEPHROTOXICITY

Hurts the kidneys: Proteinuria hematuria, & increase BUN & Creatinine.



OTOTOXICITY

Hurts the ears: Tinnitus, vertigo, hearing loss, which may be permanent.



NEUROTOXICITY

Hurts the brain: Numbness, tumors, convulsions, muscular paralysis.

DIURETICS OVERVIEW

LOOP DIURETIC

GENERIC	TRADE NAME
furose mid e	Lasix
bumeta ni d	Bumex
torse mi d	Demadex

SUFFIX: -NIDE, - MIDE

⚠️ Potent (strong) diuretic

ACTION

- Inhibit reabsorption of Na^+ & Cl^-

Acts on 3 sites

=

↑ reabsorption

PURPOSE

- Hypertension
- Heart failure
- Renal disease
- Edema
- Pulmonary edema

SIDE EFFECTS

- ↓ Hypokalemia ⚠️
- ↓ Hypotension
- ↑ Hyperglycemia
- Photosensitivity
- ↓ Hyponatremia
- Dehydration

NURSING
CONSIDERATIONS

- Obtain baseline vital signs
- Adm. furosemide SLOWLY (rapid adm. can cause ototoxicity) 🦻
- Replace K^+ if $< 3.5 \text{ mEq/L}$



NORMAL POTASSIUM
3.5 - 5.0

POTASSIUM WASTING!

THIAZIDE DIURETIC

GENERIC	TRADE NAME
hydrochloro thia zide	Microzide
chloro thia zide	Diuril
methyclo thia zide	-

SUFFIX: -THIAZIDE

ACTION

- Inhibit reabsorption of Na^+ & Cl^-

- Excretion of Na^+ , Cl^- , & H_2O

↑ UOP

=

↓ blood volume

PURPOSE

- Hypertension
- Heart failure
- Renal disease
- Cirrhosis
- Edema
- Corticosteroids
- Estrogen therapy

SIDE EFFECTS

- ↓ Hypokalemia
- ↓ Hypotension
- ↓ Hyponatremia
- ↓ Libido
- ↑ Hyperglycemia
- Photosensitivity
- Dehydration
- Azotemia

POTASSIUM WASTING!

NURSING
CONSIDERATIONS

- Obtain baseline vital signs
- Monitor intake & output
- Give w/ meals to ↓ GI upset
- Replace K^+ if $< 3.5 \text{ mEq/L}$
➔ **NEVER** give K^+ IV push
- Avoid giving to pt.'s with gout
- Monitor renal function
- Daily weights
➔ Same time, same scale!
- Clients with a **sulfa allergy** should avoid thiazide diuretics ⚠️

DIURETICS OVERVIEW



DIURESIS THE BODY
Diuretics = Diuresis = Dry inside

- **WHERE SODIUM GOES . . . WATER FLOWS!**
- Sodium makes us retain water
 - ➔ Low sodium diet (**SODIUM SWELLS!**)
- Give diuretics in the morning, not at night
 - ➔ You don't want your client peeing all night long (Nocturia)
- Instruct the client to make slow position changes (diuretics cause orthostatic hypotension)
- Monitor...
 - ➔ Daily weights (report 2-3 lbs weight gain)
 - ➔ Intake & output
 - ➔ Vital signs
 - ➔ Potassium levels

OSMOTIC DIURETIC

GENERIC	TRADE NAME
mannitol	Osmitrol



ACTION	PURPOSE	SIDE EFFECTS	NURSING CONSIDERATIONS
<ul style="list-style-type: none"> • ↑ the thickness of the filtrate so water can't be reabsorbed • Excretion of Na⁺ & Cl⁻ 	<ul style="list-style-type: none"> • Treatment of cerebral edema • ↓ intraocular pressure (IOP) 	<ul style="list-style-type: none"> • Edema • Blurred vision • Nausea, vomiting, & diarrhea • Urinary retention 	<ul style="list-style-type: none"> • Only administered IV • May crystallize (<i>check solution before adm.</i>) • Perform neuro assessment & LOC (<i>if using for cerebral edema</i>)

K⁺ SPARING DIURETIC

GENERIC	TRADE NAME
spironolactone	Aldactone



S think **S**paring

ACTION	PURPOSE	SIDE EFFECTS	NURSING CONSIDERATIONS
<ul style="list-style-type: none"> • Blocks aldosterone ("salt water" hormone) • Lets fluid out of the body, into the potty!  • Excretion of Na⁺ & H₂O <p>NOT K⁺ (saves potassium)</p>	<ul style="list-style-type: none"> • Hypertension • Edema • Hypokalemia • Hyperaldosteronism • Cross-sex hormonal therapy <p>SPIRONOLACTONE INHIBITS TESTOSTERONE</p>	<ul style="list-style-type: none"> • Hyperkalemia (> 5.0) • Diarrhea • Gastritis • Drowsiness • Erectile dysfunction • Gynecomastia (<i>enlargement of the breasts in men</i>) <p>EDUCATE: GYNECOMASTIA IS USUALLY REVERSIBLE AFTER THERAPY HAS STOPPED</p>	<ul style="list-style-type: none"> • Avoid eating foods high in potassium (<i>green leafy veggies, melons, bananas, avocado, etc.</i>)  <ul style="list-style-type: none"> • Avoid salt substitutes & potassium supplements • Monitor K⁺ levels

⚠ Watch out for HYPERKALEMIA (K⁺ > 5.0 mEq/L)

ANTIHYPERLIPIDEMIC DRUGS


OVERVIEW

- ♦ Atherosclerosis is when lipids stick to the blood vessel walls which can obstruct blood flow
- ♦ The goal of all antihyperlipidemic drugs is to lower lipid levels in the blood

CHOLESTEROL

LDL → Want LOW Levels (<100 mg/dL)
Low Density Lipoprotein
BAD CHOLESTEROL

HDL → Want HIGH Levels (>60 mg/dL)
High Density Lipoprotein
HAPPY CHOLESTEROL



HMG-COA REDUCTASE INHIBITORS "STATINS"

USES



- ♦ Hyperlipidemia
- ♦ PRIMARY PREVENTION: Preventable treatment for patients at risk for coronary artery disease (CAD)
- ♦ SECONDARY PREVENTION: Stabilizes fatty plaques in clients with current coronary artery disease (CAD)

LOWERS CHOLESTEROL

ACTIONS

- ♦ Inhibits the enzyme HMG-CoA Reductase
- ♦ Statins are not a cure!

SIDE EFFECTS

NEURO

- Headache
- Nausea
- Dizziness

GI

- Constipation
- Cramping
- Abdominal pain
- Hyperglycemia

GENERIC	TRADE NAME
atorvastatin	Lipitor
fluvastatin	Lescol
lovastatin	Altoprev
pitavastatin	Livalo
simvastatin	Zocor
rosuvastatin	Crestor


SUFFIX: -STATIN

NURSING CONSIDERATIONS

- ♦ Monitor liver enzymes
 - ↳ ALT/AST
- ♦ Monitor therapeutic response
 - ↳ Statins should lower LDL, & increase HDL
- ♦ Avoid grapefruit consumption
 - ↳ Increases risk for toxicity of statins
- ♦ Statins are **pregnancy category X** & should not be taken while breastfeeding
- ♦ Monitor for signs of **rhabdomyolysis** because statins have been associated with this

RHABDOMYOLYSIS

- ♦ Rare condition where the muscles are damaged
- ♦ Myoglobin leaks into the blood which can cause **kidney damage**
- ♦ Signs & symptoms:
 - ↳ Muscle pain, tenderness, or weakness
 - ↳ Accompanied by malaise or fever
 - ↳ ↑ creatine kinase levels
 - ↳ Dark urine color (tea or cocoa like urine)



BILE ACID RESINS

USES



- ♦ Hyperlipidemia
- ♦ Gallstone dissolution
- ♦ Pruritus associated with partial biliary obstruction

SIDE EFFECTS

GI

- ♦ Constipation
- ♦ Increase risk for bleeding R/T Vit K malabsorption
- ♦ Vitamin A & D deficiencies

ACTIONS

Bile is made & secreted by the **liver**

↓

Then, it's stored the **gallbladder**

↓

Once emulsified, the fats & lipids are absorbed in the **intestines**

↓

Bile Acid Resins binds to the bile acid to form an insoluble substance (can not be absorbed by the intestine)

↓

So it's excreted with the **feces**

↓

↓ bile acids = **liver** uses cholesterol to make more bile = ↓ cholesterol



NURSING CONSIDERATIONS

GENERIC	TRADE NAME
cholestyramine	Prevalie
colestipol	Colestid
colesevelam	Welchol

- ♦ Bile acid resins may interfere with the digestion of fats, preventing the absorption of **fat-soluble vitamins**

All Kids Eat Donuts

- ↳ Vitamin A & D may be given in a water-soluble for long term therapy
- ♦ Bile acid resins may cause constipation, so educate to...
 - ↳ Increase fluids, fibers
 - ↳ Exercise regularly
 - ↳ Use stool softener


ANTIHYPERTENSIVES

ACE INHIBITORS

angiotensin-converting enzyme inhibitors

GENERIC	TRADE NAME
capto pril	-
enalap ril	Vasotec
fosino pril	-
lisino pril	Prinivil

SUFFIX: -PRIL

USES 

- † Hypertension
- † Heart Failure

ACTION

Dilates blood vessels, which lowers blood pressure. They do not directly affect the heart rate.

- † Inhibits RAAS Renin-Angiotensin-Aldosteron-System
- † RAAS is the main hormonal mechanism involved in regulating the blood pressure
- † ACE converts angiotensin I → angiotensin II (a powerful vasoconstrictor)
- † Inhibiting ACE will inhibit this vasoconstricting effect, decreasing blood pressure!

SIDE EFFECTS

A = ANGIOEDEMA
C = COUGH (DRY)
E = ELEVATED K+

Orthostatic Hypotension
Dizziness


NURSING CONSIDERATIONS

- † Assess BP & pulse routinely
- † Monitor for hypotension
 - Educate on changing positions slowly
- † Monitor K+ levels
 - Normal 3.5 - 5.0
 - Educate to avoid foods high in potassium & avoid salt substitutes
- † Assess for angioedema
 - Swelling of the area beneath the skin or mucosa (deep edema)
 - **DANGEROUS:** swelling of the face & mouth
- † Educate to not suddenly stop the medication it can cause **rebound hypertension** (needs to be tapered off)
- † Ace inhibitors are contraindicated in pregnancy due to the teratogenic effects on the fetus

BETA BLOCKERS

GENERIC	TRADE NAME
acebut olol	Sectral
metop rolol	Corgard
propran olol	Inderal
nad olol	Bystolic

SUFFIX: -OLOL



USES 

- † Hypertension
- † Stable angina
- † Chronic / compensated heart failure (not acute heart failure)
- † Dysrhythmias

ACTION

- † Blocks norepinephrine & epinephrine (fight or flight hormones)
- † Blocks the negative effects of the sympathetic nervous system
 - ➔ Beta blockers can be selective or non-selective
 - Meaning they can block different beta sites (beta 1 and/or beta 2)

↓ Resistance
↓ Workload
↓ Cardiac Output

BETA 1 (one heart) 
BETA 2 (two lobes) 

SIDE EFFECTS

THE B'S OF BETA BLOCKERS

- † **B**radycardia & heart **B**locks
- † **B**reathing problems
 - ➔ Bronchi spasms
- † **B**ad for heart failure patients (in an acute setting)
- † **B**lood sugar masking
 - ➔ Masks S&S of hypoglycemia (low blood sugar)
- † **B**lood pressure lowered - Hypotension

NURSING CONSIDERATIONS

- † Monitor for hypotension
- † Educate on changing positions slowly
- † Do not give non-selective beta blockers to asthma patients or COPD patients (*remember: non-selective works on Beta1 & **Beta2** = Lung constriction*)
- † Educate to not suddenly stop the medication. It can cause rebound hypertension (*needs to be tapered off*)
- † Monitor for S&S of heart failure
 - These medications produce inotropic effects (↑ contraction strength of the ♥)
 - S&S of ♥ failure: Wet lung sounds, weight gain, edema, etc

ANTIHYPERTENSIVES

CALCIUM CHANNEL BLOCKERS

**VERY
NICE
DRUGS**

GENERIC	TRADE NAME
Verapamil	Calan
Nifedipine	Procardia
Diltiazem	Cardizem
amlodipine	Norvasc
nicardipine	Cardine

SUFFIXES: -DIPINE, -AMIL

USES

LOWER HR & BP

- Hypertension
- Angina
- Dysrhythmias



SIDE EFFECTS

- Orthostatic hypotension
- Dizziness
- Flushing
- Headache
- Peripheral edema
- Constipation

**ALL COMMON
SIDE EFFECTS**

ACTION

Blocks movement of calcium
(↓ calcium = ↓ available for
transmission of nerve impulses)

- Relaxes blood vessels
- ↓ blood pressure
- ↑ supply of oxygen to the heart
- ↓ ♥'s workload

NURSING CONSIDERATIONS

- Antihypertensives cause **orthostatic hypotension**
 - Change positions slowly
 - Sit on the side of the bed for a few minutes before standing
- Educate to not suddenly stop the medication. It can cause **REBOUND HYPERTENSION** (needs to be tapered off).

- Do not drink grapefruit juice
- Leg elevation & compression to reduce edema
- To help with **CONSTIPATION**:



**CAN CAUSE
SEVERE
HYPOTENSION!**



ANTICOAGULANTS

Prevents new clots or prevents current clots from getting bigger! Anticoagulants do not dissolve clots & do not thin the blood.
Anticoagulants are use for clients who are at an increased risk for **CLOT FORMATION!**

WARFARIN

GENERIC	TRADE NAME
warfarin	Coumadin

ACTION

- Interferes with the production of **VITAMIN K**
 - ➔ **USED IN THE LIVER TO MAKE CLOTTING FACTORS**
- ↓ of clotting factors II (prothrombin), VII, IX, and X. (Prothrombin is required for the clotting)

USES

- LONG-TERM THERAPY**
- Works slowly (a few days to take effect)



ROUTES

- Orally
- IV

MOST COMMON

SAFE DURING PREGNANCY? **NO!**

ANTIDOTE: VITAMIN K

NURSING CONSIDERATIONS

- Educate client to be consistent with their vitamin K food intake (green leafy vegetables, liver, etc.)
- Antibiotics increase the risk for **BLEEDING** (they increase INR)
- Will have freq. blood test to check therapeutic range
- Educate to take the pill at the same time every day

Can a patient be on *both* at the same time?

YES!

Commonly used together. Gives time for Warfarin to kick in!

HEPARIN

ACTION

Heparin inhibits the formation of fibrin clots. Inhibits the conversion of fibrinogen to fibrin (inactivates factors needed for the clotting)

USES

- SHORT-TERM THERAPY**
- Works quickly



ROUTES

- NOT** given orally
- Given by injection (IV or subq)
- IV drip

Heparin is inactivated by gastric acid in the stomach

SAFE DURING PREGNANCY? **YES!**

ANTIDOTE: PROTAMINE SULFATE

LOW MOLECULAR WEIGHT HEPARIN (LMW)

GENERIC	TRADE NAME
enoxaparin	Lovenox
dalteparin	Fragmin

SUFFIX: -PARIN

LMW HEPARIN IS ADMINISTERED:

- Subq in the belly
- 2 inches from the umbilicus
- 90 degree angle!
- After subq injection, it's common to have bruising, irritation, & pain!
- Do not massage injection site after

For LMW heparins, we don't look at blood coags. We want to monitor **PLATELET COUNT!**

Heparin Induced Thrombocytopenia (HIT)

Should check platelets while on LMW.
Normal PLT count: **150,000 - 450,000**

COAGULATION

NOT ON ANY ANTICOAGULANT:

PT: 10 - 12 seconds
INR: < 1
aPTT: 30 - 40 seconds

ABBREVIATIONS:

PTT: Prothrombin Time
aPTT: Activated Partial Thromboplastin Time
INR: International Normalized Ratio

INTERPRETATION:

Numbers are **TOO** high = Patient will die (increased bleeding)
Numbers are **LOW** = Clots will **GROW**

WARAFIN
Measured with: INR

THERAPUTIC RANGE:
1.5 - 2 times the normal value
INR: 2 - 3
INR: 2.5 - 3.5
(Heart valve replacement)



HEPARIN
Measured with: aPTT

THERAPUTIC RANGE:
1.5 - 2 times the normal value
aPTT: 47 - 70 SECONDS

DIGOXIN

MEDICATION CLASS: CARDIAC GLYCOSIDES

USES

- Heart failure
- Cardiogenic shock
- Antiarrhythmic
 - Atrial fibrillation



ACTION

- **(+) INOTROPIC ACTIVITY**
 - Increases the force of the contraction = increased cardiac output
- **(-) CHRONOTROPIC:** beats slower
- **(-) DROMOTROPIC:** slows impulses sent through AV node, able to squeeze more blood

TOXICITY

THERAPEUTIC RANGE:
0.8 - 2.0 NG/ML

> 2 = Think Toxic

SIGNS OF TOXICITY

Report these to the HCP



GI SYMPTOMS

Nausea, vomiting, diarrhea

EARLY SIGN



VISUAL SYMPTOMS

Blurred vision, yellow or green vision, halo effect around dark object



NEUROLOGICAL SYMPTOMS

Headache, drowsiness, confusion, disorientation

ANTIDOTE: DIGIBIND

GENERIC

digoxin

TRADE NAME

Lanoxin

CAUSES OF TOXICITY?

- D** Decreased potassium (**HYPO**kalemia) <3.5 mEq/L
 - Potassium wasting diuretics (Loop)
- I** Injured kidneys
- G** GFR decreased (the elderly)


Digoxin is almost solely excreted by the kidneys

NURSING CONSIDERATIONS

- **HOLD THE MEDICATION IF**
 - Adults: <60 bpm
 - Children: <70 bpm
 - Infants: <90-110 bpm
- **KEEP ALL APPOINTMENTS:** drug levels & electrolytes will be monitored

THE APICAL PULSE
MUST BE ASSESSED
FOR 1 MINUTE
BEFORE
ADMINISTERING
DIGOXIN

The **APICAL PULSE** is located at the fifth intercostal midclavicular space.



- The apex of the heart
- Point of maximal impulse (PMI)
- Mitral valve

NITROGLYCERIN (NTG)

MEDICATION CLASS: ANTIANGINALS

USES

- Angina (chest pain)
- Prevent angina attacks
- Acute coronary syndrome



SIDE EFFECTS

- THE H's**
- H** = Headache
 - H** = Hypotension (orthostatic)
 - H** = Hot flushing of the face

- Rash
- Sublingual
 - Tingling / burning sensation
- Transdermal
 - Contact dermatitis

ALARMING SIGNS

- Lack of coordination
- Lightheadedness
- Pallor
- Irritable


ACTION

VASODILATOR

Dilators do the following:

- D** Decrease blood pressure
- D** Dilates vessels
- D** ↓ Vascular resistance

- D** = decrease cardiac workload
- D** = decrease oxygen consumption



VASODILATION

NORMAL BLOOD VESSEL

CONTRAINDICATIONS

- Known hypersensitivity to nitroglycerin
- Allergy to adhesive (transdermal)
- Clients taking phosphodiesterase (PDE) inhibitors
- Head trauma, cerebral hemorrhage
- Severe anemia



QUICK VS. SLOW ONSET

QUICK

- IV
- Sublingual tabs
- Translignal spray

SLOW

- Nitro patch
- Nitro ointment
- Sustained-release tablets

NURSING CONSIDERATIONS

DO NOT TAKE with phosphodiesterase (PDE) inhibitors (erectile dysfunction (ED) drugs)

-  Ends in "-afil" Like sildenafil (viagra)
- Causes dangerously low blood pressure resulting in death

- Monitor blood pressure
 - Stop the medication if systolic BP drops below 100 or the baseline drops below 30 mmHg
- Increased risk for falls due to orthostatic hypotension
 - Educate: rise slowly when getting up



LONG ACTING NITRATES DESIRABLE OUTCOME:

The client can perform activities without chest pain (shower, get dressed, etc)



PATIENT EDUCATION

TOPICAL & TRANSDERMAL PATCH

- Remove prior dose
- Rotate sites
- Place over a clean/hairless area
- Wear gloves
- Do not rub nitro ointment into the skin, it can cause rapid absorption!
- Patches can be worn in the shower



SUBLINGUAL NTG OR SPRAY



1 tab/spray sublingual every 5 minutes, up to 3 doses.

If angina is not relieved or is worse 5 min after the first dose, call 911!

SUBLINGUAL OR BUCCAL

- Place **BUCCAL** tablet between the cheek and gum
- Place **SUBLINGUAL** under the tongue
- Rinse with water before placing the tablets in your cheek



-  Keep in original container (dark, glass bottle) in a dry, cool place.
-  Do not swallow or chew these tablets



CORTICOSTEROIDS

GENERIC	TRADE NAME
prednisone	Deltasone
hydrocortisone	Hydrocort
dexamethasone	Ozurdex
fluticasone	Flovent HFA
beclomethasone	–
flunisolide	Aerospan
ciclesonide	Zetonna


SUFFIXES: -SONE, -ASONE, -IDE

ACTION



ANTI-INFLAMMATORY EFFECTS!

➔ They reduce the number of mast cells in the airway



THERAPUTIC USES

- COPD
- Rheumatoid arthritis
- Lupus

Can also be administered:
IV, IM, PO, rectally, ocularly

TOPICAL CORTICOSTEROIDS

Dermatitis
Rashes
Eczema
Insect bites

INHALED CORTICOSTEROIDS (ICSS)

Chronic asthma
Nasal polyps
& rhinitis

SIDE EFFECTS

S's of Steroids

Steroids cause...

- 👉 **Sugar:** Hyperglycemia
- 👉 **Soft Bones:** Causes osteoporosis
- 👉 **Sick:** Decreased immunity / sepsis
- 👉 **Sad:** Depression
- 👉 **Salt:** Water & salt retention (hypertension)
- 👉 **Sex:** Decreased libido
- 👉 **Swollen:** Water gain = weight gain
- 👉 **Sight:** Risk for cataracts

PATIENT EDUCATION

(Long-term corticosteroid replacement)

- 👉 **Report signs of an INFECTION**
 - Corticosteroids are immunosuppressing and can cause an infection
 - Since they are anti-inflammatory, it may hide the fact that the client has an infection
- 👉 **Increase CALCIUM in the diet**
 - Corticosteroids can cause osteoporosis & muscle weakness
- 👉 **Yearly optometrist appointment**
 - Corticosteroids may cause **CATARACTS**
- 👉 **STRESS OR SURGERY causes a decrease in cortisol**
 - You may need to increase your dose in times of stress
- 👉 **Never stop steroids suddenly**
 - Slowly taper off the medication!

IMPORTANT TEACHING!

After administration, rinse the mouth to decrease the risk of contracting a possible fungal infection from candidiasis

THRUSH: a type of yeast infection



TAKING Bronchodilators & Corticosteroids?

- 1 **Bronchodilator** first (to help open up the airways)
- 2 Wait 5 minutes
- 3 Administer the **Corticosteroid**



B comes before **C**
in the alphabet



BRONCHODILATORS (SABA & LABA)

SHORT-ACTING BETA2 AGONISTS (SABAs)

GENERIC	TRADE NAME
albuterol	Proventil
epinephrine	Adrenalin
levalbuterol	Xopenex
terbutaline	-

Think **Albuterol** is for **Acute Asthma Attacks**

LONG-ACTING BETA2 AGONISTS (LABAs)

GENERIC	TRADE NAME
salmeterol	-
formoterol	Foradil
indacaterol	Arcapta
arformoterol	Brovana

Think **Salmeterol** is for **Slow and Steady** working a **LONG** time

SUFFIX: -TEROL

SIDE EFFECTS

- Tachycardia
- Palpitations
- Cardiac arrhythmias
- Hypertension
- Nervousness & anxiety
- Insomnia

REMEMBER Think **FIGHT OR FLIGHT!**



ACTION

BRONCHO-DILATORS

Dilates (opens up) the bronchi

When an agonist binds to the beta-2 receptors the sympathetic nervous system "Fight or flight" takes effect. **THE AIRWAYS RELAX AND DILATE WHICH INCREASES OXYGEN FLOW** which makes it easier to breathe.



MEMORY TRICK

To remember that beta-2 receptors are in the lungs: you have **TWO LUNGS**

To remember beta-1 receptors found on the heart: you only have **ONE HEART**.

BETA 1 (one heart)



BETA 2 (two lobes)



USES

SHORT-ACTING BETA2 AGONISTS (SABAs)

Acute symptom relief

- Bronchospasms
- Asthma

LONG-ACTING BETA2 AGONISTS (LABAs)

Long-term management

- COPD
- Chronic Bronchitis
- Prevention of bronchospasms

Airway Dysfunction

IMPORTANT TEACHING!

After administration, rinse the mouth to decrease the risk of contracting a possible fungal infection from candidiasis

THRUSH: a type of yeast infection

PATIENT EDUCATION

TAKING Bronchodilators & Corticosteroids?

- 1 Bronchodilator first (to help open up the airways)
- 2 Wait 5 minutes
- 3 Administer the Corticosteroid



MEMORY TRICK

B comes before **C** in the alphabet

ABC

BRONCHODILATOR

(Xanthine derivatives)
(Methylxanthines)

GENERIC	TRADE NAME
aminophylline	-
dyphylline	Lufyllin
oxtriphylline	Choledyl SA
theophylline	Theochron

SUFFIX: -PHYLLINE

ACTION



BRONCHO-DILATORS

Dilates (opens up) the bronchi

Stimulate the central nervous system (CNS) to promote bronchodilation.

Relaxation of the smooth muscles of the bronchi.

USES

- Relief & prevention of bronchial asthma
- Tx of bronchospasms seen in COPD

SIDE EFFECTS

- Tachycardia
- Palpitations
- ECG changes
- Nervousness & anxiety
- Irritable



THEOPHYLLINE

Therapeutic levels
10 - 20 mcg/dL

Toxic **>20 mcg/dL**

SIGNS OF TOXICITY

- Tonic clonic seizures
- Tachycardia & dysrhythmias

CHOLINERGIC BLOCKING (Anticholinergic)

ALSO CALLED:



ANTICHOLINERGIC DRUGS

CHOLINERGIC BLOCKERS

PARASYMPATHOLYTIC DRUGS

GENERIC	TRADE NAME
acclidinium	Tudorza
umeclidinium	Incruse
ipratropium	Atrovent
tiotropium	Spiriva

SUFFIXES: -TROPIMUM, -CLINDIDIUM

Inhaled anticholinergic medications

ACTION



Cholinergic blocking drugs **BLOCK THE PARASYMPATHETIC NERVE** that causes the airway to constrict.

By blocking this, it allows the airways to remain open.

SIDE EFFECTS

BLOCKS SECRETIONS → Dry Inside



- Can't See - Blurred vision
- Can't Pee - Dysuria
- Can't Spit - Dry mouth
- Can't Poop - Constipation

RESPIRATORY USES

Prevention of bronchospasms associated with COPD

PATIENT EDUCATION

- Prevent constipation
 - Increase fluids & fiber
- To help with the dry mouth, increase fluids & suck on hard candies.



Anticholinergic drugs are used for many other purposes as well, such as:
PARKINSONISM, PEPTIC ULCER, VAGAL NERVE-INDUCED BRADYCARDIA & PREOPERATIVE REDUCTION OF ORAL SECRETIONS.

LITHIUM CARBONATE

MOOD STABILIZER:

Known for its side effects and narrow therapeutic range

THERAPEUTIC RANGE:

0.6 - 1.2 mEq/L




USES

Bipolar disorder

Helps regulate the "mood swings" (depression & mania)

Lithium is a Long-term treatment



ADVERSE REACTIONS

- ➔ Nausea/drowsiness/fatigue
- ➔ Thirst
- ➔ Dry mouth
- ➔ Weight gain

TOXICITY!

- * Confusion
- * Blurred vision
- * Diarrhea
- * Tinnitus
Ringing in ears
- * Slurred speech
- * Coma
- * Convulsions
- * Excessive urination
- * Excessive thirst
- * Tremors/ataxia

TOXICITY LEVELS

Mild: 1.5 - 2 mEq/L

Moderate: 2 - 3 mEq/L

Severe: > 3 mEq/L

HOW DOES TOXICITY HAPPEN?

- ☞ Dehydration causes ↑ lithium levels in blood
- ☞ Hyponatremia
- ☞ Old age & kidney failure
↓ GFR = lithium builds up in the blood

- Excessive sweating such a high fever
- Diarrhea
- Diuretic therapy

EDUCATION

- * Carry ID that shows you are taking lithium
- * Educate on signs & symptoms of toxicity
- * Educate and stress importance of taking medication regularly
- * Serum lithium levels should be checked every 1-2 months
- * Do not operate heavy machinery or drive
- * Educate on drinking plenty of water to avoid dehydration (therefore avoiding toxicity)
- * Avoid starting a low salt diet
Sudden ↓ in salt = ↑ in lithium






CONTRAINDICATION

- * Contraindicated in pregnancy & breastfeeding
- * Renal/cardiovascular disease
- * Dehydrated patients
Excessive diarrhea or vomiting
- * Receiving diuretics
- * Sodium depletion
- * Hypersensitivity to tartrazine
- * Avoid NSAIDs
↓ renal blood flow = ↑ risk for toxicity





Contraceptives may be prescribed

ANTIDEPRESSANT DRUGS

	SSRIs Selective serotonin reuptake inhibitor	SNRIs / DNRI Serotonin / Norepinephrine & Dopamine / Norepinephrine reuptake inhibitor																								
ACTION	Inhibits uptake of serotonin = ↑ serotonin  MEMORY TRICK Think S miley S erotonin	Affects serotonin, norepinephrine & dopamine																								
USES	<ul style="list-style-type: none"> • Depression • Anxiety • OCD • Eating disorders 	<ul style="list-style-type: none"> • Depressive episodes • Anxiety disorders • Fibromyalgia • Diabetic neuropathy pain 																								
SIDE EFFECTS	<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;">  NEURO <ul style="list-style-type: none"> • Headache • Tremors • Difficulty sleeping </div> <div style="width: 45%;">  GI <ul style="list-style-type: none"> • Nausea • Dry mouth / thirst • Constipation • Urinary retention • Sexual dysfunction </div> </div> <div style="margin-top: 10px;"> <p>3 S's OF SSRIs</p> <ul style="list-style-type: none"> • Serotonin syndrome • Sexual dysfunction • Stomach issues </div> <div style="margin-top: 10px; border: 1px solid red; padding: 5px;"> <p>SEROTONIN SYNDROME</p> <ul style="list-style-type: none"> • Too much serotonin in the brain • Mental changes • Tachycardia • Tightness in muscles • Difficulty walking • ↑BP & temp </div>	<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;">  NEURO <ul style="list-style-type: none"> • Headache • Dizziness • Vertigo • Photosensitivity • Agitation/tremors • Insomnia </div> <div style="width: 45%;">  GI <ul style="list-style-type: none"> • Dry mouth/thirst • Dehydration • Constipation • Nausea/diarrhea </div> </div>																								
NURSING CONSIDERATIONS	<ul style="list-style-type: none"> • May take 4-6 weeks to take effect • Take medication in the morning • First line drug for depression/anxiety <div style="border: 1px solid red; padding: 5px; margin-top: 10px;"> <p>⚠ SUICIDE WARNING ⚠</p> <p>A client who had suicidal plans may now have the energy due to the medication to carry out the plans!</p> </div>	<ul style="list-style-type: none"> • May take 4-6 weeks to take effect • Do not mix with TCAs or MAOIs • Zyban is used for smoking cessation. <p>Do not use it while taking bupropion for depression – it could cause overdose.</p>																								
DRUG TABLE	<table border="1" style="width: 100%; text-align: center;"> <thead> <tr> <th>GENERIC</th> <th>TRADE NAME</th> </tr> </thead> <tbody> <tr> <td>sertraline</td> <td>Zoloft</td> </tr> <tr> <td>citalopram</td> <td>Celexa</td> </tr> <tr> <td>escitalopram</td> <td>Lexapro</td> </tr> <tr> <td>fluoxetine</td> <td>Prozac</td> </tr> <tr> <td>vilazodone</td> <td>Viibryd</td> </tr> </tbody> </table> <p>SUFFIXES: -TALOPRAM, -OXETINE, -ZODONE</p>	GENERIC	TRADE NAME	sertraline	Zoloft	citalopram	Celexa	esc italopram	Lexapro	flu oxetine	Prozac	vilaz odone	Viibryd	<table border="1" style="width: 100%; text-align: center;"> <thead> <tr> <th>GENERIC</th> <th>TRADE NAME</th> </tr> </thead> <tbody> <tr> <td>bupropion</td> <td>Zyban & Wellbutrin</td> </tr> <tr> <td>duloxetine</td> <td>Cymbalta</td> </tr> <tr> <td>venlafaxine</td> <td>Effexor XR</td> </tr> <tr> <td>milnacipran</td> <td>Savella</td> </tr> <tr> <td>nefazodon</td> <td>–</td> </tr> </tbody> </table> <p>SUFFIXES: -FAXINE, -ZODONE, -NACIPRAN</p>	GENERIC	TRADE NAME	bupropion	Zyban & Wellbutrin	duloxetine	Cymbalta	venlaf axine	Effexor XR	milnac ipran	Savella	nefaz odon	–
GENERIC	TRADE NAME																									
sertraline	Zoloft																									
citalopram	Celexa																									
esc italopram	Lexapro																									
flu oxetine	Prozac																									
vilaz odone	Viibryd																									
GENERIC	TRADE NAME																									
bupropion	Zyban & Wellbutrin																									
duloxetine	Cymbalta																									
venlaf axine	Effexor XR																									
milnac ipran	Savella																									
nefaz odon	–																									

Educate on the importance of compliance

ANTIDEPRESSANT DRUGS

<h2 style="color: red;">TCAs</h2> <p>Tricyclic antidepressants</p>	<h2 style="color: red;">MAOIs</h2> <p>Monoamine oxidase inhibitor</p>																					
<p>Blocks reuptake of serotonin & norepinephrine in the brain</p>	<p>Blocks monoamine oxidase which causes ↑ in epinephrine, norepinephrine, dopamine, & serotonin, which causes stimulation of the CNS!</p>	<p>ACTION</p>																				
<ul style="list-style-type: none"> • Depressive episodes • Bipolar disorder • OCD • Neuropathy • Enuresis 	<p>Depression</p>	<p>USES</p>																				
<ul style="list-style-type: none"> • Constipation • Dry mouth • Drowsiness • Blurred vision • Orthostatic hypotension • Urine retention • Cardiotoxic <div style="border: 1px dashed black; border-radius: 50%; padding: 10px; width: fit-content; margin-left: 150px;"> <p><i>Causes heart problems in patients with pre-existing cardiac conditions or elderly clients...give with caution!</i></p> </div>	<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;">  <p>NEURO</p> <ul style="list-style-type: none"> • Orthostatic hypotension • Dizziness • Blurred vision </div> <div style="width: 45%;">  <p>GI</p> <ul style="list-style-type: none"> • Constipation • Dry mouth • Nausea/ vomiting </div> </div> <div style="border: 1px solid red; border-radius: 15px; padding: 10px; width: fit-content; margin-left: 150px;"> <p>HYPERTENSIVE CRISIS</p> <ul style="list-style-type: none"> • Headache • Stiff neck • Nausea / vomiting • Fever • Dilated pupils: </div> <div style="border: 1px dashed black; border-radius: 50%; padding: 10px; width: fit-content; margin-left: 150px;"> <p><i>Seek medical help to ↓ blood pressure</i></p> </div>	<p>SIDE EFFECTS</p>																				
<ul style="list-style-type: none"> • May take 2- 3 weeks to take effect • WAIT 14 days after being off MAOIs to start taking TCAs • Amoxapine is not an antipsychotic drug but similar to these drugs, it may cause TD & NMS (D/C the drug immediately if these symptoms occur) <div style="border: 1px dashed black; border-radius: 50%; padding: 10px; width: fit-content; margin-left: 150px;"> <p><i>Educate on the importance of compliance</i></p> </div>	<ul style="list-style-type: none"> • Can take up to 4 weeks to reach therapeutic levels • Educate on the signs & symptoms of HTN crisis • Avoid foods with Tyramine <ul style="list-style-type: none"> • Aged cheese • Fermented meats • Chocolate • Caffeinated beverages • Sour cream & yogurt 	<p>NURSING CONSIDERATIONS</p>																				
<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="background-color: #1a3d4d; color: white;">GENERIC</th> <th style="background-color: #1a3d4d; color: white;">TRADE NAME</th> </tr> </thead> <tbody> <tr> <td>amitriptyline</td> <td>—</td> </tr> <tr> <td>amoxapine</td> <td>—</td> </tr> <tr> <td>clomipramine</td> <td>Anafranil</td> </tr> <tr> <td>protriptyline</td> <td>Vivactil</td> </tr> <tr> <td>nortriptyline</td> <td>Pamelor</td> </tr> </tbody> </table> <p style="background-color: #1a3d4d; color: white; padding: 5px; text-align: center; font-weight: bold;">SUFFIXES: -TRIPTYLINE, -PRAMINE</p>	GENERIC	TRADE NAME	amitriptyline	—	amoxapine	—	clomipramine	Anafranil	protriptyline	Vivactil	nortriptyline	Pamelor	<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="background-color: #1a3d4d; color: white;">GENERIC</th> <th style="background-color: #1a3d4d; color: white;">TRADE NAME</th> </tr> </thead> <tbody> <tr> <td>phenelzine</td> <td>Nardil</td> </tr> <tr> <td>tranylcypromine</td> <td>Parnate</td> </tr> <tr> <td>isocarboxazid</td> <td>Marplan</td> </tr> </tbody> </table>	GENERIC	TRADE NAME	phenelzine	Nardil	tranylcypromine	Parnate	isocarboxazid	Marplan	<p>DRUG TABLE</p>
GENERIC	TRADE NAME																					
amitriptyline	—																					
amoxapine	—																					
clomipramine	Anafranil																					
protriptyline	Vivactil																					
nortriptyline	Pamelor																					
GENERIC	TRADE NAME																					
phenelzine	Nardil																					
tranylcypromine	Parnate																					
isocarboxazid	Marplan																					

ANTI-ANXIETY DRUGS (ANXIOLYTICS)

BENZODIAZEPINES



USES

Bipolar disorder

Benzos are mainly prescribed for:

- acute anxiety
- sedation/muscle relaxant
- seizures
- alcohol withdrawal

Not a first-line drug for treating long-term psychiatric anxiety conditions

ACTION

Binds to cell receptors enhancing the effects of **GABA**

GABA (inhibitory neurotransmitter) slows/calms the activity of the nerves in the brain



GENERIC	TRADE NAME
alprazolam	Xanax
lorazepam	Ativan
diazepam	Valium
clonazepam	Klonopin
chlordiazepoxide	Librium

SUFFIXES: -ZOLAM, -ZEPAM

ANTIDOTE: FLUMAZENIL



I **FLU** fast in my Mercedes **BENZ**



ADVERSE DRUG REACTIONS (ADRs)

- Mild drowsiness, sedation
- Lightheadedness, dizziness, ataxia
- Visual disturbances
- Anger, restlessness
- Nausea, constipation, diarrhea
- Lethargy, apathy, fatigue
- Dry mouth

NURSING CONSIDERATIONS TO HELP WITH ADRs

Take at night if it makes you dizzy/drowsy
Rise slowly from sitting or lying
Do not drive or operate heavy machinery

Fluids, fiber, & exercise!
Give with food to ↓ GI upset

Sips of water, suck on hard candy, chewing sugar-free gum

SYMPTOMS OF WITHDRAWAL

Withdrawals typically happen when the medication is stopped abruptly or taken for >3 months

- ↑ Anxiety
- ↑ HR
- ↑ BP
- ↑ Temp/sweating
- ↓ Memory
- Agitation
- Seizures/tremors
- Insomnia
- Vomiting
- Muscle aches

NURSING CONSIDERATIONS

- Not meant for long term therapy because ↑ risk for physical & psychological **DEPENDENCE**
- Use of long term therapy leads to **TOLERANCE**
Larger doses of the drug are required to achieve the desired outcome
- Must be **TAPERED**
↓ the dosage gradually.
NEVER stop the medication abruptly!

CONTRAINDICATIONS & PRECAUTIONS

- Pregnant, laboring & lactating women
- Elderly (↑ chance of dementia)
- Impaired liver or kidney function
- Debilitation



NONBENZODIAZEPINES

ACTION

Depends on the drug

bupirone (Buspar)
acts on serotonin receptors

hydroxyzine (Vistaril)
acts on the hypothalamus & brainstem reticular formation

GENERIC	TRADE NAME
bupirone	Buspar
doxepin	Silenor
hydroxyzine	Vistaril
meprobamate	-

ANTIPSYCHOTICS

Most commonly used for psychosis (schizophrenia)



REVIEW: WHY ARE SGAs BETTER THAN FGAs?

SGAs work on both positive & negative symptoms, and have a lower risk of developing tardive dyskinesia (TD)

FIRST GENERATION ANTIPSYCHOTICS (FGAs)

Also called **TYPICAL/CONVENTIONAL**

GENERIC	TRADE NAME
chlorpromazine	-
haloperidol	Haldol
loxapine	Adasuve

SECOND GENERATION ANTIPSYCHOTICS (SGAs)

Also called **ATYPICAL**

GENERIC	TRADE NAME
risperidone	Risperdal
clozapine	Clozaril
quetiapine	Seroquel
ziprasidone	Geodon
aripiprazole	Abilify

ACTIONS

- Blocks/inhibits dopamine from being released in the brain
- Helps diminish positive symptoms of schizophrenia

ACTIONS

- Acts on both serotonin & dopamine in the brain
- Helps diminish positive symptoms of schizophrenia & helps negative symptoms as well!

SIDE EFFECTS

- Higher risk of **TD, EPS, & NMS**
- Orthostatic hypotension

SIDE EFFECTS OF BOTH

- Anticholinergic effects
- Photophobia
- Photosensitivity
- Sedation/lethargy

SIDE EFFECTS

- Lower risk of TD, EPS & NMS
- ↑ Weight
- ↑ Cholesterol
- ↑ Triglyceride
- ↑ Blood sugar

TARDIVE DYSKINESIA (TD)

- Involuntary movements of the face, tongue, or limbs that may be irreversible.

EXTRAPYRAMIDAL SYNDROME (EPS)

- Parkinson's like symptoms • Akathisia (restlessness) • Dystonia (muscle twitching)

NEUROLEPTIC MALIGNANT SYNDROME (NMS)

- Combination of symptoms: EPS, high fever, & autonomic disturbance
- One can recover 7-10 days after DC of medication, but it can be fatal if not treated in time

CONTRAINDICATIONS

- Hypersensitivity
- Comatose client
- Depressed
- Bone marrow depression
- Blood dyscrasias
- Parkinson's disease
- Liver problems
- Coronary artery disease
- **Hyper** or **hypotension**

NURSING CONSIDERATIONS



Educate that it may take 6 - 10 weeks to take effect



Tell client about adverse reactions and emphasize that adherence is very important

FGAs

- Teach S&S of **TD, EPDS, & NMS!**
- Advise the client to get up slowly

SGAs

- Check labs (blood sugar, LDL, triglycerides)
- To decrease the risk of gaining weight, advise the client about exercise, low-calorie diet, & monitor their weight.

LEVOthyroxine

GENERIC	TRADE NAME
levo thy roxine	Synth roid



think **THY**roid

SYNthetic **THY**roid

ACTION

- The exact mechanisms are not fully known
- Levothyroxine increases the metabolic rate of tissues


SIDE EFFECTS

- Anxiety
- GI upset
- Sweating
- Weight loss
- Heat intolerance

**SAME AS
HYPERTHYROIDISM!**

MEDICATION CLASS

- Synthetic Hormone



THERAPEUTIC USES

- Treats hypothyroidism
- Thyroid-stimulating hormone suppression
- Thyroid diagnostic testing
- Hormone supplement after thyroidectomy

Should not be used as a weight loss regimen

THERAPEUTIC RESPONSE

NO LONGER SHOWING SIGNS OF HYPOTHYROIDISM

- Normal heart rate (60 - 100 BPM)
- Improved energy levels (not fatigued)
- Normal skin (not cool or pale)

SAFE DURING PREGNANCY? YES!

PATIENT EDUCATION



- It may take 8 weeks to see the full effect
- Report signs of **HYPERTHYROIDISM** ⚠️
 - Tachycardia, heart palpitations, weight loss, insomnia, anxiety
- Monitor T4 & T3 levels
- Take once a day (in the morning before breakfast)
- Take at the same time everyday
- Take on an empty stomach

Educate on the importance of compliance!

Do not stop the medication if symptoms resolve. Thyroid hormone is needed for fetal brain development!



Levothyroxine is a **Life Long** therapy

ANTITHYROID DRUGS

METHIMAZOLE

GENERIC	TRADE NAME
methimazole	Tapazole

MEDICATION CLASS

First-line antithyroid drugs

ACTION

- Inhibits the manufacture of thyroid hormones
- Does not affect existing thyroid hormones circulating in the blood or stored in the thyroid gland

SIDE EFFECTS

- Hay fever
- Skin rash
- Headache
- Nausea & vomiting
- Paresthesias



SYSTEMIC ADVERSE REACTIONS

Risk for:

- Agranulocytosis
- Drug-induced hepatitis



Monitor liver values



PROPYLTHIOURACIL (PTU)

GENERIC
propylthiouracil (PTU)



Prevents Thyroid from being Up

MEDICATION CLASS

First-line antithyroid drugs

USES

- Treats hyperthyroidism
- Treats thyrotoxicosis
- Treats Graves' disease (autoimmune disease that causes hyperthyroidism)
- Used before thyroidectomy surgery (shrinking it before the surgery)



PATIENT EDUCATION

- It may take 1-2 weeks to see the full effect
- Report signs of **HYPOTHYROIDISM** (Bradycardia, weight gain, lethargy, cold intolerance, depression)
- Report signs & symptoms of an infection to the health care provider (Fever, sore throat, etc.)
- Do not abruptly stop the medication (could cause **THYROID STORM**)

Educate on the importance of compliance



PREGNANCY CONSIDERATIONS

- Use with extreme caution during pregnancy because they can cause **hypothyroidism** in the fetus
- If it's necessary, **PROPYLTHIOURACIL** is the preferred drug (does not cross the placenta)



REMEMBER: The fetus needs thyroid hormone for proper brain development



INSULIN TYPES

RAPID



GENERIC

Lispro
Aspart
Glulisine

BRAND NAME

Humalog
Novolog
Apidra

ONSET: 5 - 30 min
PEAK: 30 - 90 min
DURATION: 3 - 5 hrs

HIGHEST RISK FOR HYPOGLYCEMIA

SHORT



Regular

Humulin R
Novolin R

ONSET: 30 - 60 min
PEAK: 2 - 4 hrs
DURATION: 5 - 7 hrs

ONLY INSULIN GIVEN IV



Regular goes **R**ight into the vein

INTERMEDIATE



NPH

Humulin N
Novolin N

ONSET: 1 - 2 hrs
PEAK: 4 - 12 hrs
DURATION: 18 - 24 hrs

NEVER GIVE IV

LONG



Long think
Lonely

Glargine
Detemir


Lantus
Levemir

ONSET: 1 - 2 hrs
PEAK: None
DURATION: 24 hrs+

LOWEST RISK FOR HYPOGLYCEMIA

DO NOT MIX WITH ANY OTHER INSULIN

MIXING REGULAR INSULIN & NPH INSULIN



how to remember this order?

"You are **Not Retired**, you are an **RN**"

ADMINISTRATION

- Must be given subcut or IV
 - Insulin is destroyed by the GI tract so it can not be given PO
- Remove all air bubbles
- Rotate site 1 inch from previous site
- Common sites: back of arms, thighs & abdomen (at least 2 inches away from the belly button)



COMPLICATIONS

- Hypoglycemia (especially with rapid insulin)
- Weight gain
 - Insulin is a growth hormone
- Lipoatrophy (loss of subcut fat)

ALLOPURINOL

GENERIC	TRADE NAME
allopurinol	Aloprim, Zyloprim, Lopurin

MEDICATION CLASS

Uric acid inhibitors

THERAPEUTIC USES

PREVENTS gout attacks ⭐

Does not help with acute attacks



AlloPurinol → P prevents gout

Take NSAIDs for acute attacks, NOT aspirin

SIDE EFFECTS

GI UPSET:
Nausea, vomiting, abdominal pain, diarrhea

SKIN RASH

EDUCATION

Stop the medication if a **RASH** occurs
 ▪ This may indicate a **HYPERSENSITIVITY REACTION** ☠️
 (Stevens-Johnson syndrome)

G Gulp a lot of fluid during the day (2-3 L/day) & take the medication with a glass of water

O No **O**rgan meats

U Urine output up to 2 L/day

T Takes several months to take effect

- Uric acid deposits can cause kidney stones
 - Fluids help prevent this
- Allopurinol + aspirin = ↑ uric acid levels
 - Take acetaminophen instead

COLCHICINE

GENERIC	TRADE NAME
colchicine	Mitigare, Colcris

MEDICATION CLASS

Antigout agent

THERAPEUTIC USES

RELIEVES acute gout attacks ⭐

Also **PREVENTS** gout attacks as well

- Does not help with pain relief, only helps decrease inflammation



Colchicine → for a Cute gout attacks

Take NSAIDs for acute attacks, NOT aspirin

SIDE EFFECTS

GI UPSET:
Nausea, vomiting, abdominal pain, diarrhea

ADVERSE REACTION:
Risk for **BONE MARROW SUPPRESSION** ☠️

EDUCATION

G Gulp a lot of fluid during the day (2-3 L/day) & take the medication with a glass of water

O No **O**rgan meats

U Urine output up to 2 L/day

T Takes several months to take effect

- Uric acid deposits can cause kidney stones
 - Fluids help prevent this

same GI side effects

BISPHOSPHONATES VS. CALCITONIN (SALMON)

BISPHOSPHONATES

GENERIC	TRADE NAME
alendronate	Binosto
etidronate	Didronel
ibandronate	Boniva
pamidronate	Aredia
risedronate	Actonel

SUFFIX: -DRONATE

MEDICATION CLASS

☞ Bone resorption inhibitors

MODE OF ACTION

☞ Bisphosphonates inhibit normal & abnormal **BONE RESORPTION** which leads to increased bone mineral density!

THERAPEUTIC USES

☞ **BUILDS BONE DENSITY & PREVENTS BONE FRACTURES**


- ☞ Treats & prevents osteoporosis (postmenopausal & long term use of steroids)
- ☞ Treats paget's disease
- ☞ Treats hypercalcemia

SIDE EFFECTS

☞ **GI UPSET:**
Nausea, diarrhea, dyspepsia, acid reflux, abdominal pain

EDUCATION

- ☞ Take with a full glass of water on an empty stomach
 - ☞ Stay upright for 30 minutes
 - ☞ Separate iron, antacids, & multiple vitamins at least 30 minutes apart from taking bisphosphonates
 - ☞ Encourage increased intake of calcium & vitamin D
 - ☞ Encourage weight-bearing exercises to preserve bone mass
- "If you don't use it, you lose it!"**



These decrease absorption



NURSING CONSIDERATIONS

☞ Monitor serum calcium levels before, during, & after therapy

NORMAL CALCIUM:
9 - 11 mg/dL

CALCITONIN (SALMON)

GENERIC	TRADE NAME
calcitonin (salmon)	Miacalcin

MEDICATION CLASS

- ☞ Hormone
- ☞ Hypocalcemic agent



Calci**TON**in helps **TON**e down calcium levels in the blood!

MODE OF ACTION

- ☞ Inhibits **OSTEOCLASTS** (Cells that cause bone breakdown)
- ☞ ↓ the rate of bone breakdown

THERAPEUTIC USES

- ☞ Treats & prevents postmenopausal **OSTEOPOROSIS**
- ☞ Treats hypercalcemia
 - ☐ Too much calcium in the bloodstream (we want it in the bones, not in the bloodstream)



SIDE EFFECTS

- ☞ **GI UPSET**
- ☞ **INTRANASAL ROUTE**
Nasal irritation & nasal dryness



EDUCATION

- ☞ Encourage increased intake of calcium & vitamin D
 - ☞ Encourage weight-bearing exercises to preserve bone mass
- "If you don't use it, you lose it!"**




TEMPLATES & PLANNERS

Tear these pages out and make copies,
as many as you need!

BROUGHT TO YOU BY





HOURLY Planner

MONTH: _____

- SUNDAY
 MONDAY
 TUESDAY
 WEDNESDAY
 THURSDAY
 FRIDAY
 SATURDAY

PRIORITIES	PRIORITIES	PRIORITIES	PRIORITIES	PRIORITIES	PRIORITIES	PRIORITIES
6 AM	6 AM	6 AM	6 AM	6 AM	6 AM	6 AM
7 AM	7 AM	7 AM	7 AM	7 AM	7 AM	7 AM
8 AM	8 AM	8 AM	8 AM	8 AM	8 AM	8 AM
9 AM	9 AM	9 AM	9 AM	9 AM	9 AM	9 AM
10 AM	10 AM	10 AM	10 AM	10 AM	10 AM	10 AM
11 AM	11 AM	11 AM	11 AM	11 AM	11 AM	11 AM
12 PM	12 PM	12 PM	12 PM	12 PM	12 PM	12 PM
1 PM	1 PM	1 PM	1 PM	1 PM	1 PM	1 PM
2 PM	2 PM	2 PM	2 PM	2 PM	2 PM	2 PM
3 PM	3 PM	3 PM	3 PM	3 PM	3 PM	3 PM
4 PM	4 PM	4 PM	4 PM	4 PM	4 PM	4 PM
5 PM	5 PM	5 PM	5 PM	5 PM	5 PM	5 PM
6 PM	6 PM	6 PM	6 PM	6 PM	6 PM	6 PM
7 PM	7 PM	7 PM	7 PM	7 PM	7 PM	7 PM
8 PM	8 PM	8 PM	8 PM	8 PM	8 PM	8 PM
9 PM	9 PM	9 PM	9 PM	9 PM	9 PM	9 PM

PRODUCTIVITY METER

BAD
 GOOD
 GREAT!

PRODUCTIVITY METER

BAD
 GOOD
 GREAT!

PRODUCTIVITY METER

BAD
 GOOD
 GREAT!

PRODUCTIVITY METER

BAD
 GOOD
 GREAT!

PRODUCTIVITY METER

BAD
 GOOD
 GREAT!

PRODUCTIVITY METER

BAD
 GOOD
 GREAT!

MONTHLY *Planner*

MONTH: _____

YEAR: _____

WEEKLY Planner

MONDAY:

TUESDAY:

WEDNESDAY:

THURSDAY:

FRIDAY:

SATURDAY:

SUNDAY:

TO DO LIST

- _____
- _____
- _____
- _____
- _____
- _____
- _____
- _____
- _____
- _____
- _____
- _____
- _____
- _____
- _____

NOTES

TESTS / EXAMS

PROJECTS / ASSIGNMENTS

SELF-CARE ♥

DISEASE:

PATHOLOGY

SIGNS & SYMPTOMS

RISK FACTORS

COMPLICATIONS

DIAGNOSIS

TREATMENT

PHARMACOLOGY TEMPLATE

DRUG CLASS: _____

GENERIC NAME	TRADE NAME
SUFFIXES OR PREFIXES:	
ANTIDOTE:	

ACTION

THERAPEUTIC USES

CONTRAINDICATIONS

SIDE EFFECTS

NURSING CONSIDERATIONS

PATIENT EDUCATION



NCLEX Study Schedule

WORLD EDITION

Month: _____
NCLEX Date: _____

SUNDAY

Subject:
Body System:
of Practice Questions:
Self Care:

MONDAY

Subject:
Body System:
of Practice Questions:
Self Care:

TUESDAY

Subject:
Body System:
of Practice Questions:
Self Care:

WEDNESDAY

Subject:
Body System:
of Practice Questions:
Self Care:

THURSDAY

Subject:
Body System:
of Practice Questions:
Self Care:

FRIDAY

Subject:
Body System:
of Practice Questions:
Self Care:

SATURDAY

Subject:
Body System:
of Practice Questions:
Self Care:

Subject:
Body System:
of Practice Questions:
Self Care:

Subject:
Body System:
of Practice Questions:
Self Care:

Subject:
Body System:
of Practice Questions:
Self Care:

Subject:
Body System:
of Practice Questions:
Self Care:

Subject:
Body System:
of Practice Questions:
Self Care:

Subject:
Body System:
of Practice Questions:
Self Care:

Subject:
Body System:
of Practice Questions:
Self Care:

Subject:
Body System:
of Practice Questions:
Self Care:

Subject:
Body System:
of Practice Questions:
Self Care:

Subject:
Body System:
of Practice Questions:
Self Care:

Subject:
Body System:
of Practice Questions:
Self Care:

Subject:
Body System:
of Practice Questions:
Self Care:

Subject:
Body System:
of Practice Questions:
Self Care:

Subject:
Body System:
of Practice Questions:
Self Care:

Subject:
Body System:
of Practice Questions:
Self Care:

Subject:
Body System:
of Practice Questions:
Self Care:

Subject:
Body System:
of Practice Questions:
Self Care:

Subject:
Body System:
of Practice Questions:
Self Care:

Subject:
Body System:
of Practice Questions:
Self Care:

Subject:
Body System:
of Practice Questions:
Self Care:

Subject:
Body System:
of Practice Questions:
Self Care:

Subject:
Body System:
of Practice Questions:
Self Care:

Subject:
Body System:
of Practice Questions:
Self Care:

Subject:
Body System:
of Practice Questions:
Self Care:

Subject:
Body System:
of Practice Questions:
Self Care:

Subject:
Body System:
of Practice Questions:
Self Care:

Subject:
Body System:
of Practice Questions:
Self Care:

Subject:
Body System:
of Practice Questions:
Self Care:

Dear future nurse,

You may be stressed, you may feel tired, and you may want to give up. Nursing school is hard, there's no doubt about it. Everyone cries, everyone has meltdowns, and there will be moments you don't feel qualified for the task at hand. But take heart, the challenge only makes you stronger. Put in the work, show up on time, and find an amazing study group. You got this! 😊

— Kristine Tuttle, BSN, RN



You
got this,
future
nurse!

By purchasing this material, you agree to the following terms and conditions: you agree that this ebook and all other media produced by NurseInTheMaking LLC are simply guides and should not be used over and above your course material and teacher instruction in nursing school. When details contained within these guides and other media differ, you will defer to your nursing school's faculty/staff instruction. Hospitals and universities may differ on lab values; you will defer to your hospital or nursing school's faculty/staff instruction. These guides and other media created by NurseInTheMaking LLC are not intended to be used as medical advice or clinical practice; they are for educational use only. You also agree to not distribute or share these materials under any circumstances; they are for personal use only.

© 2021 NurseInTheMaking LLC. All content is property of NurseInTheMaking LLC and www.anurseinthemaking.com. Replication and distribution of this material is prohibited by law. All digital products (PDF files, ebooks, resources, and all online content) are subject to copyright protection. **Each product sold is licensed to an individual user and customers are not allowed to distribute, copy, share, or transfer the products to any other individual or entity, they are for personal use only. Fines of up to \$10,000 may apply and individuals will be reported to the BRN and their school of nursing.**



www.anurseinthemaking.com

Need more
help with
nursing school?



SCAN ME!